



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2018

Microbiome and asthma

Sokolowska, Milena ; Frei, Remo ; Lunjani, Nonhlanhla ; Akdis, Cezmi A ; O'Mahony, Liam

Abstract: The mucosal immune system is in constant communication with the vast diversity of microbes present on body surfaces. The discovery of novel molecular mechanisms, which mediate host-microbe communication, have highlighted the important roles played by microbes in influencing mucosal immune responses. Dendritic cells, epithelial cells, ILCs, T regulatory cells, effector lymphocytes, NKT cells and B cells can all be influenced by the microbiome. Many of the mechanisms being described are bacterial strain- or metabolite-specific. Microbial dysbiosis in the gut and the lung is increasingly being associated with the incidence and severity of asthma. More accurate endotyping of patients with asthma may be assisted by further analysis of the composition and metabolic activity of an individual's microbiome. In addition, the efficacy of specific therapeutics may be influenced by the microbiome and novel bacterial-based therapeutics should be considered in future clinical studies.

DOI: <https://doi.org/10.1186/s40733-017-0037-y>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-148435>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Sokolowska, Milena; Frei, Remo; Lunjani, Nonhlanhla; Akdis, Cezmi A; O'Mahony, Liam (2018). Microbiome and asthma. *Asthma Research and Practice*, 4:1.

DOI: <https://doi.org/10.1186/s40733-017-0037-y>

REVIEW

Open Access



Microbiome and asthma

Milena Sokolowska^{1,2}, Remo Frei^{1,2}, Nonhlanhla Lunjani^{1,2,3}, Cezmi A. Akdis^{1,2} and Liam O'Mahony^{1*}

Abstract

The mucosal immune system is in constant communication with the vast diversity of microbes present on body surfaces. The discovery of novel molecular mechanisms, which mediate host-microbe communication, have highlighted the important roles played by microbes in influencing mucosal immune responses. Dendritic cells, epithelial cells, ILCs, T regulatory cells, effector lymphocytes, NKT cells and B cells can all be influenced by the microbiome. Many of the mechanisms being described are bacterial strain- or metabolite-specific. Microbial dysbiosis in the gut and the lung is increasingly being associated with the incidence and severity of asthma. More accurate endotyping of patients with asthma may be assisted by further analysis of the composition and metabolic activity of an individual's microbiome. In addition, the efficacy of specific therapeutics may be influenced by the microbiome and novel bacterial-based therapeutics should be considered in future clinical studies.

Keywords: Asthma, Microbiome, Bacteria, Mucosal immune system, Immune tolerance, Short-chain fatty acids, Histamine

Background

An enormous number of microbes colonize the skin and mucosal body surfaces. These microbes are highly adapted to survive within complex community structures, utilizing nutrients from other microbes and/or host processes. The microbiome is defined as the sum of these microbes, their genomic elements and interactions in a given ecological niche. The composition and diversity of the microbiome varies across body sites, resulting in a series of unique habitats within and between individuals that can change substantially over time [1]. The establishment of stable microbial communities closely tracks host growth and immune development during the first few years of life. Factors that influence this evolution include antibiotic use, birth mode, infant nutrition and biodiversity in the home, surrounding environment and in family members [2]. Delayed or altered establishment of these microbial communities' leads to microbiome immaturity and has been associated with increased risk of allergies and asthma later in life.

Highly sophisticated mucosal immune cellular and molecular networks need to be constantly coordinated in order to tolerate the presence of a large number and

diversity of bacteria, while protective immune responses to potential pathogens must be maintained and induced on demand. The balance between immune tolerance and inflammation within tissues is regulated in part by the crosstalk between immune cells and the microbiome [3]. Disrupted communication between the microbiome and the host due to altered microbiome composition and/or metabolism is thought to negatively influence immune homeostatic networks. This can be clearly seen in mice bred under germ-free (GF) or sterile conditions, whereby mucosal tolerance mechanisms do not fully develop and these mice display increased allergic responses to allergen challenge.

In this review, we will examine the potential mechanisms by which the microbiome influences immune responses within the lung and assess the evidence for a dysbiotic microbiome in the gut and the respiratory tract of asthma patients. In addition, we will summarize the current therapeutic approaches and challenges associated with microbial-based therapies in asthma patients and highlight the future research and clinical needs in the field.

Immune mechanisms influenced by the microbiome

Multiple mechanisms have now been described, through which bacteria can induce regulatory responses or dampen inflammatory processes. Both bacterial cell wall components and metabolites from the microbiome have

* Correspondence: liam.omahony@siaf.uzh.ch

¹Swiss Institute of Allergy and Asthma Research, University of Zürich, Obere Strasse 22, 7270 Davos, Switzerland

Full list of author information is available at the end of the article

been associated with immunoregulatory effects within the mucosa. Certain commensal microbes such as specific *Bifidobacterium*, *Lactobacillus* and *Clostridium* strains have been shown to increase the proportion of T regulatory cells in mice [4–8]. In addition, *Clostridia* have been shown to stimulate ILC3s to produce IL-22, which helps to reinforce the epithelial barrier and reduces the permeability of the intestine to dietary proteins [9]. Furthermore, *Bifidobacteria* and *Lactobacilli* can stimulate metabolic processes in dendritic cells, such as vitamin A metabolism, tryptophan metabolism and heme oxygenase-1, which promote induction of T regulatory cells [10–12]. The capsular polysaccharide A from *Bacteroides fragilis* has been shown to interact directly with mouse plasmacytoid dendritic cells and thereby promoted IL-10 secretion from CD4⁺ T cells [13]. In addition, an exopolysaccharide from *Bifidobacterium longum* was recently shown to suppress Th17 responses within the gut and within the lung [14, 15]. Notably, consumption of *Bifidobacterium longum* 35,624 by healthy human volunteers increased Foxp3⁺ T regulatory cells in peripheral blood, while administration of this bacterial strain to psoriasis patients, chronic fatigue syndrome patients or ulcerative colitis patients consistently resulted in reduced levels of serum proinflammatory biomarkers such as CRP, possibly mediated by increased numbers of T regulatory cells [12, 16].

In addition to bacterial-associated components, bacterial-derived metabolites have significant effects on immunoregulatory processes. Short-chain fatty acids (SCFAs), such as acetate, propionate and butyrate, are produced by the gut microbiota and have been shown to influence dendritic cell and T cell responses, via their binding to G protein-coupled receptors and their inhibition of histone deacetylases, thereby promoting epigenetic changes [17]. Bacteria within the human gut can produce a wide range of biogenic amines (due to metabolism of amino acids), which can also influence immune and inflammatory responses [18]. Interestingly, in murine models, microbiota-derived taurine, histamine, and spermine were shown to influence host-microbiome interactions by co-modulating NLRP6 inflammasome signaling, epithelial IL-18 secretion, and downstream anti-microbial peptide secretion [19].

Microbiome in animal models of asthma

A number of different animal studies support the concept for a role of the microbiome in development of airway diseases. In particular, valuable insights for the mechanistic role of the microbiome in the development of allergic airway inflammation comes from GF animals, lacking any exposure to pathogenic or nonpathogenic microorganisms. Herbst et al. observed that OVA-induced type 2 airway inflammation and airway

hypersensitivity is much stronger in GF mice as compared to the mice from a specific pathogen-free environment (SPF) that were colonized with commensal microbes. Moreover, the exaggerated allergic inflammation in GF mice could be reduced to the same level observed in SPF mice, when GF mice were co-housed for 3 weeks with SPF mice, suggesting that gut and airways recolonization with commensal microbes had protective effects [20]. In addition, early life colonization of GF mice prevented invariant natural killer T cell accumulation in the gut lamina propria and the lungs thereby reducing the severity of allergic airway responses. Later life colonization had no effect on disease phenotypes nor on the development of regulatory T cells or on invariant natural killer T cells [21]. Furthermore, antibiotic treatment of neonatal mice resulted in fewer regulatory T cells and a more pronounced T helper cell type 2 response, which was prevented by re-introducing a commensal intestinal microbiota [22–24].

Gollwitzer et al. examined the susceptibility to house dust mite (HDM)-induced allergic airway inflammation in mice of different ages (3, 15 and 60 days), simulating the conditions of gradual colonisation of the human infant airways [25]. Neonatal mice were prone to develop exaggerated airway eosinophilia, they released more type 2 cytokines and exhibited higher airway hyper-responsiveness following exposure to HDM compared to mice that were older. This protective effect in older mice was associated with the colonization of the mouse lungs with increased numbers of bacteria and the shift from a predominance of Gammaproteobacteria and Firmicutes to Bacteroidetes. The maturation of the lung microbiota was associated with the PDL-1-dependent emergence of Helios-negative T regulatory cells. This study suggests that the absence of specific bacterial species early in life could influence appropriate regulatory mechanisms later in life and subsequently shift the immunological balance towards allergy instead of tolerance [25].

Oral supplementation of mice with specific microbes such as *Bifidobacterium breve*, *Clostridium* clade IV and XIV species, or with the capsular polysaccharide PSA of *Bacteroides fragilis* induced an anti-inflammatory response associated with induction of regulatory T cells and IL-10 secretion that attenuated allergic airway inflammation [21, 26]. In addition to the gut-induced regulatory T cells that could migrate to the lungs to provide anti-inflammatory effects, there are metabolites produced by the microbiome such as SCFAs that are absorbed and potentially have direct effects on lung immune responses [27]. Deliberate administration of SCFAs, or dietary fibers that are metabolized to SCFAs, has repeatedly been shown to reduce airway inflammation in murine models. A high-fiber diet increased the

level of colonic Bacteroidetes and Actinobacteria species and decreased Firmicutes and Proteobacteria, which was associated with increased SCFA serum levels and suppression of allergic airway-inflammation in mice [28]. The beneficial effect was transferred to the offspring after treatment of pregnant mice via epigenetic mechanisms [26, 29]. The influence of the microbiota on D-tryptophan, Vitamin A or biogenic amine metabolism can also modulate T helper cell type 2 mediated allergic airway inflammation within the lung [3, 26, 30, 31].

Several studies have suggested that direct exposure of the murine respiratory tract to microbial products such as endotoxin, CpG-containing oligonucleotides or other Toll-like receptor ligands could inhibit the classical features of asthma [32, 33]. For example, intranasal exposure to the bacterium *Escherichia coli* was protective in the OVA-induced allergic airway inflammation model [34]. These studies were recently expanded by novel findings that linked the protective effect of the farm environment with the microbiota and endotoxin levels in the house dust. Schuijs et al. demonstrated that prolonged exposure to low-dose endotoxin or farm dust protected mice from HDM-induced asthma via A20 (TNFAIP3)-dependent airway epithelial cells-dendritic cells interactions [35]. Stein et al. further demonstrated that intranasal installation of the dust from Amish houses, but not dust from Hutterite homes, reduced OVA-induced allergic airway inflammation in mice, via Myd88 and Trif-dependent mechanisms [36]. The dust from the Amish homes had different bacterial populations (especially higher in Bartonellaceae) and higher endotoxin levels as compared to Hutterite houses' dust [36].

The role of the gut microbiome in asthma

The human gut microbiome is the largest collection of bacteria in the body, consisting of 500–1000 distinct bacterial species with more than 8 million genes potentially influencing the host immune system [21, 37]. European adults' gut microbiota is predominantly colonized by Bacteroidetes, Firmicutes, Actinobacteria, Proteobacteria, and Verrucomicrobia. The stomach, duodenum, and proximal small intestine are mainly colonized with aerobic bacteria including Streptococci species, Lactobacilli species, and Enterobacteriaceae while anaerobes such as Bacteroides, Bifidobacterium, Prevotellaceae, Rikenellaceae, Lachnospiraceae, Ruminococcaceae, and Clostridium species dominate the distal small intestine and the colon [26, 38]. The gut microbiota can influence immune responses at distant sites (such as the lung) via multiple mechanisms. For example, it was recently shown that there is an increase in the number of bacteria capable of secreting histamine from the gut of adult asthma patients, compared to healthy volunteers [39]. However, it is not clear if increased secretion of

histamine by gut microbes can have an overall detrimental or protective effect as histamine can induce protective responses in the lung via histamine 2 receptor and detrimental effects via histamine 1 and 4 receptors [40].

The composition of the gut microbiome is thought to reach an adult-like diversity by 3 years of age. Development of the early life gut microbiome is influenced by many environmental factors, such as living in microbial rich environments (e.g. on a farm or with frequent contact to livestock and pets), or a diverse diet, which have been inversely associated with childhood asthma [41–45]. It is thought that exposure to and colonization by certain microbes at the correct time during early life is important for gut development, immune cell maturation and resistance to pathogens, all of which may protect against the development of asthma [22, 37, 46]. The mode of delivery has a significant influence on colonization. Babies delivered via caesarean section typically have more Staphylococcus species, Bacillales, Propionobacterineae, Corynebacterineae, Firmicutes and Acinetobacter species with fewer Actinobacteria and Bacteroidetes, while vaginal delivery has been linked to increased colonization with Clostridia [38, 47]. Clostridia metabolize fibers to SCFAs, which can have systemic anti-inflammatory effects as described above. In addition to delivery mode and diet, maternal antibiotic use during pregnancy or antibiotic treatment in early childhood significantly disrupts the microbiota and was associated with long-lasting effects such as decreased Actinobacteria and increased Bacteroidetes and Proteobacteria [1].

Several studies have linked early life dysbiosis of the gut microbiota with an altered risk of asthma later in life. Colonization by *Clostridium difficile* at 1 month of age was associated with wheeze throughout the first 6 to 7 years of life and with asthma at age 6 to 7 years [48]. Children that developed asthma at school age, had a lower gut microbiome diversity at 1 week or 1 month of age, but not at 1 year of age, compared to non-asthmatic children [49]. In another study, the early life relative abundance of the bacterial genera Lachnospira, Veillonella, Faecalibacterium, and Rothia was significantly decreased in children at risk of asthma. This dysbiosis was accompanied by reduced levels of fecal acetate and dysregulation of enterohepatic metabolites [50]. In addition, neonates with the lowest relative abundance of Bifidobacteria, Akkermansia and Faecalibacterium and a higher relative abundance of particular fungi (*Candida* and *Rhodotorula*), had the highest risk to develop atopy and asthma [51]. Thus, early life dysbiosis of the gut microbiota has been consistently associated with an increased risk of asthma later in life. However, it remains unclear if microbial dysbiosis within the gut can actually drive relevant disease promoting mechanisms or if dysbiosis simply reflects associated phenomena such as

altered patterns of immune response to microbes and environmental stimuli.

Role of the respiratory microbiome in asthma

The Human Microbiome Project, launched in 2007, did not include airway tissue sampling as healthy human lung tissue at that time was assumed to be sterile [52]. However, shortly afterwards a number of pioneering publications in this field appeared and several research consortia and individual groups subsequently started intensive studies to characterize and understand the composition and function of airway microbiota in health and disease [53–55]. Currently, it is known that the healthy respiratory mucosa is inhabited by niche-specific bacterial communities [56]. The highest densities of bacterial communities are found in the upper respiratory tract, reaching up to 10^3 viable bacteria per nasal swab from the nasal cavity and nasopharynx, with even up to 10^6 /ml viable cells from oropharynx lavages [56–58]. In the trachea and lungs, the estimated numbers of bacteria are lower with approximately 10^2 bacterial cells per ml being found in bronchoalveolar lavages (BAL) from healthy lungs [59]. The six dominant phyla routinely found in the lung are Firmicutes, Proteobacteria, Bacteroides, Fusobacteria, Acidobacteria, and Actinobacteria [60].

The original proof-of-concept study from Hilty et al., with microbiome assessments of the nose, oropharynx, bronchial brushings and BAL samples from the lower airways revealed that the Proteobacteria phylum and especially Haemophilus species are more often present in upper and lower airways of asthmatic and COPD adults and asthmatic children [53]. The study performed by Huang et al. in patients with suboptimal controlled asthma, defined as persistent symptoms on the Asthma Control Questionnaire after 4 weeks of standardized treatment with inhaled fluticasone, showed a greater airway microbiota diversity in these patients compared to control subjects that correlated positively with bronchial hyperresponsiveness [61]. Specifically, there was an increase in the phylum Proteobacteria in asthma patients, including Comamonadaceae, Sphingomonadaceae, Nitrosomonadaceae, Oxalobacteraceae, and Pseudomonadaceae families [61]. Interestingly, adult patients who benefited most from clarithromycin treatment, as assessed by the reduction in bronchial hyperactivity to methacholine were those who had significantly greater bacterial diversity prior the intervention [61]. Subsequent studies also confirmed that Proteobacteria were present in higher proportions in the asthmatic airways [59, 62]. In addition, Klebsiella species were enriched in patients with severe asthma as compared to patients with mild-to-moderate asthma and controls [63]. Moreover, within severe asthma patients, Proteobacteria was associated with T_H17 -related gene signature in airway

epithelium, worsening asthma control and total leukocytes in the sputum, while Bacteroides/Firmicutes were more abundant in obese patients with severe asthma. In contrast, the presence of Actinobacteria correlated with improvement and/or no change in asthma control [63]. Severe asthma had long been associated with the presence of *Mycoplasma pneumoniae* and *Chlamydophila pneumoniae*, resulting in several clinical trials with macrolide antibiotics in this group of patients [64]. Yet, in the face of controversial study results and the possibility that beneficial microbial species are also affected, further clinical trials that include detailed microbiome studies are needed [65, 66].

The composition of the airway microbiome develops exponentially very early in life and later in life can be influenced by the environment, health status and age. Birth mode (vaginal or via caesarean section), the exposures during the first hours of life and the environment of the following 3–4 months of life have been shown to be of utmost importance in shaping the development of stable respiratory and gut microbiota to ensure respiratory health later in life [50, 67–69]. Both human and animal studies have shown that inhaled dust particles can carry a complex mixture of microbes and microbial factors, which influence susceptibility to asthma development via their effects on innate and adaptive immune responses [35, 36]. The important research questions that are currently being addressed in children include: i) what is the longitudinal process of upper airways colonization in healthy infants? ii) how do environmental factors such as breast feeding, living on a farm, number of siblings, day-care, pets at home, smoking and antibiotic usage impact the respiratory microbiome? iii) are there correlations between patterns of respiratory microbial colonization in early life with the occurrence of acute respiratory infections such as respiratory syncytial virus (RSV), rhinovirus and influenza virus and their further impact on chronic non infectious-associated recurrent wheeze, atopic sensitization and asthma? [58, 70–74].

Teo et al. analyzed the nasopharynx microbiome in a prospective cohort of children at several time-points up to 2 years of age and correlated the presence of specific groups of bacteria with acute respiratory infections [68]. Healthy infants from this cohort were initially colonized with Staphylococcus or Corynebacterium species up to 2 months of age with subsequent stable colonization by Allocococcus or Moraxella. In contrast, Streptococcus, Moraxella or Haemophilus colonization were correlated with virus-associated acute respiratory infections in the first 60 weeks of life. Early asymptomatic Streptococcus colonization, rare in children from dog and cat-owning families, increased the risk of asthma at 5 years of age [68]. Early upper respiratory tract colonization with *S. pneumoniae*, *H. influenzae* and/or *M. catarrhalis* in

children at 4 weeks of age from other prospective birth cohorts was also found to be associated with an increased risk of pneumonia and bronchiolitis or asthma at 5 years of age [54, 75]. Additional studies have also noted associations between *H. influenza*, *Streptococcus* species and *S. aureus* nasopharyngeal colonization with RSV infection and hospitalization in children independently of their age [76–78]. Furthermore, early colonization of the upper respiratory tract of healthy infants with *Staphylococcus* species, subsequently followed by *Corynebacterium/Dolosigranulum* and *Moraxella*, were described for infants who were breastfed and who had lower rates of respiratory infections in the first 2 years of life [67, 79, 80]. Indeed, airway microbial diversity appears to be inversely associated with sensitization to house dust mites in early childhood [81, 82]. Of particular interest is a recent study comparing Amish children raised on traditional farms, who have a low prevalence of asthma and atopy, with Hutterite children coming from highly industrialized farms who have a higher prevalence of asthma and atopy, even though these two populations are genetically similar. One striking difference was the microbial composition and endotoxin load of dust from those two housing environments, associated with the enhanced induction of innate immune pathways in Amish children. The high-endotoxin dust from Amish houses was able to inhibit OVA-induced allergic airway inflammation in mice, as described above [36]. Several other studies have also confirmed that the farm environment is associated with increased bacterial diversity in the house dust samples and nasal microbiome diversity of the same children who have lower risk of developing asthma [83–85]. It is currently unknown if the protective effect of the dust-associated microbiome is due to inhalation of multiple bacteria species and further colonization of the airways, or if inhaled bacterial metabolites may also play a role.

Microbiome strategies for asthma prevention, treatment and management

Alterations in the lung and gut microbiome of asthma patients have been well described previously in this review. The deliberate restoration of lung and gut microbiota through the use of prebiotics; probiotics or synbiotics is one potential strategy currently being assessed. Interest in probiotics and prebiotics for their potential benefits in protecting against airway inflammation is relatively recent but increasing significantly, particularly as several lines of evidence suggest that a “healthy” microbial community facilitates the development of immune tolerance [30]. In vitro studies and animal models have repeatedly shown the protective effects of certain probiotic strains on lung inflammatory responses, but have also shown that not all probiotics will induce the

same effects [86]. Intervention and prevention studies in humans are inconsistent in their findings, possibly because many factors complicate the analysis of dysbiosis in patients with asthma. Comparison between human studies are difficult, because of considerable heterogeneity in the probiotics and/or prebiotics used, study design, sample size, age of study population, geographic location and lifestyle factors (including diet). One preliminary study did suggest that symbiotic (pre and probiotic) use improved peak expiratory flow and reduced the systemic production of Th2 cytokines in allergic asthmatics [87]. Another recent study using a combination of nutritional interventions (fish oils and vegetable extracts) with a probiotic led to significant improvement in pulmonary function parameters and significantly reduced requirement for short-acting inhaled bronchodilators and inhaled corticosteroids in children with asthma, suggesting that the combination of multiple approaches may lead to the most optimal benefits [88]. These findings are promising, however more definitive studies are needed to determine whether modification of gut and lung microbiota can be attributed to pre and/or probiotic use. Currently, there is no recommendation to use pre- or probiotics for treatment or prevention of asthma. Nevertheless, there is accumulating evidence that antenatal interactions between maternal diet, gut bacteria and bacterial metabolites may lead to immunological imprinting on the developing fetal immune system that could influence the development of allergy and asthma later in life [89]. Thus, further studies are required to determine if appropriate prebiotic and probiotic use during pregnancy may functionally impact the maternal gut microbiome with subsequent effects on maternal immune function and risk of asthma in the offspring [90].

In addition to using single probiotic bacterial strains, the manipulation of the entire gut microbiome with fecal microbiota transplants (FMT) is currently being explored. FMT has been successfully used for the treatment of *Clostridium difficile* infection and research into its use for other inflammatory diseases such as type 2 diabetes, inflammatory bowel diseases and non-alcoholic steatohepatitis is well under way [91]. The use of FMT beyond intestinal disorders requires additional studies and currently there is no data supporting its use in allergic disease or asthma [1].

The role of the microbiome in influencing precision medicine approaches to patient care has been best explored to date in the oncology field. Accumulating evidence suggests that the microbiome not only influences the severity of treatment-associated side effects in cancer patients, but also has a dramatic effect on treatment efficacy via pharmacodynamic and immunological mechanisms [92]. Notably, a melanoma mouse model showed commensal microbe-derived antitumor immunity evidenced by higher

intratumoral CD8⁺ T cell accumulation. From this microbiota, Bifidobacteria were identified as having the strongest association with antitumor T cell immunity and the ability to maximize the efficacy of the cancer immunotherapeutic anti-PD-L1-specific antibody treatment. Bifidobacteria augmented dendritic cell function leading to enhanced CD8⁺ T cell priming and accumulation within the tumor [93]. While there is a growing amount of data on the compositional differences in lung microbiota in health and disease, there is a dearth of research into the functional role of the microbiome on treatment efficacy in patients with chronic respiratory disorders [94]. One important study did correlate corticosteroid use and corticosteroid sensitivity in asthma patients with the presence of specific microbes in the lower airways. At the genus level, *Neisseria* species, *Haemophilus* species, *Campylobacter* species and *Leptotrichia* species were present in the lower airways of patients with corticosteroid-resistant asthma, but not in patients with corticosteroid-sensitive asthma [59]. Others have demonstrated that corticosteroid use, particularly the combination of inhaled and oral corticosteroids, is associated with an increased abundance of Proteobacteria and the genus *Pseudomonas*, and decreased abundance of Bacteroidetes, Fusobacteria, and *Prevotella* species [60]. One recent study suggests that microbiome-related functions might affect responsiveness to corticosteroid treatment in asthma patients [95]. Pre-steroid treatment *Haemophilus* levels were increased in asthma patients with diminished responses to corticosteroids. Furthermore, the predicted metagenome metabolic functions in inhaled corticosteroid nonresponders suggested increased microbiome-associated xenobiotic degradation capacity [95].

Further profiling and characterization of the microbiome associated with different asthma phenotypes is necessary for identifying novel microbiota-related mechanisms of disease. In addition, identification of these key microbial species and their associated functional effects will contribute to a more precise definition of asthma phenotypes and may help identify more suitable “phenotype-specific” management strategies [96].

Future perspectives

While it is clear that the microbiome significantly influences host immune maturation and immune activity, the molecular basis for these immunomodulatory mechanisms are only beginning to be elucidated. It still remains unclear whether and, if so, to what extent patterns of airway microbial dysbiosis actually drives rather than merely reflects associated patterns of immune reactivity within the lung. Current studies in prospective birth cohorts and cross-sectional studies in children have heightened our awareness of time-sensitive patterns of colonization of seemingly protective or detrimental bacteria in the gut or airways of healthy and diseased

children. However, further mechanistic and epidemiological studies are needed to uncover the functional, multidirectional associations between the specific bacterial strains, host, allergens and viruses. Respiratory microbiome assessments in adults have so far been performed in a cross-sectional manner, comparing the airway microbiota composition between healthy controls and patients with asthma and often with other chronic airway diseases. Some studies have provided detailed clinical characteristics of patients, allowing for the correlation of microbiota differences across different asthma phenotypes. However, longitudinal and prospective analyses of adult airway microbiome in bigger cohorts of well-characterized patients are still needed to understand the relationships between the course of the disease, its phenotype and endotype, susceptibility to exacerbations and disease progression as well as its response to treatment.

Compositional profiling needs to be complemented with metagenomics, transcriptomics, physiological, biochemical and function-oriented analyses of both the host response and microbial communities as interactions between the host and microbiome are almost certainly bidirectional, with species- and strain-specific behaviors shaped by the genetic background and microenvironment in which they exist. In addition, current compositional analysis at the genus level is not sufficient and future analysis needs to be conducted at greater depth to include information at the species and strain level. The immune response to a bacterium is often strain-specific and results from one strain cannot be extrapolated to other strains even within the same species. Thus, the traditional methods for microbiological classification, based on 16 s sequencing and certain biochemical properties, of a bacterium into a given genus or species do not currently help us to predict immunological outcomes. Culturing methods need also to be improved in order to isolate and grow lung-derived bacterial strains *in vitro*, particularly the obligate anaerobes, in order to facilitate strain-specific immunological assessments.

Conclusions

The last few years have resulted in pivotal studies that clearly associate changes in gut or lung microbial populations with asthma. However, mechanistic studies are still necessary to elucidate how members of the microbiota induce or modulate inflammatory responses in asthmatic patients. We anticipate that the continuing identification of novel bacterial strains, their components and metabolites, which modulate mucosal immunoregulatory responses, will open up new possibilities for a bug-to-drug approach in the treatment of asthma patients and the prevention of airway diseases.

Abbreviations

BAL: Bronchoalveolar lavages; FMT: Fecal microbiota transplant; GF: Germ-free; HDM: House dust mite; NLRP6: NOD-like receptor family pyrin domain containing 6; RSV: Respiratory syncytial virus; SCFAs: Short-chain fatty acids; SPF: Specific pathogen-free

Acknowledgements

None

Funding

The authors are supported by Swiss National Science Foundation grants (project numbers CRSII3_154488, 310,030_144219, 310,030-127,356 and 310,030_144219) and Christine Kühne – Center for Allergy Research and Education (CK-CARE).

Availability of data and materials

Not applicable

Authors' contributions

All authors contributed to the writing of the review. All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Competing interests

LOM is a consultant to Alimentary Health Ltd. and has received research funding from GlaxoSmithKline. CA has received research support from Novartis and Stallergenes and consulted for Actellion, Aventis and Allergopharma. MS, RF and NL have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Swiss Institute of Allergy and Asthma Research, University of Zürich, Obere Strasse 22, 7270 Davos, Switzerland. ²Christine Kühne – Center for Allergy Research and Education (CK-CARE), Davos, Switzerland. ³University of Cape Town, Cape Town, South Africa.

Received: 14 September 2017 Accepted: 18 December 2017

Published online: 05 January 2018

References

- Huang Y, Marsland B, Bunyavanich S, O'Mahony L, Leung D, Muraro A, et al. (2017). The microbiome in allergic disease: current understanding and future opportunities—2017 PRACTALL document of the American Academy of Allergy, Asthma & Immunology and the European academy of allergy and clinical immunology. *J Allergy Clin Immunol*. 2017;139:1099–110.
- Bokulich N, Chung J, Battaglia T, Henderson N, Jay M, Li H, et al. Antibiotics, birth mode, and diet shape microbiome maturation during early life. *Sci Transl Med* 2016;8:343–43.
- Frei R, Lauener R, Cramer R, O'Mahony L. Microbiota and dietary interactions - an update to the hygiene hypothesis? *Allergy*. 2012;67:451–61.
- Atarashi K, Tanoue T, Oshima K, Suda W, Nagano Y, Nishikawa H, et al. Treg induction by a rationally selected mixture of clostridia strains from the human microbiota. *Nature*. 2013;500:232–6.
- O'Mahony C, Scully P, O'Mahony D, Murphy S, O'Brien F, Lyons A, et al. Commensal-induced regulatory T cells mediate protection against pathogen-stimulated NF-kappaB activation. *PLoS Pathog*. 2008;4:e1000112.
- Di Giacinto C, Marinaro M, Sanchez M, Strober W, Boirivant M. Probiotics ameliorate recurrent Th1-mediated murine colitis by inducing IL-10 and IL-10- dependent TGF-β-bearing regulatory cells. *J Immunol*. 2005;174:3237–46.
- Karimi K, Inman MD, Bienenstock J, Forsythe P. Lactobacillus reuteri-induced regulatory T cells protect against an allergic airway response in mice. *Am J Respir Crit Care Med*. 2009;179:186–93.
- Tang C, Kamiya T, Liu Y, Kadoki M, Kakuta S, Oshima K, et al. Inhibition of Dectin-1 signaling ameliorates colitis by inducing lactobacillus-mediated regulatory T cell expansion in the intestine. *Cell Host Microbe*. 2015;18:183–97.
- Stefka AT, Feehley T, Tripathi P, Qiu J, McCoy K, Mazmanian SK, et al. Commensal bacteria protect against food allergen sensitization. *Proc Natl Acad Sci U S A*. 2014;111:13145–50.
- Konieczna P, Ferstl R, Ziegler M, Frei R, Nehrbass D, Lauener RP, et al. Immunomodulation by Bifidobacterium infantis 35624 in the murine lamina propria requires retinoic acid-dependent and independent mechanisms. *PLoS One*. 2013;8:e62617.
- Karimi K, Kandiah N, Chau J, Bienenstock J, Forsythe PA. Lactobacillus rhamnosus strain induces a heme oxygenase dependent increase in Foxp3 + regulatory T cells. *PLoS One*. 2012;7:e47556.
- Konieczna P, Groeger D, Ziegler M, Frei R, Ferstl R, Shanahan F, et al. Bifidobacterium infantis35624 administration induces Foxp3 T regulatory cells in human peripheral blood: potential role for myeloid and plasmacytoid dendritic cells. *Gut*. 2011;61:354–66.
- Dasgupta S, Erturk-Hasdemir D, Ochoa-Reparaz J, Reinecker H, Kasper D. Plasmacytoid Dendritic cells mediate anti-inflammatory responses to a gut Commensal molecule via both innate and adaptive mechanisms. *Cell Host Microbe*. 2014;15:413–23.
- Schiavi E, Gleinser M, Molloy E, Groeger D, Frei R, Ferstl R, et al. The surface-associated exopolysaccharide of Bifidobacterium longum 35624 plays an essential role in dampening host Proinflammatory responses and repressing local T H 17 responses. *Appl Environ Microbiol*. 2016;82:7185–96.
- Altmann F, Kosma P, O'Callaghan A, Leahy S, Bottacini F, Molloy E, et al. Genome analysis and characterisation of the exopolysaccharide produced by Bifidobacterium longum subsp. longum 35624™. *PLoS One*. 2016;11:e0162983.
- Groeger D, O'Mahony L, Murphy E, Bourke J, Dinan T, Kiely B, et al. Bifidobacterium infantis35624 modulates host inflammatory processes beyond the gut. *Gut Microbes*. 2013;4:325–39.
- Smolinska S, Groeger D, O'Mahony L. Biology of the microbiome 1: interactions with the host immune response. *Gastroenterol Clin N Am*. 2017;46:19–35.
- Pugin B, Barcik W, Westermann P, Heider A, Wawrzyniak M, Hellings P, et al. A wide diversity of bacteria produce and degrade biogenic amines within the human gastrointestinal tract. *Microb Ecol Health Dis*. 2017;28:1353881.
- Levy M, Thaiss C, Zeevi D, Dohnalová L, Zilberman-Schapira G, Mahdi J, et al. Microbiota-modulated metabolites shape the intestinal microenvironment by regulating NLRP6 Inflammasome signaling. *Cell*. 2015;163:1428–43.
- Herbst T, Sichelstiel A, Schär C, Yadava K, Bürki K, Cahenzli J, et al. Dysregulation of allergic airway inflammation in the absence of microbial colonization. *Am J Respir Crit Care Med*. 2011;184:198–205.
- Fujimura KE, Lynch SV. Microbiota in allergy and asthma and the emerging relationship with the gut microbiome. *Cell Host Microbe*. 2015;17:592–602.
- Rook GA, Adams V, Hunt J, Palmer R, Martinelli R, Brunet LR. Mycobacteria and other environmental organisms as immunomodulators for immunoregulatory disorders. *Springer Semin Immunopathol*. 2004;25:237–55.
- Sudo N, Sawamura S, Tanaka K, Aiba Y, Kubo C, Koga Y. The requirement of intestinal bacterial flora for the development of an IgE production system fully susceptible to oral tolerance induction. *J Immunol*. 1997;159:1739–45.
- Sudo N, XN Y, Aiba Y, Oyama N, Sonoda J, Koga Y, et al. An oral introduction of intestinal bacteria prevents the development of a long-term Th2-skewed immunological memory induced by neonatal antibiotic treatment in mice. *Clin Exp Allergy*. 2002;32:1112–6.
- Gollwitzer ES, Saglani S, Trompette A, Yadava K, Sherburn R, McCoy KD, et al. Lung microbiota promotes tolerance to allergens in neonates via PD-L1. *Nat Med*. 2014;20:642–7.
- McAleer JP, Kolls JK. Contributions of the intestinal microbiome in lung immunity. *Eur J Immunol*. 2017;doi:https://doi.org/10.1002/eji.201646721. [Epub ahead of print].
- Natarajan N, Pluznick JL. From microbe to man: the role of microbial short chain fatty acid metabolites in host cell biology. *Am J Physiol Cell Physiol*. 2014;307:C979–85.
- Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru C, et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease and hematopoiesis. *Nat Med*. 2014;20:159–66.

29. Thorburn AN, McKenzie CI, Shen S, Stanley D, Macia L, Mason LJ, et al. Evidence that asthma is a developmental origin disease influenced by maternal diet and bacterial metabolites. *Nat Commun*. 2015;6:7320.
30. Frei R, Akdis M, O'Mahony L. Prebiotics, probiotics, synbiotics, and the immune system: experimental data and clinical evidence. *Curr Opin Gastroenterol*. 2015;31:153–8.
31. Frei R, Ferstl R, Konieczna P, Ziegler M, Simon T, Rugeles TM, et al. Histamine receptor 2 modifies dendritic cell responses to microbial ligands. *J Allergy Clin Immunol*. 2013;132:194–204.
32. Rodríguez D, Keller AC, Faquim-Mauro EL, de Macedo MS, Cunha FQ, Lefort J, et al. Bacterial lipopolysaccharide signaling through toll-like receptor 4 suppresses asthma-like responses via nitric oxide synthase 2 activity. *J Immunol*. 2003;171:1001–8.
33. Kline JN. Eat dirt: CpG DNA and immunomodulation of asthma. *Proc Am Thorac Soc*. 2007;4:283–8.
34. Nembrini C, Sichelstiel A, Kisielow J, Kurrer M, Kopf M, Marsland BJ. Bacterial-induced protection against allergic inflammation through a multicomponent immunoregulatory mechanism. *Thorax*. 2011;66:755–63.
35. Schuijs MJ, Willart MA, Vergote K, Gras D, Deswarte K, Ege MJ, et al. Farm dust and endotoxin protect against allergy through A20 induction in lung epithelial cells. *Science*. 2015;349:1106–10.
36. Stein MM, Hrusch CL, Gozdz J, Igartua C, Pivniouk V, Murray SE, et al. Innate immunity and asthma risk in Amish and Hutterite farm children. *N Engl J Med*. 2016;375:411–21.
37. Funkhouser LJ, Bordenstein SR. Mom knows best: the universality of maternal microbial transmission. *PLoS Biol*. 2013;11:e1001631.
38. Shukla SD, Budden KF, Neal R, Hansbro PM. Microbiome effects on immunity, health and disease in the lung. *Clin Transl Immunol*. 2017; 6:e133.
39. Barcik W, Pugin B, Westermann P, Perez NR, Ferstl R, Wawrzyniak M, et al. Histamine-secreting microbes are increased in the gut of adult asthma patients. *J Allergy Clin Immunol*. 2016;138:1491–4.
40. Ferstl R, Frei R, Barcik W, Schiavi E, Wanke K, Ziegler M, et al. Histamine receptor 2 modifies iNKT cell activity within the inflamed lung. *Allergy*. 2017; 72:1925–35.
41. Huang YJ, Boushey HA. The microbiome in asthma. *J Allergy Clin Immunol*. 2015;135:25–30.
42. Braun-Fahrlander C, Riedler J, Herz U, Eder W, Waser M, Grize L, et al. Environmental exposure to endotoxin and its relation to asthma in school-age children. *N Engl J Med*. 2002;347:869–77.
43. Roduit C, Frei R, Depner M, Schaub B, Loss G, Genuneit J, et al. Increased food diversity in the first year of life is inversely associated with allergic diseases. *J Allergy Clin Immunol*. 2014;133:1056–64.
44. Frei R, Roduit C, Bieli C, Loeliger S, Waser M, Scheynius A, et al. Expression of genes related to anti-inflammatory pathways are modified among farmers' children. *PLoS One*. 2014;9:e91097.
45. Frei R, Ferstl R, Roduit C, Ziegler M, Schiavi E, Barcik W, et al. Exposure to nonmicrobial N-glycolylneuraminic acid protects farmers' children against airway inflammation and colitis. *J Allergy Clin Immunol*. 2017;S0091-6749(17)30994-6. doi:<https://doi.org/10.1016/j.jaci.2017.04.051>. [Epub ahead of print].
46. Weiss ST, Litonjua AA, Vitamin D. The gut microbiome, and the hygiene hypothesis. How does asthma begin? *Am J Respir Crit Care Med*. 2015;191:492–3.
47. Rusconi F, Zugna D, Annesi-Maesano I, Baiz N, Barros H, Correia S, et al. Mode of delivery and asthma at school age in 9 European birth cohorts. *Am J Epidemiol*. 2017;185:465–73.
48. van Nimwegen FA, Penders J, Stobberingh EE, Postma DS, Koppelman GH, Kerkhof M, et al. Mode and place of delivery, gastrointestinal microbiota, and their influence on asthma and atopy. *J Allergy Clin Immunol*. 2011;128:948–55.
49. Abrahamsson T, Jakobsson H, Andersson A, Björkstén B, Engstrand L, Jenmalm M. Low gut microbiota diversity in early infancy precedes asthma at school age. *Clin Exp Allergy*. 2014;44:842–50.
50. Arrieta M, Stiemsma L, Dimitriu P, Thorson L, Russell S, Yurist-Doutsch S, et al. Early infancy microbial and metabolic alterations affect risk of childhood asthma. *Sci Transl Med*. 2015;7:307ra152.
51. Fujimura KE, Sitarik AR, Havstad S, Lin DL, Levan S, Fadrosch D, et al. Neonatal gut microbiota associates with childhood multisensitized atopy and T cell differentiation. *Nat Med*. 2016;22:1187–91.
52. Human Microbiome Project, C. Structure, function and diversity of the healthy human microbiome. *Nature*. 2012;486:207–14.
53. Hilty M, Burke C, Pedro H, Cardenas P, Bush A, Bossley C, et al. Disordered microbial communities in asthmatic airways. *PLoS One*. 2010;5:e8578.
54. Bisgaard H, Hermansen MN, Buchvald F, Loland L, Halkjaer LB, Bønnelykke K, et al. Childhood asthma after bacterial colonization of the airway in neonates. *N Engl J Med*. 2007;357:1487–95.
55. Huang YJ, Charlson ES, Collman RG, Colombini-Hatch S, Martinez FD, Senior RM. The role of the lung microbiome in health and disease. A National Heart, Lung, and Blood Institute workshop report. *Am J Respir Crit Care Med*. 2013;187:1382–7.
56. Bassis CM, Erb-Downward JR, Dickson RP, Freeman CM, Schmidt TM, Young VB, et al. Analysis of the upper respiratory tract microbiotas as the source of the lung and gastric microbiotas in healthy individuals. *MBio*. 2015;6:e00037.
57. Charlson ES, Diamond JM, Bittinger K, Fitzgerald AS, Yadav A, Haas AR, et al. Lung-enriched organisms and aberrant bacterial and fungal respiratory microbiota after lung transplant. *Am J Respir Crit Care Med*. 2012;186:536–45.
58. Man WH, de Steenhuisen Piters WA, Bogaert D. The microbiota of the respiratory tract: gatekeeper to respiratory health. *Nat Rev Microbiol*. 2017;15:259–70.
59. Goleva E, Jackson LP, Harris JK, Robertson CE, Sutherland ER, Hall CF, et al. The effects of airway microbiome on corticosteroid responsiveness in asthma. *Am J Respir Crit Care Med*. 2013;188:1193–201.
60. Denner DR, Sangwan N, Becker JB, Hogarth DK, Oldham J, Castillo J, et al. Corticosteroid therapy and airflow obstruction influence the bronchial microbiome, which is distinct from that of bronchoalveolar lavage in asthmatic airways. *J Allergy Clin Immunol*. 2016;137:1398–405.
61. Huang YJ, Nelson CE, Brodie EL, Desantis TZ, Baek MS, Liu J, et al. Airway microbiota and bronchial hyperresponsiveness in patients with suboptimally controlled asthma. *J Allergy Clin Immunol*. 2011;127:372–81.
62. Marri PR, Stern DA, Wright AL, Billheimer D, Martinez FD. Asthma-associated differences in microbial composition of induced sputum. *J Allergy Clin Immunol*. 2013;131:346–52.
63. Huang YJ, Nariya S, Harris JM, Lynch SV, Choy DF, Arron JR, et al. The airway microbiome in patients with severe asthma: associations with disease features and severity. *J Allergy Clin Immunol*. 2015;136:874–84.
64. Carr TF, Kraft M. Chronic infection and severe asthma. *Immunol Allergy Clin N Am*. 2016;36:483–502.
65. Wong EH, Porter JD, Edwards MR, Johnston SL. The role of macrolides in asthma: current evidence and future directions. *Lancet Respir Med*. 2014;2:657–70.
66. Slater M, Rivett DW, Williams L, Martin M, Harrison T, Sayers I, et al. The impact of azithromycin therapy on the airway microbiota in asthma. *Thorax*. 2014;69:673–4.
67. Biesbroek G, Tsvitvadze E, Sanders EA, Montijn R, Veenhoven RH, Keijsers BJ, et al. Early respiratory microbiota composition determines bacterial succession patterns and respiratory health in children. *Am J Respir Crit Care Med*. 2014;190:1283–92.
68. Teo SM, Mok D, Pham K, Kusel M, Serralha M, Troy N, et al. The infant nasopharyngeal microbiome impacts severity of lower respiratory infection and risk of asthma development. *Cell Host Microbe*. 2015;17:704–15.
69. Stiemsma LT, Turvey SE. Asthma and the microbiome: defining the critical window in early life. *Allergy Asthma Clin Immunol*. 2017;13:3.
70. Burbank AJ, Sood AK, Kesic MJ, Peden DB, Hernandez ML. Environmental determinants of allergy and asthma in early life. *J Allergy Clin Immunol*. 2017;140:1–12.
71. Ong MS, Umetsu DT, Mandl KD. Consequences of antibiotics and infections in infancy: bugs, drugs, and wheezing. *Ann Allergy Asthma Immunol*. 2014; 112:441–5.
72. Singanayagam A, Ritchie AI, Johnston SL. Role of microbiome in the pathophysiology and disease course of asthma. *Curr Opin Pulm Med*. 2017;23:41–7.
73. Esposito S, Principi N. Impact of nasopharyngeal microbiota on the development of respiratory tract diseases. *Eur J Clin Microbiol Infect Dis*. 2017;doi:<https://doi.org/10.1007/s10096-017-3076-7>. [Epub ahead of print].
74. Sokolowska M, Chen LY, Liu Y, Martinez-Anton A, Logun C, Alsaaty S, et al. Dysregulation of lipidomic profile and antiviral immunity in response to hyaluronan in patients with severe asthma. *J Allergy Clin Immunol*. 2017; 139:1379–83.
75. Vissing NH, Chaves BL, Bisgaard H. Increased risk of pneumonia and bronchiolitis after bacterial colonization of the airways as neonates. *Am J Respir Crit Care Med*. 2013;188:1246–52.

76. de Steenhuijsen Piters WA, Heinonen S, Hasrat R, Bunsow E, Smith B, Suarez-Arrabal MC, Chaussabel D, et al. Nasopharyngeal microbiota, host Transcriptome, and disease severity in children with respiratory Syncytial virus infection. *Am J Respir Crit Care Med*. 2016;194:1104–15.
77. Mansbach JM, Hasegawa K, Henke DM, Ajami NJ, Petrosino JF, Shaw CA, et al. Respiratory syncytial virus and rhinovirus severe bronchiolitis are associated with distinct nasopharyngeal microbiota. *J Allergy Clin Immunol*. 2016;137:1909–13.
78. van den Bergh MR, Biesbroek G, Rossen JW, de Steenhuijsen Piters WA, Bosch AA, van Gils EJ, et al. Associations between pathogens in the upper respiratory tract of young children: interplay between viruses and bacteria. *PLoS One*. 2012;7:e47711.
79. Bosch AA, Levin E, van Houten MA, Hasrat R, Kalkman G, Biesbroek G, et al. Development of upper respiratory tract microbiota in infancy is affected by mode of delivery. *E Bio Med*. 2016;9:336–45.
80. Hasegawa K, Linnemann RW, Mansbach JM, Ajami NJ, Espinola JA, Fiechtner LG, et al. Household siblings and nasal and fecal microbiota in infants. *Pediatr Int*. 2017;59:473–81.
81. Chiu CY, Chan YL, Tsai YS, Chen SA, Wang CJ, Chen KF, et al. Airway microbial diversity is inversely associated with mite-sensitized rhinitis and asthma in early childhood. *Sci Rep*. 2017;7:1820.
82. Jatzlauk G, Bartel S, Heine H, Schlöter M, Krauss-Etschmann S. Influences of environmental bacteria and their metabolites on allergies, asthma, and host microbiota. *Allergy*. 2017;72:1859–67.
83. Birzele LT, Depner M, Ege MJ, Engel M, Kublik S, Bernau C, Loss GJ, et al. Environmental and mucosal microbiota and their role in childhood asthma. *Allergy*. 2017;72:109–19.
84. Depner M, Ege MJ, Cox MJ, Dwyer S, Walker AW, Birzele LT, et al. Bacterial microbiota of the upper respiratory tract and childhood asthma. *J Allergy Clin Immunol*. 2017;139:826–34.
85. Ege MJ, Mayer M, Normand AC, Genuneit J, Cookson WO, Braun-Fahrlander C, et al. Exposure to environmental microorganisms and childhood asthma. *N Engl J Med*. 2011;364:701–9.
86. Lyons A, O'Mahony D, O'Brien F, MacSharry J, Sheil B, Ccedia M, et al. Bacterial strain-specific induction of Foxp3+ T regulatory cells is protective in murine allergy models. *Clin Exp Allergy*. 2010;40:811–9.
87. Van De Pol M, Lutter R, Smids B, Weersink E, Van Der Zee J. Synbiotics reduce allergen-induced T-helper 2 response and improve peak expiratory flow in allergic asthmatics. *Allergy*. 2010;66:39–47.
88. Lee SC, Yang YH, Chuang SY, Huang SY, Pan WH. Reduced medication use and improved pulmonary function with supplements containing vegetable and fruit concentrate, fish oil and probiotics in asthmatic school children: a randomised controlled trial. *Br J Nutr*. 2013;110:145–55.
89. Bieber T, Akdis C, Lauener R, Traidl-Hoffmann C, Schmid-Grendelmeier P, Schappi G, et al. Global allergy forum and 3rd Davos declaration 2015: atopic dermatitis/eczema: challenges and opportunities toward precision medicine. *Allergy*. 2016;71:588–92.
90. Gray L, O'Hely M, Ranganathan S, Sly P, Vuillermin P. The maternal diet, gut bacteria, and bacterial metabolites during pregnancy influence offspring asthma. *Front Immunol*. 2017;8:365.
91. Bakker G, Nieuwdorp M. Fecal microbiota transplantation: therapeutic potential for a multitude of diseases beyond *Clostridium Difficile*. *Microbiol Spectr*. 2017;5.
92. Zitvogel L, Galluzzi L, Viaud S, Vetizou M, Dailly R, Merad M, et al. Cancer and the gut microbiota: an unexpected link. *Sci Transl Med*. 2015;7:271ps1.
93. Sivan A, Corrales L, Hubert N, Williams J, Aquino-Michaels K, Earley Z, et al. Commensal *Bifidobacterium* promotes antitumor immunity and facilitates anti-PD-L1 efficacy. *Science*. 2015;350:1084–9.
94. Barcik W, Wawrzyniak M, Akdis CA, O'Mahony L. Immune regulation by histamine and histamine-secreting bacteria. *Curr Opin Immunol*. 2017;48:108–13.
95. Durack J, Lynch S, Nariya S, Bhakta N, Beigelman A, Castro M, et al. Features of the bronchial bacterial microbiome associated with atopy, asthma, and responsiveness to inhaled corticosteroid treatment. *J Allergy Clin Immunol*. 2017;140:63–75.
96. Ozturk A, Turturice B, Perkins D, Finn P. The potential for emerging microbiome-mediated therapeutics in asthma. *Curr Allergy Asthma Rep*. 2017;17:62.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

