

POSTER PRESENTATION

Open Access

Additional stimulation of sGC on top of standard treatment with ARB's may offer a new therapeutic approach for the treatment of diabetic nephropathy resistant to ARB treatment alone

Markus Alter^{1,2,3}, Ina Ott¹, Karoline von Websky^{1,2}, Oleg Tsuprykov^{1,2}, Yuliya Sharkovska², Katharina Krause-Relle^{2,3}, Jens Raila³, Andrea Henze³, Axel Kretschmer⁴, Johannes-Peter Stasch^{4,5}, Berthold Hocher^{2,3*}

From 5th International Conference on cGMP: Generators, Effectors and Therapeutic Implications
Halle, Germany. 24-26 June 2011

Background

Riociguat is the first of a new class of drugs, the soluble guanylate cyclase (sGC) stimulators. Riociguat has a dual mode of action: it sensitizes sGC to the body's own NO and can also increase sGC activity in the absence of NO. The NO-sGC-pathway is impaired in many cardiovascular diseases such as heart failure, pulmonary hypertension and diabetic nephropathy (DN). DN leads to high cardiovascular morbidity and mortality. There is still a high unmet medical need. The urinary albumin excretion rate is a predictive biomarker for these clinical events. Therefore, we investigated the effect of riociguat, alone and in combination with the angiotensin II receptor antagonist (ARB) telmisartan on the progression of DN in diabetic eNOS knock out mice, a new model closely resembling human pathology.

Methods

Seventy-six male eNOS knockout C57BL/6J mice were divided into 4 groups after receiving intraperitoneal high-dose streptozotocin: telmisartan (1 mg/kg), riociguat (3 mg/kg), riociguat+telmisartan (3 and 1 mg/kg), and vehicle. Fourteen mice were used as non-diabetic controls. After 12 weeks, urine and blood were obtained and blood pressure measured. Glucose concentrations were highly increased and similar in all diabetic groups.

Results

Riociguat, alone (105.2 ± 2.5 mmHg; mean \pm SEM; n = 14) and in combination with telmisartan (105.0 ± 3.2 mmHg; n = 12), significantly reduced blood pressure versus diabetic controls (117.1 ± 2.2 mmHg; n = 14; p = 0.002 and p = 0.004, respectively), whereas telmisartan alone (111.2 ± 2.6 mmHg) showed a modest blood pressure lowering trend (p = 0.071; n = 14). The effects of single treatment with either riociguat (97.1 ± 15.7 μ g/d; n = 13) or telmisartan (97.8 ± 26.4 μ g/d; n = 14) did not significantly lower albumin excretion on its own (p = 0.067 and p = 0.101, respectively). However, the combined treatment led to significantly lower urinary albumin excretion (47.3 ± 9.6 μ g/d; n = 12) compared to diabetic controls (170.8 ± 34.2 μ g/d; n = 13; p = 0.004), and reached levels similar to non-diabetic controls (31.4 ± 10.1 μ g/d, n = 12).

Conclusion

Riociguat significantly reduced urinary albumin excretion in diabetic eNOS knock out mice that were refractory to treatment with ARB's alone. Patients with diabetic nephropathy refractory to treatment with ARB's have the worst prognosis among all patients with diabetic nephropathy. Our data indicate that additional stimulation of sGC on top of standard treatment with ARB's may offer a new therapeutic approach for patients with diabetic nephropathy resistant to ARB treatment.

* Correspondence: berthold.hocher@charite.de

²Center for Cardiovascular Research, Charité, Campus Mitte, Berlin, Germany
Full list of author information is available at the end of the article

Author details

¹Department of Nephrology, Charité, Campus Benjamin Franklin, Berlin, Germany. ²Center for Cardiovascular Research, Charité, Campus Mitte, Berlin,

Germany. ³Institute for Nutritional Science, University of Potsdam, Germany.

⁴Bayer HealthCare AG, Cardiovascular Research, Wuppertal, Germany.

⁵Institute of Pharmacy, Martin-Luther-University of Halle, Germany.

Published: 1 August 2011

doi:10.1186/1471-2210-11-S1-P1

Cite this article as: Alter *et al.*: Additional stimulation of sGC on top of standard treatment with ARB's may offer a new therapeutic approach for the treatment of diabetic nephropathy resistant to ARB treatment alone. *BMC Pharmacology* 2011 11(Suppl 1):P1.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

