A 575134

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN



Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

H1608
H15(1)

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

THE

CHRONIC DISEASES:

THEIR

SPECIFIC NATURE

AND

HOMŒOPATHIC TREATMENT.

BY

Dr. SAMUEL HAHNEMANN.

TRANSLATED AND EDITED

BY

CHARLES J. HEMPEL, M.D.

VOLUME IV.

ANTI-PSORIC REMEDIES.

NEW-YORK:
WM. RADDE, 322 BROADWAY.

LONDON:
H. BALLIERE, 219 REGENT-STREET.

1846.

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

Entered according to the Act of Congress, in the year 1846, by

WILLIAM RADDE,

In the Clerk's Office of the District Court for the Southern District of New-York.

H. LUDWIG, PRINTER,
70 & 72 Vesey-st.

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

PREFACE.

Inquiry into the modus operandi of a homœopathic drug.

We have no means of knowing the inmost functions of the organism; sometimes we may feel at liberty to draw speculative conclusions from results relative to the nature of the causes in which they have originated, but we cannot sustain our explanations by illustrating them by the phenomena of material nature; for this reason, that the functions of the living organism have nothing in common with the changes of matter, but on the contrary, take place in an entirely different manner.

Possessed by these convictions, it is quite natural that I should never have attempted to explain how diseases can be cured by medicines, the pathogenetic symptoms of which are similar to those of the disease. I have furnished a conjecture relative to that matter, without wishing it to be considered an explanation. Such an explanation is not necessary. It simply behoves us to apply the law "like cures like" correctly and successfully, and not to boast with abstract explanations, in spite of which the patient may remain uncured. Indulging foolish hypotheses is all that the so-called physicians of the Old School have accomplished up to this moment. These physicians would not have hesitated to reject the doctrines of homœopathy, which however teach the only true method of cure, simply because my explanation of the modus operandi of homœopathic agents was not satisfactory.

If I again attempt to throw some light on the inmost

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

action of homœopathic drugs, it is not because I expect to satisfy my opponents, but in order to furnish more probable explanatory suggestions to the real adherents and able advocates of the homœopathic system, and to satisfy the irresistible, innocent and praiseworthy impulse of the human mind to account for the mode in which we accomplish changes from evil to good.

It is undeniable, that our vital powers, when unassisted by remedial agents, are unable to cure diseases, even those of short duration—taking it for granted that those powers are not overcome by the disease, without expending by a so-called crisis a portion, and frequently even a considerable portion, of both the fluids and solids of our organism. I have shown this elsewhere. We shall, perhaps, never be able to explain this mode of action of the vital powers; but we know with certainty that they do not accomplish a victory over disease in a *direct* manner, or without making great sacrifices. The vital powers are not even capable of healing the chronic, miasmatic diseases, and of substituting health in their stead, without inflicting a loss upon the organism. But it is certain, that although the vital forces may achieve a victory over the acute as well as miasmatic diseases without inflicting such losses upon the organism, provided they are assisted and directed in their action by a properly selected homœopathic agent, it is nevertheless those forces that conquer, in the same sense as the native army is said to have beaten the enemy, although it may have been assisted by auxiliary troops. Those vital forces, unless assisted by a homœopathic drug, cannot control a natural disease without inflicting great sacrifices upon the organism; for, naturally, they have only power to maintain the unimpeded evolution of the functions of the animal economy, as long as those functions are not disturbed by the improper interference of an external agent.

The unassisted vital forces are no match for such hostile,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

disturbing agents; they are even weaker than those agents—oppose them with many symptoms of suffering, and cannot overpower them, be they either acute or chronic, without inflicting great losses on the organism. This result can only be avoided by assisting the vital forces by a proper remedial agent. God has appointed the true physician to this office.

The homœopathic agent is alone able to secure to the vital forces that superior power which is necessary to conquer the enemy.

I repeat, that the vital forces, the natural destination of which is the regulation of the undisturbed functions of the organism, oppose but a feeble resistance to the approaching enemy, which waxes stronger in proportion as this enemy becomes more powerful, but which, even in the most favorable circumstances, is scarcely equal, and in weak patients, not even equal to the power of the assailing enemy: the vital forces are neither able, nor naturally destined to overpower that enemy, nor can they oppose him without being injured.

By magnifying ever so little the enemy by whom the vital forces are assailed—which we accomplish by invading the vital forces by an artificially morbid influence closely resembling the disease, we induce those forces, as from an instinct of self-preservation, gradually to unfold all their energies until they become powerful enough to control the disease; the vital forces are then restored to the regulation of the animal functions, whereas, the artificial medicinal disease disappears of itself, so soon as we discontinue the use of those agents by which it had been established in the organism.

The principle by which the body is animated, is infinitely great and lasting, provided we maintain its integrity by a proper diet, and relieve it from an invading enemy by proper homœopathic treatment.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

INDEX OF REMEDIES

CONTAINED IN THIS VOLUME.

	Page.
Iodine,	6
Kali Carbonicum,	26
Lycopodium,	80
Magnesia Carbonica,	138
Magnesia Muriatica,	170
Manganum,	197
Daphne Mezereum,	219
Muriatic Acid,	242
Natrum Carbonicum,	261
Natrum Muriaticum,	294
Nitric Acid,	333

CORRECTION.

Read "of" in the place of "by," page first, line 10, from above.

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

Digitized by
UNIVERSITY OF MICHIGAN

Original from
UNIVERSITY OF MICHIGAN

ANTI-PSORIC REMEDIES.

JODIUM—IODINE.

DYNAMISE one grain of iodine according to the mode taught in the first volume.

Iodine is a heroic medicine, even when employed in the highest potencies; its use requires all the discretion of a good homœopathic physician, lest he should abuse this substance, as allœopathic physicians do, who frequently destroy the constitution and even the life of their patients, by enormous and improperly applied doses of the drug in question.

Iodine has rendered especial service in affections with he following symptoms: Dizziness in the morning; throbbing in the head; pain as from excoriation in the eyes; buzzing in the ears; hardness of hearing; coated tongue; mercurial ptyalism; bad taste, as of soap; sour eructations, with a sense as of burning; *heartburn*, after eating heavy food; *canine hunger;* nausea; incarceration of flatulence; distension of the abdomen; constipation; nightly micturition; delaying menses; cough; old morning cough; difficulty of breathing; external swelling of the neck; weakness of the arms, early in the morning when in bed; going to sleep of the fingers; curvature of the bones; dryness of the skin; night sweats.

This drug has been proved by Drs. Gersdorff, Gross, Hartlaub and Trinks, Schreter.

From *Noack and Trinks*: Iodine is useful in the following affections: *Scrophulosis*, (it is supposed by some that the muriate of baryta will dispute the rank with iodine in the treatment of scrofula.) Glandular affections, especially swelling of the thyroid gland, hypertrophy of the tonsillæ, parótid gland, &c. *Affections of the mucous nembranes*, especially when depending upon a scrofulous and leuco-phlegmatic disposition; and particularly when this condition seems to affect all the functions of the

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

being; in catarrhal affections of the Schneiderian membrane without any fever, of the Eustachian tube, larynx, trachea, vagina, urethra, with disposition to thin, white coryza and obstruction of the nose, especially in persons of between 20 and 30 years, brought on by cold and neglected catarrhal fever. *Syphilis, with tendency to hypertrophy;* chancre elevé. Iodine has cured a chancre on the frænulum præputii; the frænulum having become detached, that portion of the ulcer which was situated on the frænulum, became elevated, whilst that other portion which was situated upon the glans was deepened, without, however, showing the characteristic edges of a chancre. The cure was effected in this way:—the edges became dry first, the ulcer became gradually covered with a brown crust, under which the suppurative process continued for a time, but afterwards left a solid, star-shaped cicatrix, there being very little substance lost, so that the cicatrix appeared rather even than excavated. *Old gout,* characterised by violent, nightly pains, in various joints, without any swelling; spina ventosa and caries of the spongy extremities of the bones. Arsenic disease. *Tabes mercurialis.* Dropsy; dropsy after fever and ague, in a scrofulous female; various forms of intermittent fever and ascites, intimately connected with disturbances of the spleen; scarlatina, when *Bell.* would do no good, (giving mercury first.) Sore nipples, consequent upon scrofulous conditions. Scirrhous induration. Febris meseraica; Lobethal observes, that in febris meseraica, iodine is inferior to CALC. CARB., when this fever indicates, as it often does, very insidiously, an approaching atrophy. Inflamed, big, bloated eyelids. Humming before the ears and hardness of hearing, being remains of frequent colds or catarrhal inflammations of the throat, when we can with certainty diagnose an accumulation of mucus in the Eustachian tube by the humming being increased by chewing or any other motion of the jaws, or when the cold frequently returns, or when the patient complains of his ear being closed by a valve. Hard hearing from swelling of the tonsillæ; swelling of the parotid gland. Obstinate and torpid swelling of the cervical glands. Constitutional stomacace. Chronic, painless vomiting, (in the case of a

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

carder, whose affection was caused by pressing upon the heavy scissors.) *Affections of the liver.* Scirrhous swelling of the inguinal glands, resulting from metaschematismus of the gout. *Diseases of the urinary organs.* Metritis and oophoritis (in conjunction with Ars.)—amenorrhœa, with frequent and violent colic, sometimes violent stitches in the left side of the chest when breathing, frequent, weak, and dry cough, with stitches in the throat, paleness of the face, generally alternating with sudden redness and burning of the ears, want of breath when going up stairs, loss of strength, constantly cold hands covered with cold sweat when working, (in this case the pain disappeared, but the menses did not come on.) Leucorrhœa, of scrofulous females; chronic leucorrhœa when accompanied by a pale, cachectic and chlorotic complexion and disposition to catarrhal affections. *Diseases of the trachea.* Profuse secretion of mucus in the larynx, brought on by influenza or a catarrhal fever, with rough voice, tingling and burning in the throat, in persons who incline to phthisis. Cough remaining after an epidemic influenza; violent influenza; cough, with spitting of blood; pain in the chest and high fever, (preceded by aconite.) Cough, with tenacious, white expectoration, excited by an irritation under the upper portion of the sternum—worst early in the morning. Chronic catarrh of persons who incline to phthisis laryngea or pulmonalis. Croup, in alternation with aconite. Hooping-cough, with intolerable tickling in the whole chest. Cough, first periodical, afterwards continuous, especially in the morning when rising (remaining after inflammation of the lungs;) irritation, exciting a cough in the left and lower part of the chest, four fingers' breadth above the pit of the stomach, the cough being sometimes accompanied by a slight, concealed pain in that part, whitish or yellowish expectoration, temporary asthma, and shortness of breath, stitching pain in the affected side of the chest when walking fast during severe cold, continual and violent fluent and dry coryza, complete absence of smell and taste; and sad, vexed, solicitous mood, in alternation with mercury. Pneumonia, when hepatization threatens to set in. Suspicious cough, with weakness and emaciation, hysteric convulsions, toothache, leucor-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

rhœa, hard nodosity of the labia, occasional tearing accompanied by continual darting, (in a patient of delicate constitution, having suffered formerly with herpes scabiosus, tearing in the axillary glands, hard nodosities in the right axilla and mamma, discharge of blood from the vagina, and her mother having died with cancer of the womb.) Pulmonary consumption, tuberculous and pituitous. Arthropyosis scrophulosa. Scrofulous inflammation of the knee, considerable bright-red, shining, hot, tense swelling of the knee, darting stitches and burning in the swelling, small, red openings discharging a bloody and watery pus, pain increased by motion, or by bending the knee, or pressing upon it. Dropsy of the knee-joint. White swelling of the knee. Inflammation of the synovial membrane of the knee: large, elastic, fluctuating swelling of the knee, simple redness of the skin, with bright-red, hot, painful, scattered blotches, increased temperature of the knee, dull pains in the interior of the joint while the body is extended and at rest, intense pain when making the least motion, or when bending or pressing the part, occasional paroxysms of fever, afterwards a small opening above the patella, discharging a healthy pus in the beginning, but afterwards a clear, light-yellow, serous fluid, in pretty large quantity.

ANTIDOTES—Of large doses. Starch boiled to a thin glue by means of a large quantity of water, and given to the patient in large doses; or, in case of necessity, sugar-water, milk, especially asses' milk; tepid baths; sulphur baths. MERC. and ARS., frequent walks in the open air; Op. Hyosc., (in a case of endogastritis, brought on by abuse of iodine;) Gratiola. Altschuhl says he has cured an iodine cachexia, with iodine 30, 2 pellets. Antidotes of small doses: Ant., Ars., Camph., Chi., Chin. sulph., Coff., Hep. s., Op., Phosp., Spong.

Iodine is especially useful after, or in alternation with, mercury.

MORAL SYMPTOMS.—*Despondency.*—Gloomy mood. Sad, melancholy mood. Hypochondriac mood. Sadness. Out of humor and sensitive, during the period of digestion, from noon till evening, with an oppressive sensation in the throat and chest, such as one feels when one is about to weep. *Oppression of the chest.* Continued disposition

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

to weep. Alternate disposition to weep and to be cheerful. Oppressive anxiety after having performed some manual labor, going off when sitting. *Anguish.*—Anxiety. Great anguish. Oppression of the chest. Anguish and despondency; the patients are mostly thinking of the present moment. Impatient moving about; she is running about all the time, and never sits down; nor does she sleep at night; one ought to suppose her crazy. Restlessness—the body is continually moving. Aversion to sitting still. He apprehends this or that accident from every trifle. Discouragement. Feeling of despondency, oppressive and continuing with the pain. Aversion to work. He deems himself incapable to do any thing. She is no longer phlegmatic; she has become a blooming, stronger, and more animated girl. Vexed, out of humor; nothing is done to her satisfaction. Extreme disposition to get angry. Increased sensitiveness and irritability. Sensitiveness to noise. Increased sensitiveness to external impressions. Irritated nerves and feelings. Excessive nervous irritation. Excessively merry and talkative; she does not give any one a chance to say a word. *Illusions of the sense of tact.*—Delirium. Fixed, immoveable thoughts, (21. d.)

Head.—Obtusion of the head, making thought difficult. *Obtusion of the head with great disinclination to serious labor.* Obtusion of the head, which appears to come from the back and move up along the nape of the neck. Obtusion of the head, becoming an oppressive pain, (a. 1 h.) Slight obtusion of the head, with pressure in the right half of the forehead, and great hunger, (a. 1, 2 h.) *Vertigo.* Vertigo, drawing one forwards. Vertigo with weakness, early in the morning. *Headache.* Headache, so violent that it makes him almost crazy. Quickly passing pain in the occiput. Headache in warm air, when driving a long time, or walking fast. Pain in the forehead and top of the head, increasing at every noise or conversation. Headache as if a ribbon were tied round the head very firmly. Headache in the forehead; the brain feels bruised and is extremely sensitive; the whole body, and especially the arms are weak and feel paralysed; he had to lie down; accompanied by eructations and painful sensitiveness of

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

the external head during motion; the obtusion of the head and painfulness of the face continue even the next day, (a. 26 d.) Pressure on a small spot of the forehead, just above the root of the nose. Headache, especially in the left side towards the forehead, frequently returning. Pressure in the region of the vertex, for 10 minutes. Headache, especially in the temples, going and coming. Aching in the occiput, moderate when at rest, violent during motion, and resulting at last in a sensation of great dreariness in the whole head, (1. d.) Aching over the eyes, towards evening. Violent aching in the base of the occiput, in the afternoon, in the open air. Sharp aching, *on the left side over the forehead.* Drawing pressure in the left upper half of the head, as far as the temples. Aching and sometimes lancinating headache. Drawing pain in the left side of the head, as far as the teeth. Tearing, first in the left, afterwards in the right temporal region, almost at the same time. Tearing headache over the left eye and in the temple. Tearing with pressure on the right side over the forehead. Stitches in the occiput, improved by the recumbent posture. Stitches in the top of the head, (a. 3 d.) Throbbing in the forehead, without any pain. * Throbbing in the head at every motion, (a. 24 h.) *Congestion of blood to the head.* The congestion of blood to the head is increased in those who have a disposition for it. Congestion of blood to the head, and afterwards, at 2 o'clock in the afternoon, headache of half an hour, recurring at 5 o'clock. Pain in the forehead, as of sub-cutaneous ulceration. Smarting sore feeling in the occiput, on the right side, over the ear, towards the back part of the head, in the skin. Falling off of the hair. Considerable falling off of the hair.

Eyes.—Pain in the orbits. Pressure in the eyes as of sand. Pressure in the eyes. Tension over the right eye, the eyes being somewhat inflamed. Tearing around the right eye, especially below it. Stitches in the upper part of the left eyeball. Cutting stitches in the left eye, in the direction of the external canthus. Itching in the canthi. Itching of the eyelids. Redness and swelling of the eyelids, with nightly agglutination. *Inflamed eyes.* Watery white swelling of the eyelids. The white of the eye is of

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

a dirty-yellow, traversed by injected veins. Weak feeling around the eyes, as if they were lying deep, especially in the afternoon. *Lachrymation.* Twitchings of the eyes, (in a few hours.) Continual twitching to and fro of the lower eyelid. Tremor of the eyelids. Turbid sight. Light looks to her dimmer and more indistinct. Objects seem to her as if she were looking at them through a gauze. *Obscuration of sight. Weak sight.* Sometimes the things she looks at, seem to her multiplied; she is not able "then" to distinguish them clearly. Dark curls are floating downwards in front of the eyes, sideways and close to the direction of the visual ray, (a. 16 h.) Occasional twinkling before the eyes. She is unable to do fine sewing, on account of the stitches flitting before her eyes. Sparks before the eyes. Fiery, curved rays are shooting downwards in a line near the visual ray, and likewise at a small distance from the focus of the ray, round about the eye, but more upwards, (a. 24 h.)

Ears.—Straining pain in the left ear. Forcing in the right ear. Tearing pressure in the little groove, under the right ear and near it, in the neck. Small yellow scurf on the concha. Sensitiveness of the hearing to the noise, (4. d.) * *Hardness of hearing.* Frequent din in the right ear as if in a mill.

Nose.—Pain in the lower part of the nose, when blowing it, without any coryza. Itching stinging, in the forepart of the septum of the nose. Red, burning spot on the nose, under the eye. Itching elevation on the nose. Small little scab in the right nostril. Bleeding of the nose when blowing it. *Profuse bleeding of the nose.*

Face.—The complexion becomes changed. Pale, contracted face. *Pale face.* * *Yellow complexion.* The yellow complexion becomes more pale, (curative action.) The face which had been yellow heretofore, becomes brown, so that in a few days the skin of a female of 28 years looked smoked. The features become disfigured, altered. Sunken eyes. *Convulsive twitchings of the facial muscles.* Suppurating ulcer on the left cheek with swelling of the surrounding glands, and a hard nodosity at the place which the ulcer occupies, dispersing very slowly;

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

shortly after, a second ulcer appeared near the former; it healed more speedily.

TEETH AND JAWS.—Aching in the right upper jaw. Painless swelling in the sub-maxillary glands, (2. d.) *Oppressive toothache in alternate places, in the molar teeth, on both sides.* Clawing toothache in the posterior molares of the right side. Drawing pain in the teeth of the right side, towards the ear, accompanied by stitches. Cutting drawing and sore feeling in the roots or the gums of the lower incisores, at times on the left, at times on the right side. Pain as from looseness, in the teeth and gums, when eating. In the morning, the teeth are covered with more mucus, are coloured more yellow and are blunted more rapidly by weak vegetable acids. The gums are painful to the touch. Increased redness of the gums. Inflammation and swelling of the gums. Bleeding of the gums. Ulcer on the gums of one of the lower hollow molares, with swelling of the cheek extending to the lower border of the orbit. Vesicles made occasionally their appearance in the mouth, with swelling of the gums.

MOUTH.—Aphthæ in the mouth, with ptyalism. Small elevation on the inner side of the right cheek; in the beginning they felt aching and sore when touched; in a few days they became lancinating and cutting like an ulcer, especially when opening the mouth widely, when eating or reading aloud, with inflammation of the parts around. The glands on the inner side of the cheeks, are intensely painful, as if one had sharp vinegar in one's mouth.—Smarting and sensation as of pulling in the tonsillæ, (immediately.) Putrid smell from the mouth, even before breakfast, immediately after rinsing the mouth with pure water. Troublesome dryness of the tongue. *Coated tongue.* * Tongue thickly coated, the coating looking like the substances that were thrown up. Pressure in the left half of the palate. Swelling and elongation of the uvula with a good deal of spitting. Considerable constriction of the throat.

THROAT.—Constrictive sensation in the fauces. Tormenting constriction of the throat. Impeded deglutition when drinking water, as if the fauces were constricted and too weak to press the beverage down. Feeling of fulness

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

in the throat, (6. d.) Pain in the œsophagus, increased by pressing upon the larynx. Aching on the right side in the throat, more between than during the acts of deglutition. Tearing in the throat, above the larynx. Stitches in the throat, apparently in the larynx, somewhat perceptible during the act of deglutition. Itching titillation in the throat, in the region of the larynx, early in the morning, when in bed. Disagreeable scraping in the fauces, with copious secretion of saliva. Burning and scraping in the fauces, passing quickly. Burning and hot feeling in the fauces. Inflammation and ulcers of the œsophagus. *Increased secretion of saliva.* Increased secretion of saliva, obliging one to spit frequently, for several days. Frequent and watery saliva in the mouth.

Taste and Appetite.—Bad taste in mouth, especially when the stomach is either quite empty or quite full. Abominable taste in mouth. *Salt taste in mouth*, (1. and 2. d.) Bitter taste in mouth. Bitter taste, in the afternoon; sweet plums taste bitter to her. Sourish, offensive taste in mouth, the whole day, especially when smoking tobacco. Sweet taste on the tip of the tongue, (6. d.) Thirst, a good deal, day and night, (a. 24 h.) Unusual, increased thirst. Violent thirst, tormenting. *Decrease of appetite.* Total want of appetite and sleep. *Increase of appetite.* Continued increase of appetite. Unusual hunger. Gnawing hunger, (a. 3 h.) Violent, craving appetite. Voracity. * Canine hunger. She would like to eat again shortly after a meal; she feels much better after having eaten a good deal. He feels an anguish, unless he eats every 3 or 4 hours; but he must take care not to eat too much. The digestive functions are carried on with more rapidity and regularity, the stools being normal. Alternate canine hunger and want of appetite. *Very weak digestion.* Week feeling, and a feeling of sickness all over the body after dinner.

Gastric Symptoms.—Eructations. Eructations smelling of the medicine. Continued empty eructations, from morning till evening, as if every thing he eats or drinks were transmuted into air. Hiccough. Qualmishness in the pit of the stomach, every day, passing off after eating, with pressure on the stomach. **Nausea.* Nausea imme-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

diately after rising, with spasmodic pain in the stomach. Inclination to vomit. Inclination to vomit, in paroxysms, with heartburn, with sensation as of the stomach being spoiled. *Vomiting.* Violent vomiting. Incessant vomiting. Obstinate vomiting, occurring especially after eating. Continual vomiting with diarrhœa. Yellowish, saltish vomiting. *Vomiting of bile.* Increased secretion of bile. Increased secretion of the gastric and pancreatic juice. Gastric derangement with constipation.

Stomach.—Pains in the stomach. Pains over the stomach. Pain in the stomach, early in the morning, going off after an eructation. Most violent pain in the stomach. Excessive pain in the stomach and bowels. Tormenting pain in the stomach. Pains in the stomach with copious bilious evacuations. Violent pains in the stomach and œsophagus. which felt obstructed and did not brook contact. Pressure in the region of the stomach, increased after every meal. Fulness and distension of the stomach, with trembling and increased warmth in the abdominal cavity, and a movement taking place from that cavity to the surface as if sweat would break out. Tension in the stomach and abdomen, preceded by movements in that cavity. Spasmodic pain in the stomach. Spasms of the stomach, frequently recurring. Gnawing pain in the upper part of the stomach. Gnawing, corroding pains in the stomach. Stitches in the pit of the stomach, (a. 5 d.) Prickings in the upper part of the pit of the stomach. Pulsations in the pit of the stomach. Increased warmth in the region of the stomach. Burning in the stomach. Inflammation of the stomach in the region of the orifice. Small, linear ulcers in the stomach having corroded the mucous membrane, which was swollen near the orifice, and covered with a crust of coagulated lymph ;—in animals.

Abdomen.—Decrease of pain in the region of the liver and pit of the stomach. Pressure in the right hypochondrium. Pressure in the region of the liver painful to the touch. Pressure and stitches in the region of the liver. The left hypochondrium is hard and is intensely painful when being pressed upon. Single attacks of sore pressure in the left hypochondrium. Sharp stitches in the left

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

hypochondrium, as if from incarcerated flatulence. Pain in the epigastrium. Great painfulness of the whole abdomen, as if from inflammation of the omentum. Pains in the abdomen extending to the dorsal spine. Oppressive pain in the epigastrium, between the pit of the stomach and the umbilicus. Pressure in the abdomen, near the right hip. Frequent paroxysms of pressure in the abdomen, mostly when sitting, and gradually abating when stretching one's-self, several days in succession. Distension in the epigastrium, with sharp pressure as if from flatulence, in different places, beginning after dinner, during the whole period of digestion. * Big, distended abdomen; she was unable to lie in a horizontal position, lest she should suffocate. Painful tension of the abdomen, with unusual pressing. Drawing cramp-pain in the epigastrium, beginning in the pit of the stomach. * Colic. * Pinching and aching colic. Cutting in the umbilical region, with papescent stools. Violent cutting in the umbilical region, with desire for stool. Drawing and pressure in the region of the right kidney. Burning tearing in the left side of the abdomen, near the hip. Stitches in the side of the abdomen. Sharp stitches in the abdomen, on the left side. Pulsations in the abdomen. Frequent pressing and bearing-down towards the lower opening of the pelvis, with frequent diarrhœa-stools. The abdominal symptoms return, especially after having eaten something. The abdominal symptoms recur after eating, for days, weeks, and even months. Ascites. Pressure in the right inguinal region. Disappearance of the swelling of the inguinal glands, (curative effect.) The region of the hernial sac becomes inflamed from the truss, (6. d.) Rumbling in the side where the hernia exists, which may be felt and heard; during an inspiration. * Incarceration of flatulence in the left side of the abdomen. Pressing movements in the abdomen, beginning in the stomach and extending towards the abdomen, especially towards the os pubis, the bladder and the scrotum, rarely accompanied by oppression in the direction of the chest, sometimes with cutting. Frequent rumbling in the abdomen. Emission of flatulence and stool, with slight cutting in the abdomen. Increased emission of flatulence. Emission of

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

flatulence, smelling like rotten eggs. Increase of the peristaltic motion of the intestines.

STOOL.—Desire for stool, without any evacuation taking place; it takes place with great facility after taking some cold milk. Indolent stool. Difficult stool. * Constipation, sometimes lasting a good while. Obstinate constipation. Hard, knotty, dark-coloured stools. Irregular stools; at times constipation, at times diarrhœa. Irregular stool; alternate constipation and diarrhœa, with distension of the abdomen, rumbling and tormenting flatulence. Sometimes obstinate constipation, sometimes violent diarrhœa. Several whitish stools a day, more soft than usual. Increased stool. Four papescent stools a day, every stool succeeded by burning at the anus. *Copious*, papescent *stools*. Inclination to diarrhœa. *Diarrhœa*. Long and exhausting diarrhœa. Diarrhœa early in the morning. Violent diarrhœa, with colic. Frequent diarrhœa, with pressing and bearing-down towards the abdomen. Papescent stool, with cutting in the umbilical region. Copious discharge of a watery, foamy, whitish mucus, with pinching around the umbilicus, and pressure upon the vertex. Discharges of thick mucus, or purulent, part of the fæces being retained, like dysentery. Discharges of bloody mucus, fetid. Pressure in the abdomen, after the morning stool, which is rather hard than soft. Pressure in the rectum, in the evening, when in bed, (a. 36 h.) Violent itching of the rectum, as of ascarides. Small, forcing stitches in the anus, when sitting. Smarting as from excoriation in the rectum, after a good stool. Frequent smarting as from excoriation, itching and burning of the anus. *Burning of the anus, in the evening.*

URINARY ORGANS.—Obstinate retention of urine. She emits but little, red urine, (a. 48 h.) Scanty emission of a dark-coloured urine. Frequent desire to urinate, with scanty emission. Unceasing desire to urinate. Frequent micturition, with desire. Increased micturition. Copious and frequent emission of a light-yellow watery urine. Increased secretion of thickly urine, with dark sediment. Involuntary emission of urine, (a. 3 d.) Dark, turbid, sometimes milky urine. Dark, yellow-green urine. Ammoniacal odor of the urine. Smarting, corroding

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

urine, while emitting it. Itching smarting as if from excoriation, in the orifice of the urethra. Intensely painful cutting in the orifice of the urethra, between the acts of micturition. Prickings in the external orifice of the urethra, (a. 16 d.) Frequent intensely painful drawing in the forepart of the penis; it is impossible to distinguish whether the drawing is more in the urethra or the glans.

GENITAL ORGANS.—Gurgling tearing, on the right side and near the penis. Aching on the right side, near the penis. Violent itching of the glans. Itching of the glans, recurring frequently. Violent titillation of the glans and under it. Cutting drawing in the corona glandis. Frequent pressing and bearing down towards the testes. One of the testes is considerably drawn up to the abdomen. A painless swelling of the right testis becomes scattered, amidst violent itching and burning, and oozing out of a fetid humor, (curative effect.) An induration of the prostate gland becomes scattered, (curative effect.) Erethism of the genital organs. Increase of the sexual instinct. The erections take place slowly, (5. d.) Violent pressing in the hypogastrium towards the female organs of generation. Labor-like spasms in the hypogastrium. Speedy suppression of dropsy of the ovaries, (curative effect?) Induration of the uterus is speedily transmuted into cancer. Diminished induration of the uterus, (curative effect.) Diminution of the cancerous degenerations in in the neck of the uterus, (curative effect.) Hysterics of nubile girls. Cessation of the actually flowing menses. * Menses delay 8 days, with vertigo and palpitation of the heart. Menstrual irregularity. Increase of the menstrual flow. Menses unusually premature, copious and violent. Causes easily uterine hemorrhage. Violent uterine hemorrhage, lasting 4 weeks, in a plethoric female of 24 years. Cessation of a uterine hemorrhage, which occurred at every stool, together with cutting in the abdomen and pains in the loins and small of the back, (curative effect.) The menses are preceded by rising of heat to the head, with palpitation of the heart and tension in the neck, which became thicker. Disappearance of the usual troublesome precursory symptoms of the menses. Pains in the small of the back at the appearance of the menses, which took

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

place in proper time. Great weakness during the menses. Cessation of the pains and unpleasant symptoms accompanying the menses; they flow without any trouble. Palpitation of the heart after the menses. Complete cessation of the leucorrhœal discharge, (curative effect.) Complete cessation of an old leucorrhœa, which was most abundant at the time of the menses, made the thighs sore, and corroded the linen, (curative effect.) The corroding leucorrhœa becomes more and more mild and scanty, (curative effect.)

COLD, CATARRH.—Sneezing, without coryza, with immediate and considerable protrusion of the nasal mucus. Obstruction of the nostrils, (a. 28 h.) Frequent and dry coryza (especially in the evening;) it becomes fluent in the open air, with much discharge from the chest. Fluent coryza, with much sneezing. Fluent coryza like water. *Increased secretion of mucus in the nose*, for several days. A quantity of yellow mucus is expelled from the nose. Increased secretion of nasal mucus. Pain of the larynx. Pain in the larynx, with discharge of hardened mucus. Pressure in the region of the larynx, extending as far as the fauces, as if these parts were swollen. Aching and stitches in the region of the larynx and the sub-lingual glands, coming on frequently on the same day. Pressure in the larynx, which obliges one to hawk up a quantity of tough mucus. Increased pain of the œsophagus, when pressing upon the larynx. Contraction and heat in the larynx. Contraction and heat of the throat. Soreness of the throat and chest, when in bed, with wheezing in the throat and drawing pains in the lungs, corresponding with the beat of the heart. * Inflammation of the trachea. Roughness of the trachea, the whole day. Hoarseness. * Hoarseness in the morning. Hoarseness for more than two weeks. Hoarseness in the morning after rising, which obliges one to hawk up tough mucus. Hoarseness the whole day. The voice becomes deeper, and finally quite deep. Sensation, as if something were lodged in the larynx which can be hawked up, the whole day and evening. Intolerable tingling and tickling in the larynx, which can only be relieved by coughing and clearing the throat, with accumulation of water in the mouth early in the morning, when in bed. Violent clear-

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

ing of the throat, (2. d.) Frequent clearing of the throat, early in the morning. *Increased secretion of mucus in the trachea.* Increased secretion of mucus in the bronchial tubes. Increased secretion of mucus in the throat, with roughness of voice. The secretion of mucus extends as far as the Eustachian tube, a kind of soreness remaining behind in the throat. Discharge of mucus from the throat in the morning after rising, with soreness in the throat. Great accumulation of tough mucus in the throat, in the evening. Discharge of tough mucus from the throat, with pressure in the throat, as if something were lodged in it which he imagines he is able to swallow, in the morning. Rapid and permanent disappearance of a continual inclination to hawk up tough mucus from the trachea; this hawking is very troublesome, and accompanied with tingling and stinging in the larynx. (Curative effects.) Irritation, which occasions a cough. * Irritation and cough, brought on by a violent tickling in the throat. Cough, with exertion; she came near vomiting; the cough ceases after a discharge of mucus. Cough, with pressure on the chest and difficulty of breathing. Evening-cough. Frequent and dry cough in the evening. Dry, short, and hacking cough. Dry cough. Frequent and dry cough. A good deal of dry cough, with dyspnœa, pressure and burning of the chest. Frequently a deep and dry cough, excited by pressure in the chest. Deep and dry cough, with stitches in the chest. Occasional irritation with cough, accompanied by expectoration of tough mucus. Short cough caused by a tickling in the throat, accompanied by a thick yellow discharge, good appetite, and wretched appearance. Cough accompanied by expectoration of mucus; previous to the cough, one experiences a sensation of heaviness extending from the throat as far as the chest, aggravating the breathing. Rattling of mucus in the chest, with roughness under the sternum, and oppression of the chest. Discharge of mucus from the chest. Discharge of mucus, occasionally mixed with blood. * Discharge of mucus streaked with blood.

CHEST.—Iodine increases pulmonary consumption. Pulmonary consumption. * Violent difficulty of breathing. Aggravated respiration. Oppressed breathing. Oppres-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

sion of the chest. * *Asthma*, with pain during a deep inspiration, more violent and rapid beating of the heart, and smaller and more frequent pulse. Asthma, the respiration in the trachea being impeded for a fortnight. Difficulty of dilating the chest when taking an inspiration. Sensation as if he had to conquer a great obstacle in dilating the chest during an inspiration. * Want of breath. Suppression of breath. Suffocating catarrh. Feeling of weakness in the chest, and in the region of the heart. Pain in the chest. Sore pain in the chest, continuing in both sides during respiration and contact. Pressure, rather deep in the right side of the chest. Aching in the right half of the chest, aggravated by every inspiration, for one hour after dinner. Pressure on the chest and oppressed breathing, as after inspiring the vapour of sulphur. Oppression and burning in the middle and sometimes also in the sides of the chest, with dry cough. Sharp stitches in the middle of the right half of the chest, only during an expiration. Sharp stitches in the lower part of the right half of the chest, near the pit of the stomach, during an inspiration. Congestion of blood to the chest, with inclination to inflammation. Violent pulsations in the chest, and palpitation of the heart, increased by every muscular exertion to such an extent that she was unable to support herself for one minute without being near fainting; a quiet, horizontal position gave her the greatest relief. * *Palpitation of the heart*, which might have been heard at the distance of a few paces. Palpitation the whole day, until the moment of going to bed. Spasmodic palpitation of the heart; she feels it as low down as the umbilicus, but mostly in the pit of the stomach. The palpitation of the heart disappears entirely. ,(Curative effect.) Sensation as of the heart being squeezed together. Burning and stinging tension in the integuments of the chest. Tearing in the integuments of the right side of the chest. Increase of the mammæ, accompanied with decrease of the goïtre, in several females. Occasional dwindling of the mammæ. Dwindling, withering of the mammæ. Disappearance of the adipose tissue in the mammæ. Dwindling of the formerly full and round mammæ, so that two years after the use of iodine, every trace

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

of the mammæ was lost. Nursing a baby is accompanied with disagreeable consequences.

Back.—Periodically increasing and decreasing aching in the sacrum and os coccygis. Stitches in the small of the back, (a. 15 d.) Itching of the back, over the right hip. Spasms in the back. Stitches in the scapulæ, when lifting something, (a. 14 d.) Burning on the right scapula. Rheumatic clawing in the lower part of the neck, near the shoulder; it is apparently alleviated by an eructation, but it frequently returns afterwards. Rheumatic tension in the right side of the neck. Tearing in the right side of the neck. Tension in the outer neck. Constriction of the neck. Unpleasant sensation of constriction in the neck. The neck becomes larger by animated talking. Increase of the goïtre, it becomes more painful. *Enlargement and painful induration of the goïtre.* Pains and pulsations in the goïtre. Painful tension in the goïtre. Continual sensation of constriction in the goïtre. Enlargement and induration of the goïtre in the first days, followed by a decrease. Decrease of the goïtre, (curative effect.) *Permanent disappearance of the glandular swellings of the neck and nape of the neck. Disappearance of old, hard, or soft swellings of the thyroid gland, and likewise of large goïtres,* (curative effect.) Redness of the neck and chest, as if those parts were ecchymosed. Yellow spots on the neck.

Superior Extremities.—Induration of the axillary glands. *Disappearance of the glandular swellings in the axilla.* Drawing tearing pains in the shoulder, which is abnormally elevated, (2. d.) Rheumatic pain in the left shoulder. Violent stitches in the shoulder-joint, even when at rest. Pain in the external surface of the humerus, waking one from sleep and becoming aggravated by lying upon the arm. Rheumatic pains in the arms. Tearing pain in both arms, after performing some manual labor. Paralytic * weakness in the arms, early in the morning, when waking up, in bed. Subsultus tendinum in the arms. *Tearing in the elbow of the left side.* Pressure in the bend of the left elbow. Painful stitches in the right wrist-joint. Pain as from contusion, in the upper surface of the metacarpal bone of the index-finger, increased by

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

contact. Tearing in the metacarpal bone of the right index-finger. Heat in the hands. *Tremor of the hands.* Slight tremor of the hands. Violent tremor of the hands, so that he cannot use them except at certain hours of the day. Zigzag movements of the hands. Subsultus tendinum in the hands. Round, burning-itching spot on the right hand, between the thumb and index-finger, covered with two whitish vesicles; rubbing made it feel easy; it went off on the third day. Tensive pain in the joints of the fingers, when bending them, as if they would break; with some swelling when stretched, and painfulness when pressing upon them, (after a few days.) Tearing in the posterior joint of the right thumb. Tearing in the knuckle of the right little finger. Tearing in the whole of the index and middle-finger of the left hand. Tremor of the fingers. Subsultus tendinum in the fingers. Panaris of the left index-finger, from a slight stitch in the finger, near the nail.

INFERIOR EXTREMITIES.—Pressure in the left half of the nates, apparently in the os ischium. Intermittent, sharp tearing between the left hip and the head of the femur, increased by moving the joint. Cramp-like sensation in the thighs and legs, only when sitting, not when lying down, walking or standing. Rheumatic drawing in the whole of the left lower limb, especially the thigh and knee, with humming in the heel, in the evening, when in bed; rather increased by motion. Heaviness of the lower limbs, as if they were made of lead. Swelling of the lower limbs. Swelling and trembling of the lower limbs. Œdematous swelling of the lower limbs. Uneasiness in the lower limbs. Trembling of the lower limbs, making walking difficult, and uncertain. Paralysis of the lower limbs. Pain of the thigh, especially at night, when in bed. Pains of a peculiar kind, in the thighs. Rheumatic pain in the left thigh. Clawing tearing in the left thigh, near the head of the femur. Sharp, lancinating tearing in the middle of the thigh, towards the inner side. Twitchings of the muscles of the thigh. Soreness of the thighs in a female, where they rub against one another in walking. Tearing in the left knee. Dull tearing in the outer side of the bend of the right knee. Disappearance of a white

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

swelling of the knee, (curative effect.) Pain of the leg, along the tibia, as if from sub-cutaneous ulceration. Tearing in both sides of the leg, close above the ankles. Violent spasm, with twitchings, in the ankle-joint, at night. Spasms in the feet. Aching cramp-pain in the foot, from the middle toe to the tarsus. Sharp, continued pain in the inner half of the right heel. Single, sharp stitches in the ankles. Weight of the feet, as if they were made of lead. Swelling of the feet, followed by rapid emaciation of the same. Œdematous swelling of the feet. *Subsultus tendinum in the feet.* Acrid sweat of the feet, corroding the skin. Lancinating tearing under the nail of the left big toe. Pain in the corns.

General Symptoms.—Erratic pains in the joints. Rheumatic pains in the trunk, neck and arms. Numb feeling in the thighs and legs. Numb feeling and trembling in the limbs. Paralysis of the limbs. Liability to cold, bringing on ophthalmia. Stinging itching in different parts of the body. Sensation as of violent flea-bites over the whole body, day and night. Itching of the cicatrix of an ulcer upon the thigh which had been healed for years past. Itching pimples on the old cicatrix. Small, red, dry pimples on the arms, chest and back, itching in the beginning. Dirty-yellow color of the skin, from 5 to 6 weeks. Rough, dry skin. Glandular indurations. Disappearance of single glandular swellings in the nape of the neck, axilla and groin, (curative effect.) Excites the glandular system, the glands of the buccal cavity, also the stomach, liver, the pancreas, and promotes the process of secretion in these organs. Pulsations in all the arteries, at every muscular effort. Extremely troublesome pulsations in all the larger trunks. Easy erethism of the arterial system, with obtusion of the head and subsequent headache. The circulation is easily excited and accelerated. Erethism of the vascular system, especially the veins. Orgasm of the blood. Violent orgasm. Inclination to hemorrhage. Hemorrhage from different organs. Great irritation of the nervous system. Increased sensibility of the whole body. Impatience in all the limbs. Uneasiness and nervousness, with trembling extending from the region of the stomach to all the parts of the peri-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

phery, as if he would begin to tremble, or to sweat all over, with an increase of warmth all over, as if coming from the stomach; afterwards pressure in the region of the heart, heaviness on the chest, with difficult breathing, and acceleration of the pulse by 7 or 8 beats.

Spasms, Fainting Fits.—*Trembling. Trembling of the limbs.* Trembling of the limbs, especially the hands. Trembling of the fingers and eyelids. Trembling of a peculiar kind. Trembling, first of the hands, then of the arms, feet and back, his walk is vacillating and unsteady; his hand moves in zigzag, he is unable to approach any thing to his mouth in a straight line; when at rest, the trembling parts may be held still easily: general exercise is painful and the circulation accelerated, with a small, wiry pulse. *Spasms.* Spasmodic movements of the limbs. Violent spasms and convulsed movements of the arms, back and feet, ceasing scarcely a moment. *Subsultus tendinum.* Subsultus tendinum in arms, fingers, and feet. Catching at flocks. A kind of Asiatic cholera. Difficult, vacillating gait, from trembling of the limbs. Heaviness in the limbs, early in the morning, (8. d.) Great lassitude, in the afternoon, after a short walk, with a sensation as if one had not yet taken breakfast, but without any hunger. Weakness, she sweats when talking. *Weakness.* Complete prostration of strength. Out of humour, weary and slow to move. Weak feeling. *Weakness of the muscles.* Long weakness of the muscles. *Decrease of strength. Loss* of strength. *Great prostration of strength.* Inclination to fainting fits and spasms. Fainting fits. *Emaciation. General emaciation.* Sudden, excessive emaciation. Excessive emaciation; her arms and body were almost without any flesh, her breast was quite flat, her calves had disappeared, and her thighs were as large as her fore-arms used to be in her healthy days. Emaciation lasting one year, until one does not know him any more, with general muscular weakness and unusual appetite. Chronic emaciation, pale face and muscular weakness. Emaciation with hectic fever. Return of embonpoint and cheerfulness, (curative effect.) Increased embonpoint, (curative effect.) Disappearance of a remarkable phlegma in a girl; she becomes stronger

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

and more lively, and the movements of her muscles become more easy, (curative effect.) Swelling and pain of the affected parts. Disappearance of a fungus articularis which had come on in consequence of measles, (curative effect.) Œdematous swelling of the whole body, which did not pass off. *Dropsical swellings.* Anasarca. Anasarca and ascites. Apoplexy. Death, with either local, or general inflammation.

SLEEP.—No desire for sleep, (6. d.) Restless sleep. Restless sleep, with anxious dreams. *Sleeplessness.* Sleeplessness for eight days; she did not sleep a wink. Sleep full of dreams. Vivid dreams, from which he would like to rouse himself; but is not able; with feeling of weakness after waking up. Dreams which he is unable to recollect, with sound sleep. Anxious dreams. Dreams about dead persons. She dreams every night: of swimming in the water, walking in the mud, of her daughter having fallen into a well. Cold feet, at night. Cold skin. He is frequently shaken by unusual chilliness, even when in a warm room; he does not feel well during the day. *Increased warmth of the skin. Increase of animal heat.* Flying heat. Fever, where chills alternated with flushes of heat. *Feverish sensations.* Fever with delirium, and subsultus tendinum. Fever with dryness and coldness of the skin, soft, quick pulse, delirium, subsultus tendinum and grasping at flocks. Quartan fever. Pulse strong, large and full. *Quick, hard pulse.* Accelerated, increased pulse. Quicker pulse, increased by 15 beats. Pulse increased to 86 beats. Increased, full, hard pulse. *Small wiry, accelerated pulse.* Small, hard pulse, and so quick that it was scarcely possible to count the beats. Small, very frequent, compressed pulse. Soft and quick pulse. The beats of the heart and the pulsations of arteries are synchronous; the pulse was small and weak, and not much accelerated, when at rest; at every motion the quickness of the pulse and the beats of the heart are synchronous. Night-sweat. Profuse night-sweat and little good sleep. In the morning she wakes up with sweat, and afterwards feels weaker. Sourish night-sweat, all over, every morning; with great faintness in the lower limbs during the first hour.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

KALI CARBONICUM.

Dynamise one drop of the carbonate of potash in the usual fashion. Kali has been found most useful in the following affections :—

Out of humor ; *tendency to start ;* want of memory ; dizziness ; unpleasant sensation of dulness of the head ; vertigo, apparently from the stomach ; headache when riding in a carriage ; headache when sneezing or coughing ; headache early in the morning ; headache with nausea ; aching in the occiput ; congestion of blood to the head ; humming in the head ; great liability of the external head to cold ; scurfy formation about the head ; dryness of the hair on the head ; falling off of the hair ; morning-sweat on the forehead ; swelling of the eye ; agglutination of the eyes early in the morning ; lachrymation of the eyes ; spots hovering before the eyes ; *the eyes are dazzled by the light ;* stitches in the ear ; stitches in the ears from within outwards ; weak hearing ; dull hearing ; ulcerated condition of the internal nose ; dulness of smell ; heat in the face ; flushes of heat in the face ; yellow complexion ; bloated condition of the face ; drawing pain in the face ; toothache when eating ; stinging toothache ; accumulation of mucus in the palate ; *hawking up of mucus ;* sour taste in the mouth ; spoiled taste ; bitter taste in mouth ; canine hunger ; great inclination to eat sugar ; frequent eructations ; sour eructations ; nausea ; inclination to vomit, with anguish ; nausea when eating ; *repletion after eating ;* tension across the stomach ; pinching in the stomach ; pressure in the liver ; pain in the liver, when stooping, as if it had been sprained ; pressure in the abdomen when stooping ; heaviness and uneasiness in the abdomen ; want of action and coldness in the abdomen ; labor-like colic ; accumulation of flatulence ; *incarceration of flatulence ;* want of emission of flatulence ; constant emission of flatulence ; inactivity of the rectum ; difficult expulsion of the fæces, on account of their large size ; constipation ; constipation every other day ; costiveness and difficult expulsion of the fæces ; mucus with the stool ; anguish previous to stool ; varices of the anus ; *itching of*

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

the anus ; aching in the rectum, previous to the emission of flatulence ; desire to urinate ; frequent micturition, day and night ; suppressed sexual instinct ; want of sexual instinct ; (sur-excitation of the sexual instinct ;) want of erections ; want of pollutions ; (too frequent pollutions ;) swelling of the testes ; weakness of the body, especially of the eyes, after an embrace ; (suppression of the menses, with anasarca and ascites ; *menses are too weak ; menses too early ;* itching eruption and soreness between the lower limbs during the menses ; *acridity, itching and corrosion* about and in the *genital organs ; leucorrhœa.*

Dryness of the nose ; obstruction of the nostrils ; *dry coryza ;* hoarseness ; cough ; *nightly cough ; purulent expectoration when coughing ;* purulent expectoration ; spasm of the chest when coughing ; difficult breathing ; wheezing in the chest ; asthma in the morning ; *asthma* when walking a little faster than usual ; spasmodic asthma ; spasm in the chest ; palpitation of the heart ; palpitation of the heart and orgasm of the blood, early in the morning when waking ; pain in the small of the back ; pain in the small of the back from a fall ; drawing pain from the small of the back to the middle of the back ; drawing pain in the back ; stiffness between the scapulæ ; *stiffness of the nape of the neck ;* weakness of the muscles in the nape of the neck ; goïtre ; pressure on the shoulders ; the arms go to sleep ; the upper arms go to sleep ; want of strength in the arms ; weakness of the arms, early in the morning, when in bed ; stiffness of the elbow-joint ; trembling of the hands when writing ; paralytic pain in the wrist-joint ; convulsed twitching of the fingers when sewing ; tearing with pressure in the thighs and legs ; *nightly tearing in the lower limbs ;* Creeping shuddering over the tibiæ ; cold feet, in the evening when in bed ; stiffness of the tarsal joint ; swelling of the legs ; burning pain in the legs and feet ; cold feet ; *fetid sweat of the feet ;* stitches and burning in the ball of the big toe ; corns painful to the touch ; drawing pain in the limbs ; contraction of the fingers and toes ; the limbs go to sleep ; parts are easily strained by lifting ; tremulous weakness ; weak condition after confinement ; *liability to cold ; want*

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

of perspiration, inability to perspire; anasarca and ascites; red, itching, burning patches on the body; herpes; old warts in the face; drowsiness in day-time; early drowsiness in the evening; sleep full of fancies; anxious sleep, full of dreams; frightful dreams; twitchings when asleep; inclination to shudder, in day-time; heat, early in the morning when in bed; great inclination to sweat when walking; *night-sweat.*

Persons suffering with ulceration of the lungs, can scarcely get well without this anti-psoric. Nitric acid is frequently indicated after Kali.

This drug has been proved by Drs. Hahnemann, Gersdorff, Goullon, Hartlaub, Rummel, and by Ng.

Noack and *Trinks* furnish the following clinical remarks on Kali. Dropsical symptoms in the case of a very old female. Paralysis, (also Causticum.) Humid tetters in the hands, with itching and stinging. Scirrhous induration. Chronic headache, over the eyes, moving towards the nose, with vomiting and eructations. Stitches in one side of the occiput, only during motion; also at night. Sarcoma of the eyelid, (with Silicia.) Scrofulous, serous, or purulent discharge from the ears. Angina parotidea. Phthisis hepatica. Lancinating pressing proctalgia in pregnant females. *Pressing and bearing down sensations of pregnant females, as if a load were falling into the pelvis.* Menstrual affections, especially in entire suppression of the menses, or when the menses delay making their first appearance in the years of puberty. The usual consequences of miscarriage, especially when badly attended to: weakness in the back, weakness of the lower limbs, dry and hacking cough, long-continuing sweats, feverish chills, chronic-inflammatory condition of the uterus, with nausea and vomiting; it is a preventive against miscarriage, for those females in whom it generally takes place between the second and third month; weakness after confinement. Scrofulous leucorrhœa. Cough, with more or less difficult expectoration of white mucus and shortness of breath when ascending an elevation. Cough with expectoration of bloody pus and sensation as of a lump in the stomach. Cough with yellowish expectoration, want of breath when walking and in bed, (has to

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

rise,) morning-sweat, diarrhœa, lancinating pain in the sternum, (vexation brought back both the cough and asthma, relieved again by *Phosphorus*.) Night-sweat and some cough, with hectic disposition. Hacking cough, with symptoms of incipient phthisis. Hæmaptoë. Wurm thinks that Kali carb. may be advantageously employed in pleuritis of tuberculous subjects; according to Goullon, it merely relieves the temporary pleuritic symptoms in tuberculous subjects. Pleuritic affections of every degree, no matter whether it is a case of pure pleuritis or pleuropneumonia, (preceded by Aconite, according to Goullon;) pneumonia, when the violent lancinations and the inability to breathe deeply continue or return after the administration of Aconite, and the accompanying cough is dry and suppressed, "*hence in genuine pleuritis.*" Incipient pulmonary phthisis, with occasional discharge of tuberculous masses, darting stitches through the chest, and exhausting, dry cough, great weakness and emaciation; it is therefore useful in tuberculous phthisis, when the tubercles become dissolved, (together with Lycop.;) purulent pulmonary phthisis, (giving first Acon. and Nux v.;) against bags of pus in the lungs Kali is useful during an aggravation of the symptoms in the fall-season, especially when there is fever, pain in the chest, lying down, &c. Dr. Libert found it little useful in tuberculous pulmonary phthisis. Mucous phthisis, with violent cough, mucous, yellow expectoration, especially early in the morning, sore pain, occasional stitches in the chest, especially under the short ribs of the right side when touching the side, or when breathing deeply; short breathing, drawing pains under the short ribs of the left side, when lying upon the right side, sometimes tearing pains in the left half of the head, preceded by turning vertigo, nausea and inclination to vomit, tearing pains in the legs, feet and knees, (especially when in a state of rest;) frequent chilliness and uneasy sleep after midnight. *Violent pains in the small of the back, in pregnant females.*

ANTIDOTES.—Camph., Coff., Spir. nitr. dulc., Mosch., Zinc., Opium.

MORAL SYMPTOMS.—Great despondency, without anxiousness. Want of energy. Sad, weeping mood, conse-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

quent upon exercise in the open air. Sad, lonesome; she seeks company to cheer up her spirits. Great sadness; she weeps without knowing why, in the evening. Weeping mood; she felt as though she could shed torrents of tears, (a. 20 d.) She wept a good deal, because she imagined all the time she would have to die. Anguish, causing a flow of tears, (1. d) Anguish and great sadness. A sort of anguish, and aversion to society. Anxiousness every day. Apprehensiveness. He is apprehensive of not being able to get well. Anxiety on account of her illness. Mental anxiety in the evening. Sad ideas about the future. Uneasiness of the mind. Precipitate resolution and action. Irresoluteness. Despondent and pusillanimous in the highest degree. Fearfulness, in the evening, when in bed. Fearful when alone. *Great tendency to start.* Tendency to start, especially when one touches her. She starts with a loud cry on account of an imaginary vision, (such as a bird flying to the window.) Very low-spirited, in the evening when falling asleep, and in the morning when waking up. Out of humour, without knowing why, (5. d.) Very much out of humor; one can perceive this by his looks even before he himself is aware of it. Out of humor, as though she could not do any thing to her satisfaction. She is constantly in a state of antagonism with herself, knows not what she wants, and feels very unhappy. Contrary mood; he is obstinate, and frequently knows not what he wants. Contrary mood; she demands things with impetuosity, is not satisfied with any thing, is beside herself, furious if her wishes are not strictly complied with; she frequently does not know what she wants. He is impatient about his children. Irritable, as after vexation. Sensitive and irritable. *Irritated mind. Vexed and irritated mood.* He is easily irritated. Excessively irritated mood. Nothing gives her any pleasure, she is very much out of humor. Vexed and peevish mood; every trifle vexes him and every noise is disagreeable to him; worst at noon and in the evening. Angry and wrathful, early in the morning after waking, he grinds his teeth, (a. 4 d.) Easily angry. Vehement. Indifferent towards every thing. Dread of labor. Changing mood, at times kind and calm, at times

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

vehement and angry about trifles; now full of hope, now desponding. Absence of mind; he finds it difficult to direct his thoughts to one fixed object. Want of presence of mind; he is at a loss how to begin what he intends to do, (a. 15 h.) He is frequently at a loss to find the proper expression, and chooses a wrong one. Delirium, day and night. Inability to collect one's senses, apparently seated in the occiput, during much talking, passing off when closing the eyes. Loss of consciousness for a couple of minutes, so violent that he would have fallen over, if he had not held on to something, (a. 18 d.) Sensation as if she lost her thoughts for moments.

HEAD.—Occasional sensation, as if her thoughts and memory had vanished, with buzzing in the head. Dreary and stupid feeling in the head, in the evening. Feeling like intoxication, (a. 4 d.) *Frequent obtusion of the head.*

Obtusion of the head, as after intoxication, and as if the ears were stopped up, with nausea, almost unto vomiting, (a. 8 d.) Frequent obtusion and heaviness of the head, early in the morning, in the region of the eyebrows. Obtusion of the head, as if screwed in, with stitches in the brain, at frequent intervals. Gloominess in the head, as if he had not slept enough, early in the morning, after rising, with cloudiness and want of cheerfulness. Weakness of the head. Weakness of the head, over the eyes, after fast walking, (17. d.) Vertigo when walking, like intoxication, he staggers from side to side. Vertigo, he staggers when walking or standing; better in the open air. Vertigo, especially after a meal. Vertigo when rising, as if her head were too light; she had to hold on to something. Slight attack of vertigo, as often as he rises from a chair and turns himself. Vertigo, with sensation as if a hollow were behind him, and he would fall into it, when *turning* about after looking in a looking-glass and after reading. Vertigo when turning about. *Vertigo when rapidly turning the head or body. Sensation of vertigo in the head, mostly early in the morning and evening.* Very giddy, even when sitting, (a. 30 h.) Vertigo when sitting, sensation as of staggering to and fro, (previous to dinner.) Vertigo when sitting; he dares not get up from his chair lest he should fall. Vertigo when writing and in the open air, every

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

thing turns around him. Headache, relieved by sitting in the bed, aggravated by lying down. Headache, early in the morning when waking up, for a quarter of an hour, several mornings. Headache in the top of the head when pressing upon it. Horrid headache, through the eyes. Attacks of megrim, on both sides, with faintishness almost to nausea. Violent pain in the whole of the head, with throbbing and stitches in the knees, going off by motion; in the evening. *Aching in the head. Aching in the forehead, in the afternoon, when walking, being at the same time out of humor*, (13. 19. 20. d.) *Pressure in the forehead, with photophobia.* Violent pressure over the whole skull, down the nape of the neck, throbbing in the head and the whole body; the pain does not admit of the slightest contact, and increases in paroxysms with violent nausea and vomiting of bile. Pressing aching in the forehead, with vomiting of mucus and acidity. Pressing and aching in the right temple, (11. 19. 20. d.) Pressure in the right temple, from morning till noon, (a. 11 d.) Aching in the left temple, (6. d.) Aching in the forehead, in the evening, when going to bed, with squalmishness, as if he had spoiled his stomach; alleviated by rest, aggravated by walking. *Pressure over the eyes* with violent pain in the whole forehead. Aching in the occiput, towards the nape of the neck, going off in the open air. Violent aching in the whole of the head, with chills over the whole body, especially in the forenoon. Violent pressure in the occiput, with orgasm of the blood in the head, and a feeling of heaviness when standing. Pressure and drawing tearing in the forepart of the head, extending as far as the eyes and the root of the nose, (14. 17. 18. 21. d.) Violent pressure and drawing in the forehead, in the evening. Pressure and burning deep in the occiput, with heaviness of the head so great that he threatened to fall over. Pressing headache, from the temples to the centre. Pressure on the top of the head, in the evening. Pressure into the right side of the cranium, when rising after stooping. Stitching pressing into the left temple. Boring pressure from without inwards over the left eye. Pain in the right temple pressing from within outwards. Violent pressing from within outwards in the whole forehead,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

when writing. Violent pressing from within outwards over the left eye, as if the brain would fall forwards. Sensation in the forehead, as if the fore part of the head would become dashed to pieces, in frequent and short paroxysms. Feeling of fulness in the head, as if the brain were pressing against the skull. Heaviness in the occiput, as if it were full of lead, the head falls backwards; accompanied by stiffness in the nape of the neck extending as far as between the scapulæ. Heaviness in the occiput, it feels obtuse. Heaviness and painfulness of the forepart of the head. Feeling of great heaviness in the left half of the head. *Clawing pain in the left temple, in paroxysms; also tearing. Drawing in the forehead, in the forenoon and after midnight*, (2. and 30. d.) Drawing and tearing on the top of the head, (33. 34. d.) *Tearing drawing in the left half of the head*, over, in front of, and in the temple, (12. 19. 25. d.) Drawing in the occiput and nape of the neck, especially on the right side, with stiffness. Tearing in the left frontal eminence, (25. d.) Tearing extending from the left temple to the articulation of the jaw, in the evening. Tearing in both temples, also in the left parietal bone. Tearing, now in the right, now in the left side of the occiput, at times also in the forehead, (1. d.) Throbbing tearing in the right side of the occiput, close to the nape of the neck, (16. d.) Darting-tearing pain in the head. Darting in the left temple. Darting pain in the head, the whole day. *Stitches in the temples*. Stitches in the temples, which cause one to start and to scream, in the right temple they are accompanied with tearing. Stitch over the left temple, and immediately afterwards a stitch in the middle of the forehead from within outwards. Prickings in the forehead. Stitches in the forepart of the head. Violent stitches in the forehead, the whole day, sometimes also in the left side of the head; with violent pains in the chest and icy coldness of the limbs. Stitches in the forehead, early in the morning. Stitches in the upper part of the forehead and over the temples, when moving the lower jaw. Stitches extending from the nape of the neck into the occiput. Stitches in the occiput, when setting the foot down and when stooping, apparently in the surface of the brain.

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

Stitch through the right side of the head, from behind forwards. Stitches through the whole head. Dull stitches in the head, (1. d.) Throbbing and beating in the forehead and especially in the sides of the head, frequently intermitting; also, after dinner, when standing or walking. Throbbing pain in the forepart of the head. Painful throbbing in the head, when she begins to write. Beating throbbing) pain in the upper part of the left side of the head; the pain becomes more violent and lancinating, when pressing upon the part; rather externally. Searching, (digging) throbbing in the frontal bone, over the left eye. Ulcerative pain in the head, after dinner; she had to lie down, this alleviated the pain. Congestion of blood to the head, with intoxication arising from it. Warm rising of the blood to the head, with orgasm of the blood in the body, immediately; a few hours afterwards a slight headache. Great warmth in the head, especially on the right side of the face, frequently, (5. d.) Rising heat in the head, in the evening, previous to lying down, going off when in bed. Burning, painful feeling of heat in the head. Sensation in the forehead, as of a hot body having descended into it, frequently recurring when stooping or writing, going off when raising the head. Sensation, when stooping, as if something were descending into the forehead from the occiput. Painful feeling, as of something moveable in the head, worse when moving the head Continued feeling in the head, *as of something having become detached in it*, and turning and winding towards the forehead. Painful turning and winding in the head. Concussive headache. Humming, groaning, in the right temple. Creeping pain over the forehead. Sharp aching in the outer side of the temple. Painful stitches about the head and in the nape of the neck, with stinging swelling of the cheeks and stitches in the teeth. Single, tearing stitches, in the left temple, extending to the zygoma. Fine stitches about different parts of the head. Painless, gurgling jactitation of the muscles in the right temple. The head was several times pulled to one side with a jerk, without inability to collect one's senses; afterwards the nape of the neck became stiff. Chilliness about the head. Liability of the head to cold, with headache and toothache.

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

Itching of the hairy scalp. Frequent itching of the head, especially the occiput. Itching of the hairy scalp, with sore pain when scratching it. Pimples on the hairy scalp. Large red pimple on the left frontal eminence, painful to the touch, and suppurating afterwards, (a. 32 d.) Painful tumor on the right side of the head, as if an ulcer would form, (a. 6 d.) Large, yellow, scaly spot in the upper part of the forehead. Dryness of the hair of the head. *Falling off of the hair.*

Eyes.—The eyes are painful to the touch. Pain of the left eye when looking upwards. *Pressure in the eyes.* Pressure upon the eyelids. Pressure upon the eyes, and in the orbits, with drowsiness, at noon, (36. d.) Pressure in the eyes and dry pus in the lashes. Pain, as if the eyes were being pressed in. The eyes are painful when reading, as if they were being pressed in. Pinching in the eyes. Tearing in the left eye, in the evening previous to going to sleep. Sharp *tearing in the right orbit, and in the eye, at night*, (30. 31. d.) Tearing, with pressure, in the interior of the right eye, (12. 26. d.) Tearing, with pressure, in the region of the right eyebrow, (26. d.) Pulling with a jerk, or tearing in the eyelid and over the right eye. *Stitches in the middle of the eye.* Stitches in the ball of the eye. Stitches in the right eye, (a. 21 d.) Stitches in the outer canthus of the right eye. Boring pain in the eyes. Pain in the left eyebrow, as if an ulcer would form, in the evening when in bed, (8. 13. d.) Itching of the eyes. Itching of the margin of the right eyelid. *Smarting pain in the eye*, (a. 4 d.) Sore feeling in the eyelids, shortly after midnight, when waking up, (25. d.) Biting and shooting pains in the eyes. The boy complains of coldness in the eyelids. Both eyes are very hot to the touch. *Burning in the eyes.* Burning in the eyelids. Burning in both eyes. Burning and biting sensation in both eyes. *The white of the eyes is red, the capillaries are injected.* Redness and heat of the eyes. Inflammation of the white of both eyes, with a burning pain, (a. 5 d.) Inflammation of the lids of the right eye, with pain of the eyes and inability to read by light. Swelling of the right eye. Great swelling of the upper eyelid towards the nose. Swelling between the eyebrows and lids, look-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

ing like a little sack. Swelling of the gabella between the eyebrows, (21. d.) Pimple in the left eyebrow. Frequent soreness of the external canthus of the left eye. Soreness of the outer canthus, with burning pain. The corners of the eyes ulcerate. Agglutination of the eyes, early in the morning, (a. 16 h.) Lachrymation, (2. d.) Lachrymation of the eyes, especially the right, with a biting sensation in the canthus, (27.d.) Frequent lachrymation, and rays around the light of the candle in the evening. Dryness and burning of the eyes, worse in the open air, than in a room. Dry feeling in the eyes, (a. 2 d.) Dry feeling, as of sand, and great drowsiness in the eyes. Her eyelids are violently closed. The eyelids open with difficulty, early in the morning, when waking. Twitching and jerking in the right eyebrow. Staring look, she finds it difficult to turn her eyes from an object, and has to fix them upon it almost against her will. Vanishing of sight when writing, with white little stars before the eyes; the lower line looks as if it were above the upper, so that he finds himself constantly writing upon this one. The eyes feel painful and weak. Weak sight. Obscuration of the right eye, early in the morning, for some minutes. Diminution of sight, after working in water; she only saw a small portion of the objects, and afterwards felt stitches in the head above the eyes, with nausea. Black points and curls in front of the eyes, when reading. Spots, gauzes, and points before the eyes, when reading or looking in the open air, (a. 24 h.) A black little globule is hovering before the eyes. When looking at snow, there is a sensation as of white drops falling down before the eyes. Various colours before the eyes. Blue and green spots before the eyes. Yellow, shining, tremulous fog before the eyes. Wheels before the eyes, with yellow and white rays, when writing upon paper and in the open air; they turn about in a circle, and become larger and larger. Bright sparks before the eyes. Sparks rush out of the eyes when coughing. Photophobia: painful sensitiveness of the eyes to the light of day; the room has to be made obscure.

EARS.—Pain in the ears. Straining pain in the right ear. Stitches and straining in the ears, (3. d.) Claw-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

ing sensation in the left outer ear. Drawing pain in one, then in the other ear, (a. 4 d.) *Tearing in the ear.* Frequent tearing deep in the right ear, (1. d.) Alternate tearing in both ears. Tearing in the right concha, (24. d.) Quick tearing in the left ear, and around it, apparently in the bone. Frequent tearing in the anterior margin of the right ear, (1. d.) Painful tearings from the interior of the left ear into the outer cartilage, and at the same time in the bone, above and below the right patella. Violent tearing in and behind the ear. Jerking pulling behind the right ear. Darting behind and above the ear. Stitches in both ears, in the evening, when in bed. Sharp darting pain behind and above both ears. Sharp stitches into the left ear, which caused her to start, early in the morning. Frequent fine stitches in the left ear, from within outwards, (13. d.) Continued, dull stitch in the left ear, passing off by shaking the head. Stitches and creeping in the interior of the ear, connected with a similar sensation in the stomach and œsophagus, (30. d.) Boring and aching in the ears, (1. d.) Gnawing in the inner and outer left ear. Ulcerative pain in the outer right ear, continuing a long while, before midnight, (3. d.) Throbbing in the right ear, at night, when resting upon it, (a. 2 d.) Sense as of hammering in the right ear, frequent and very disagreeable, hindering hearing. Twitchings about the left ear, (a. 10 d.) Twitchings and trembling in the right ear, when rising after stooping. Itching of the lobule. *Violent itching in the ears*, (a. 4 d.) *Tickling in the ears.* Coldness of the ears in a hot room, (a. 2 d.) Heat of the lobules. Sensation as if warmth were rushing out of the left ear. Redness, heat and violent itching of the outer ears. Inflammation and swelling of the internal ear, with pain round about, (a. 3 d.) Soreness and suppuration behind the ears, for 4 weeks, (a. 21 d.) Pimples on the ears. Discharge of a yellow, liquid cerumen or pus from the ear, with previous tearing in the ear. Secretion of fetid humor in the internal ear. Opening of an ulcer in the ear, (a. 5 d.) Hard swelling of the parotid gland, with pain to the touch. Sensation of obstruction of the ears. One of the ears feels suddenly obstructed, (a. 3 d.) Sudden closing of the right ear, in the evening, when sitting,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

with sense as of tingling and buzzing in the left ear, so that his head vacillated a little. Dulness of hearing, in the evening, (a. 15 d.) Diminution of hearing in both ears, slowly increasing and decreasing, (for a fortnight.) Tingling in both ears. Violent tingling in one, and humming in the other ear. *Singing in the ears.* *Whizzing in the ears.* Violent *roaring in the ears.* Ringing in the ears. Frequent reports and detonations in the ears, during the day. *Cracking in the ear*, frequently. Cracking in the ear, during a violent expiration. Guggling in the right ear, and secretion of a quantity of soft wax.

Nose.—Clawing sensation in the nose and root of the nose, especially on the right side, (23. d.) Itching in the nose. Frequent itching in the right nasal fossa. Ulcerative pain in the right nasal fossa. Great burning in the nose. Burning in the left nostril. Burning and biting sensation in the upper part of the left half of the nose, extending to the cribriform plate, (23. d.) Red, thick nose, especially in the afternoon. Considerable swelling of the tip of the nose. Red and hot nose, bordered with a number of white pimples. Pimples on the nose. Flat, small ulcer above the left wing of the nose, painful to contact. Pimple in the left nostril, (a. 5 d.) Sore scurfy nostrils, for a long time. Ulcerative condition of both nostrils. Ulcerated nostrils. When blowing her nose, something fetid comes out of it. Bloody, red nostril, every morning. *Frequent bleeding of the nose.* Bleeding of the nose, early in the morning. Sensitive smell.

Face.—Pale countenance, and weakness. Wretched complexion, with pale lips. Blue margins around the eyes. Pale, hollow-eyed countenance, especially in the open air, where the child looks as if it were frozen. Pale countenance with faint, lifeless eyes. Heat and redness in the face, early in the morning, when in bed. Long-continued heat and redness of the face, with icy cold feet. Burning, red cheeks, in the evening, for an hour and a half; followed by great paleness of the face. Burning itching in the face. Itching of the skin of the face, preceded by twitchings; he had to rub it, after which the skin burnt like fire. Considerable swelling of the cheeks, passing over into an ulcer on the gums, without any pre-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

vious toothache. Swelling on the right cheek, low down, with stitches and pains to contact. Big cheek with tearing and stitches. Big red cheek, with small pimples, also on the nose. *Pimples in the face.* Pimples in the face, going and coming. Pimples in the face, continually. Pimples in the face, with pus in their tips. Pimples on the zygomata, with burning pain. Painless blotch on the cheek, below the ear. Small, red pustule in the centre of the forehead, passing off again next morning. Freckles in the face. Dry, parched skin of the face. Pressive drawing in the muscles of the cheek, near the lower jaw. Tearing in the left jugum, afterwards in the cheek. Tearing in the left jugum, alleviated by pressing upon it, with sensation as if the cheek were swollen, in the evening and night, until next morning; she wept and was unable to sleep.

Teeth and Jaws.—Tearing in the lower jaw and in front of the right ear. Clawing tearing in the right jugum, extending to the palate. Burning in the face, under the right eye. Twitchings in the left cheek, with fine, burning stitches; with tearing in the left temple from below upwards, in the evening. Spasmodic sensation in the lips. Tearing in the left upper lip and gums, going off by pressing upon the parts. Stitch in the upper lip. Burning of the lips. Burning of the lower lip. Itching around the margins of the lips. Smarting sore feeling around the mouth, in the margins of the vermilion border, very painful to contact. Soreness of the vermilion border of the lips; they are agglutinated early in the morning. Swelling and rhagades of the upper lip; it is painful to contact and bleeds readily. Thick, ulcerated lower lip. *Scurf on the upper lip.* Vesicles on the vermilion border of the lower lip, painful and itching when touched. Vesicles on the lips. Small, pointed, itching and humid pimples on both lips, and around the whole mouth. Pimples on the lips, smarting and itching. Pimple on the upper lip, painful when touched, near the left nostril, (36. d.) Peeling off of the lower lip, (34. d.) Chapped lips, peeling off. The lower lip peels off and becomes chapped and scaly. Clawing pain in the right half of the jaw, not far from the joint. Insufferable spasm, (cramp?) in the jaws, almost

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

closing the pharynx? at the same time. *Itching of the knee.* Pain of the sub-maxillary gland when touched. Swelling of the lower jaw and the sub-maxillary glands, with looseness of the teeth. The teeth are sensitive to pain, (4. d.) Toothache with pain in the face. One or the other tooth became loose and sensitive, or a bone of the face became painful, and, like the tooth, appeared to be pain itself; afterwards there were paroxysms of dartings and tearings in one point. Pain of the teeth, every day in the morning, when waking. Pain of the teeth, in the left side, early in the morning, in bed and the whole forenoon, (2. d.) Pain in the roots of the teeth of the left side, several mornings after waking, increased by eating, (3. d.) Toothache after a meal, in the direction of the cheek-bone and ear; here the person had a sensation as of a griping together and stinging. *Toothache when eating anything, not otherwise. Toothache only when eating*, a throbbing in all the teeth. The teeth are painful when taking water into her mouth. The teeth are painful when touched by either cold or warm substances. Frequent paroxysms of toothache, as soon as cold air gets into the mouth; improved by warmth. Toothache with subsequent swelling of the gums. Toothache, the pain feels as if it had been super-imposed from without, as if something had got into the hollow tooth; accompanied by drawing behind the ear and on the head, at last darting in the tooth and disappearance of the pain. *Toothache, only when eating*, noon and evening, frequently at the first mouthful, *as if something had got into the hollow tooth*, with insufferable drawing extending to the eyes and ears, in paroxysms, with intervals of half an hour. Aching in the root of one of the posterior molar teeth, in the evening. *Drawing toothache*, as soon as she gets into bed in the evening, not by day. Drawing in the roots of the anterior teeth and in the molar teeth of the left side, mostly in the evening. Contractive toothache in the upper and lower row. Darting and drawing in the tooth, as if it were being corroded, generally after a meal and at night, for some time. Tearing toothache during or shortly after a meal. Tearing and griping in a molar tooth and the zygoma of the left side, increased and excited by cold substances, relieved by tying some-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

thing around the parts firmly. Tearing in the teeth and the right jaw. Corrosive itching, violent pain in different teeth and the gums, which cannot be relieved by picking the teeth. Itching in the teeth, after supper. Itching and sense as of digging in one of the left upper molares, after dinner; relieved by pressing upon the teeth. Burrowing pain in the left lower row, occasioned by picking. Violent, burrowing pain in one of the upper left molares, after dinner. Boring aching, always after dinner, as if something had got into the tooth. Throbbing or pecking sensation in one of the right upper incisores, after dinner. Throbbing and pulling sensation in the teeth, during motion; otherwise, burning pain. Burning, lancinating toothache, especially at night, as if a hot iron were plunged into the tooth. Burning stinging in the tooth, especially at night, with internal chilliness, and swelling of the lower jaw and gums, (a. 32 d.) *Stinging in the teeth* and gums, afterwards swelling of the cheeks with stinging pain, (a. 14 d.) Violent stitches in the teeth. Single stitches in the foreteeth, here and there, in the evening. Single stitches in the teeth and frequent sneezing, early in the morning, when waking up. Stinging pain in the fore teeth, with sensation of dulness during supper, (32. d.) One of the teeth stands out and is very painful when chewing. Feeling of looseness in one of the upper left molares. Looseness of all the teeth. Bad smell from the teeth. Tearing pain in the gums just above the anterior incisores. Tickling of the gums, and bleeding of the same when sucking at them with the tongue. Increased redness of the gums. Painful inflammation of the anterior gums. Considerable swelling of the gums above the upper molares, with swelling of the left tonsil and the cervical glands, (9. d.) Ulcer on the gums. Soreness of the inside of the gums of the anterior teeth.

Mouth.—Fetid smell from the mouth, like old cheese. Corrosion of the inner mouth, and tooth, as if by something acrid. Soreness of the mouth. Dryness of the mouth wakes him from sleep early in the morning, (7. d.) Numbness of the mouth, as if it were burnt, early in the morning, after waking. Violent burning in the mouth, early in the morning, and thirst. Dryness in the mouth,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

early in the morning, after rising. Dry mouth, without any thirst, in the evening. Dry, viscid sensation in the mouth. Dry feeling and accumulation of saliva in the mouth, he has to spit a good deal. Accumulation of water in the mouth, constantly. A quantity of saliva flows from his mouth, also by day. Painful vesicles all over the inner mouth, with burning pain. Early in the morning, when waking, the tongue feels quite dry, and is almost without any sensation. White tongue with dryness, as if caused by something astringent, early in the morning. Burning of the tongue and lower lip. Burning of the tip of the tongue, as if it were raw, or full of vesicles. Swelling of the tongue, covered with a number of small painful vesicles. Painful vesicles on the tongue and gums. Painful pimple on the tip of the tongue. Soreness of the frænum linguæ. Soreness of the tip of the tongue.

PHARYNX AND ŒSOPHAGUS.—Itching of the palate, (10. d.) Stitches and smarting in the back part of the palate, as is felt when the parts are too dry previous to a cold setting in, increased by deglutition, morning and evening, (8. 9. 29. 30. 41. d.) Sore throat with difficult deglutition and opening of the mouth. Impeded deglutition in the pharynx. When eating, particles of food easily get into the windpipe. Pressure in the spine during deglutition. Difficult deglutition, the food descends very slowly in the œsophagus. Sensitiveness of the œsophagus, warm food causes a burning sensation in it. The food does not pass down; she is unable to swallow dry and cold things. Pressure and tearing in the pharynx, (9. d.) Pressure with anxiety in the throat. Sensation as of a lump in the throat. Sore throat on the left side; he feels a lump there, and stinging during empty deglutition. Stinging pain in the pharynx; when getting cold he feels as if he had a fish-bone in it. Elongation of the uvula, with stiffness of the nape of the neck. Sore pain in the throat. Sore pain in the throat, in the upper part of the palate, during empty deglutition, worse when swallowing food; the symptom is not felt between the acts of deglutition. Smarting sore throat when swallowing. Scraping sensation in the throat, (8. d.) Dryness in the back part of the throat. A quantity of mucus is frequently

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

in the throat, (the first 3 days.) A quantity of mucus in the throat especially early in the morning. *A quantity of mucus in the back part of the throat*, which it is difficult to hawk up. Increased hawking of mucus, (19. d.) Tenacious mucus in the back part of the pharynx, early in the morning, which it is difficult both to hawk up or to swallow, with continual sensation as if a lump of mucus were lodged in the throat, (16. d.)

TASTE AND APPETITE.—Loss of taste, early in the morning, when waking, only for a short while, (2. d.) Bad taste in the mouth. Bad taste and accumulation of mucus in the mouth. Repulsive watery taste in the mouth. Flat, viscid saliva in the mouth. Bitter taste in the mouth, with nausea, (1. d.) Bitter taste in the throat. Bitter taste in the mouth, early in the morning. Bitter sour taste in the mouth, after breakfast. Sour taste in the mouth, every day. Putrid taste in the mouth. Sweetish taste in the mouth. Taste as of blood in the mouth, early in the morning, after waking, for three hours. Little appetite. Little appetite, with flat taste in the mouth; nevertheless he relishes the food. Great hunger. Violent thirst, in the forenoon. Thirst in the evening, previous to lying down. Thirst, at night. No appetite, he does not relish what he eats. Food, especially meat, is repulsive to him; he has an appetite when he eats, nevertheless he cannot eat much. Aversion to brown bread. Aversion to every thing.

GASTRIC SYMPTOMS.—Milk does not agree with her. Bread weighs in the stomach, after a meal. Great desire for sour things. Paleness of the face, nausea, vertigo, eructations, weakness of the lower limbs, and coldness of the hands and feet, before and after dinner; still he has some appetite. Nausea unto vomiting, when eating roast fish. *Desire for sleep, during a meal*, (a. 2, 4 d.) Drowsiness and depression of strength after a meal. Great drowsiness, with chilliness and yawning, after a meal. Weariedness after a meal, with throbbing in the pit of the stomach and headache. Vexed, peevish mood during dinner, with drawing pain in the head, (30. d.) Constrictive sensation in the head, after dinner, as if it were encircled with a hoop. Pale face after a meal. Pinching and

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

uneasiness in the abdomen, after eating soup, at dinner or supper, or after eating warm biscuit in the morning. Pressure in the stomach, like a weight in the stomach, after a meal, and especially after breakfast. *Distension of the abdomen after a meal.* Repletion and great distension of the abdomen, after eating but little. Oppressive flatulent colic after breakfast, relieved only for a short time by the emission of flatulence, (14. d.) Dull stitches in the right side of the epigastrium, after dinner. Tickling, exciting a cough, after a meal, (a. 6 d.) *Chilliness after a meal.* Burning, like heartburn, from the stomach up to the pharynx, after eating flatulent aliments. Heartburn for three hours, after supper. Sour eructations after a meal. Frequent eructations, especially early in the morning. Unsuccessful desire for eructations, followed by spasmodic contraction of the stomach, morning and afternoon. Loud eructations, with accumulation of water in the mouth. Eructations tasting of the ingesta. Eructations as of bitter-sour water. Sour eructations, early in the morning, (10. d.) A quantity of sourish eructations, in the afternoon, with inclination to vomit. Acidity rises from the stomach to the mouth. Sour regurgitations. Gulping up of food and acidity, after great uneasiness, commencing from the pit of the stomach. Gulping up of water from the stomach, she spit up a good deal of it, after midnight. There is a constant inclination of something rising from the stomach into the mouth, (shortly.) Heartburn. Hiccough, at noon. Constant hiccough, before midnight. Qualmishness the whole day, with eructations in the morning. Qualmishness, as if he would faint. *Nausea, as if he would faint;* it only subsided when lying down; in the forenoon. Violent nausea in the stomach, with trembling of the hands and feet. Nausea, as from a spoiled or empty stomach; it does not go off by eating, with frequent waterbrash. Nausea with qualmishness, accumulation of saliva in the mouth, (and diarrhœa.) Waterbrash. Continued loathing, as if he would vomit, (shortly.) Nausea unto vomiting, in the stomach, at night, when in bed; gradually going off after getting up from bed. Inclines easily to vomit, especially after dinner. Inclination to vomit at every internal emotion,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

every vexation and joy, and at any time of the day; only retching in the morning. Retching, for several evenings. Frequent vomiting, without however the stomach being over-loaded or spoiled; next day faint and without an appetite, (a. 13 d.) Vomiting with a swoon-like failing of strength. Vomiting of food and sour substances, with nausea. In the morning the child is glowing red in the countenance, throws up the breakfast, turns as pale as a corpse, feels well again after several vomitings, but remains very faint for two days.

STOMACH.—Frequent pain in the stomach, but seldom in the afternoon, preceded by a sensation as of swashing in the abdomen, relieved by eructations and emission of flatulence. Pressure in the stomach, with rumbling, feeling of emptiness, and eructations. Frequent pressure in the stomach, early in the morning, when waking. Pressure in the stomach, as if from a stone, early in the morning, when in bed, relieved by clearing the throat. Pressure in the stomach extending into the chest, with want of breathing unto suffocation, accompanied by nausea and great depression of strength; she had to lie down, had a trembling of the hands and feet, and then threw up bitter water, which afforded her relief. Pressure across the stomach, and below the hypochondria, in the evening when in bed, for half an hour. Pressure in the pit of the stomach, (a. 21 d.) Pressure in the pit of the stomach and lower region of the chest, with aggravated breathing and rising of heat to the head, going off by eructations, after having lasted one hour. Pressure below the pit of the stomach, early in the morning and afternoon, relieved by bending the trunk backwards, and after a meal. Pressing pain in the pit of the stomach, obliging one to lie down. Weight in the stomach. Fulness and pressure in the region of the stomach. Fulness in the pit of the stomach. Sensation of swelling about the whole region of the stomach. Continual feeling in the stomach as if it were full of water. Spasmodic pains in the stomach, with pressure in the pit of the stomach. Violent spasmodic pain in the stomach, almost resembling a pressure and cutting, (shortly.) Violent spasmodic pains in the stomach, in paroxysms, relieved by walking, early in the morning.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Spasmodic contraction in the pit of the stomach, and transversely across the chest. Violent contractive pains in the stomach, even at one o'clock in the night, extending into the chest, and below the shoulders; where they become lancinating, with a choking sensation in the throat and oppressed breathing; afterwards anxiousness, short sweat, and eructations which afforded relief; returning in frequent paroxysms until early in the morning. Painful contraction from both sides of the stomach, with feeling of repletion, relieved by throwing up clear water. Pains in the stomach as if the parts were being screwed together, especially at night, extending as far as the chest and bowels, as if the stomach would burst, with impeded breathing and talking; in paroxysms. Constrictive pain in the stomach and towards the pharynx. The constrictive, spasmodic pains in the stomach are renewed by taking ever so little (especially cold) food or drink. The constrictive spasmodic pains in the stomach are frequently succeeded by eructations that afford relief, or by chills with shaking of the body, especially the hands, back and head, and by ordinary stool. Drawing and cutting transversely through the stomach, early in the morning after rising. Cutting pains in the stomach, towards evening. Sensation in the stomach as if it were cut to pieces, with great sensitiveness of the external region of the stomach, early in the morning. Painful cutting in the pit of the stomach, during and after breakfast. *Burrowing sensation in the stomach*, with painful contraction and sensation as if every thing in the stomach would turn up, with rising of water into the mouth, going off by the dinner, but returning afterwards, with burning rising in the throat. Burrowing and digging sensation in the stomach, as if it were being pierced. Burrowing sensation in the pit of the stomach, in the afternoon; afterwards frequent rising of bitter water, almost like waterbrash. Stitching pain in the stomach, with sensation as if every thing in the stomach would turn up, recurring after a meal. Stitches in the stomach, moving towards the left axilla, and afterwards towards the small of the back. Sore pain in the pit of the stomach, during both an inspiration and expiration. Sudden shock in the stomach, passing over

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

into rising of air or hiccough. Darting on the right side of the pit of the stomach, (in a few h.) Throbbing in the region of the stomach, it is painful to the touch. Throbbing in the pit of the stomach, like a violent palpitation of the heart, the pit being raised visibly at every throbbing; mostly in the morning, for a quarter of an hour. *Throbbing on the left side near the pit of the stomach.* Rising of heat from the abdomen to the stomach, in the forenoon. Burning in the stomach. Burning in the stomach, in the forenoon, after eructations. Sour burning from the stomach, with slight spasmodic constriction. Sensation in the stomach as of flatulence. Grunting, rumbling and moving about in the stomach, as of flatulence, or as if diarrhœa would set in. The region of the stomach is very painful when touched, talking, eating, etc. Itching, externally, in the pit of the stomach, it cannot be relieved by scratching.

ABDOMEN.—Simple pain in the hypochondria, with grunting in that part. Stitches in the hypochondria and the pit of the stomach, arresting the breathing. Frequently recurring burning stitches in both costal regions, in the afternoon, (12. d.) *Pain in the liver*, when walking, several days in succession. Pain in the liver as if it had been made sore by pressure. *Pressure in the direction of the liver*, as if coming from the right half of the chest, with throbbing in the region of the stomach, which is painful to the touch. The pressure in the liver is changed to a sensation of weight in that organ. Drawing pain in the liver. Cutting pain in the right infra-costal region, with pressure in the pit of the stomach. Lancinating tearing in the right infra-costal region. Stitches in the region of the liver, resembling splenetic stitches. Stitches under the last rib, during breathing, (1. d.) Stitches under the last right rib, independent of breathing, for 4 days. Stitches between the middle ribs of the right side, when sitting, (1. d.) Dull stitch in the right side, below the ribs, early in the morning. Dull stitches in the region of the liver and right groin. Sharp stitches in the region of the liver. Frequent dull stitches, in a small place of the region of the liver, with sore pain when touching it, (18. d.) Clawing stitches in the region of the liver.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Stinging throbbing on a rib of the right side, opposite the pit of the stomach. Feeling of heat in the region of the liver. *Burning pain in the region of the liver*, (the first days.) Tearing stitches in the region of the left ribs, arresting the breathing. Cutting pain in the left hypochondriac region, coming from the lower part of the chest, where one experiences stitches at the same time. Violent colic, sometimes extending into the hip, until late at night, (1. d.) Colic with a number of eructations. Colic, with a number of eructations, and spitting. Pressure in the abdomen. Itching in the epigastrium, extending into the pit of the stomach, in the evening, (35. d.) Pressure in the abdomen, (26. d.) Violent pressure in the left side of the abdomen, from stooping, (9. d.) Aching in a small spot of the left iliac region, in the evening, (39. d.) Aching in the hypogastrium, over the os pubis, in frequent paroxysms, going off by emission of flatulence, (1. d.) Pressing and burrowing in the abdomen, below the umbilicus, as if from incarcerated flatulence; he has to stoop when sitting, and feels worse when walking in the open air, (a. 19 d.) Distension of the adbomen with aching pain, feeling of repletion, weakness, and want of disposition to take the least exercise or to occupy the mind. Feeling of distension in the abdomen, below the umbilicus, going off by motion. Distended abdomen. Distended abdomen, after the cough, which Kali had brought on, had disappeared. *Distended abdomen. Hard, distended abdomen*, with painfulness of the umbilical region to the touch. Distension, pressing asunder and pinching in the abdomen, succeeded by soft stool. Bloated abdomen. Tense and heavy abdomen, when sitting or walking. Sensation as of a heavy burthen in the hypogastrium, more aching than pinching, and most insufferable when walking, (a. 3 h.) Pain in the abdomen, resembling a spasm, (a. 25 d.) Spasmodic contraction of the abdomen, producing a sort of coldness. Contractive pain in the abdomen. Painful retraction of the umbilical region when sitting, going off by motion. The abdomen presses outwards, several times, (a. 10 d.) Clawing pain in the epigastrium early in the morning, (11. d.) Clawing pain in the hypogastrium, (30. d.) Convulsions in the abdomen. Pinch-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

ing below the umbilicus during dinner, and after rising from one's seat; with burning in the right iliac region, with great internal and external sensitiveness of the same, and sensation, when stooping, as if something would fall out, gradually passing off when at rest. Pinching pressure in the left hypochondriac region, as if from incarcerated flatulence, (11. d.) Pinching in the epigastrium, towards noon, (25. d.) Pinching in the abdomen and distension of the same. Pinching in the abdomen, in the morning when in bed, after previous chilliness, with desire for soft stool. Cutting in the bowels, very painful; to find relief, he has to stoop while sitting, and press upon the parts with both hands, or else to recline a good deal backwards; he dares not sit straight. Cutting in the epigastrium as if from flatulence moving about, with emission of flatulence, when walking. Cutting in the left hypochondriac region. Frequent cutting in the abdomen, as if diarrhœa would set in. Frequent and slight cutting around the umbilicus. Cutting in the abdomen, as if all the contents would tear, first, deep in the hypogastrium, afterwards higher up, (1. d.) Cutting and drawing in the abdomen, resembling false labor-pains, (12. d.) Tearing in the right side of the abdomen or groin, sometimes darting, in the evening, (16. 17. d.) Intermittent tearing or dull stitches in the left iliac region, near the hip, (11. d.) *Stitches in the abdomen*, in the forenoon and again in the evening. Occasional stitches over the umbilicus, as if from flatulence. Stitches in the right side of the abdomen; as are felt after retaining the urine when asleep, relieved by emission of flatulence. Stitches in the right side of the abdomen, when laughing. Stitches in the right side of the abdomen, like fine dartings. Stitches in the left side of the abdomen, under the ribs. Sharp stitches and lancinating tearing in the left side of the abdomen, under the short ribs, (8. 9. 17. 24. d.) A few violent stitches in the hypogastrium, (a. 6 h.) Dull stitches on the right side near the umbilicus, (19. d.) Dull stitches and pressure in the left hypochondriac region, (10. 20. d.) Dull stitches in the region of the left kidney, first during an expiration; afterwards many in succession, going off by friction. Pinching-cutting stitches deep in

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

the left iliac region, as if caused by incarcerated flatulence, extending into the anus and perinæum ; they become more painful when drawing the abdomen in, and are but little relieved by the emission of flatulence, (19. d.) Sore pain in the abdomen, with bearing-down towards the genital organs, as if the menses would make their appearance, and pain in the small of the back. Long-continued, contusive pain in the region of both kidneys, in the afternoon, when sitting, (1. d.) Throbbing in the abdomen. Burning and drawing in the abdomen. Burning around the umbilicus, with pinching in the abdomen during dinner. Cold feeling in the abdomen, as if a cold fluid were being poured through the intestines, (during the menses.) Chilliness and sense as of swashing in the abdomen, as if it were full of water, generally only on the right side, in the evening. The abdominal muscles are painful to the touch. Itching of the abdomen, for several days, (a. 10 d.) Itching around the umbilicus. Pain in the right groin as if something were swollen. Pain in the right inguinal region, when drawing the abdomen in, (29. d.) Pressure in the groins, as if hernia would come on. Bearing-down in the groins, with sensitiveness to the touch, (going off after emission of flatulence.) Painful bloatedness in both groins, after dinner, when sitting. Pinching in both groins ; afterwards, prickings in the anus when sitting ; still worse after rising ; it lastly comes on when walking about, and is aggravated when sitting ; desire for stool. Drawing stitches and pressing outwards in the inguinal region, as if the old cicatrix of an operation for hernia would open again. Stitches in the groins, when moving about or extending the limbs. Sudden lancinating pain in the left inguinal region, during stool, with swelling of the glands. Intermittent, guggling pressing outwards in the right inguinal region, (27. d.) Incommoded by flatulence, (1. 2. 3. d.) Flatulent colic. Flatulent colic, going off after an eructation and emission of flatulence. *Incarceration of flatulence*, (even a. 20 d.) Incarceration of flatulence with colic. The emission of flatulence is troublesome and checked, with insufficient stool. The flatus become incarcerated and press painfully upon the bladder, (a. 2 d.) Moving of flatulence in the abdomen, with desire for stool, relieved by emission of

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

flatulence. Moving of flatulence in the abdomen, afterwards cutting in the stomach, with pressure extending up into the neck, both when at rest and in motion. Constant grunting in the abdomen, with frequent yawning and eructations. Sort of rumbling in the epigastrium, (previous to dinner,) as if diarrhœa would set in, and concealed colic, (1. d.) Guggling in the left iliac region, when pressing upon it. First, incarceration of flatulence, afterwards excessive emission of flatulence. Pressing flatulence; he is scarcely able to prevent its emission. Emission of flatulence upwards and downwards, with relief. Emission of fetid flatulence. Frequent emission of fetid flatulence, at night. *A quantity of flatus is emitted.*

Stool.—Unsuccessful desire for stool, with sensation as if the rectum were too feeble to expel the fæces. Frequent, violent desire for stool, in paroxysms, only a little good stool, or flatulence, being passed. Frequent desire for stool, at night, going off by emission of flatulence, (a. 3 d.) Constant desire for stool; something is being passed all the time. Frequent desire for stool; he passes a little at every stool. Frequent desire for stool; he feels as though he could not discharge the fæces all at a time, (a. 24 h.) Insufficient discharge of fæces; most of it remains behind. Insufficient stool, after much bearing down. *Insufficient, soft stool.* Tough stool, as though it would not pass. Tough, soft, dark stool. Costiveness, (a. 3 d.) Costiveness, with painful drawing in the abdomen. Hard stool, and uneasiness in the abdomen. Hard, retarded stool, sometimes accompanied with violent bearing down, or followed by tenesmus. Hard, insufficient stool, early in the morning; another stool, soft, in the morning, (2. d.) Hard stool every other day. Stool, resembling sheep's dung, coming off with pain and difficulty. Three scanty, but otherwise natural stools, (1. d.) Copious, brown stools. Stools more soft than hard, for several days, (a. 4 d.) Soft stools, preceded by colic, (1. d.) Soft stool, followed by burning at the anus, (in ½ h.) Thin stool, with pinching and uneasiness in the abdomen. Half liquid (scanty) stools, with colic and subsequent tenesmus. Half liquid stool, early in the morning, preceded by colic. Sudden and vehement desire for stool, as in diarrhœa, although the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

stool was hard, with colic. (Shortly after a new dose.) Diarrhœa, at night, with insufferable colic, continuing even the next day. Diarrhœa, in the evening. Profuse diarrhœa, day and night, (a. 22 d.) Profuse diarrhœa, with great weariness, (a. 27 d.) Diarrhœa with pinching, deep in the abdomen, before and after. Diarrhœa, preceded by pinching in the abdomen, and followed by burning at the rectum. Violent diarrhœa, with much colic, (a. 4 d.) Diarrhœa, with smarting pains at the anus, (a. 8 d.) Painless diarrhœa, with rumbling in the abdomen. Diarrhœa, the first fortnight, with great weakness, lying down, want of appetite, and colic every day; the fæces are light-coloured and gray. Two liquid stools, preceded by rumbling in the abdomen. Very fetid stool. Emission of flatulence, with discharge of thin stool, without being aware of it. Blood with stool, for several days, (a. 4 d.) Stools streaked with blood, followed by anxiety and difficulty of breathing. White mucus comes out of the rectum before and during stool. Discharge of a lumbricus with the stool. Pieces of tænia are expelled with the hard stool. Painful bearing-down towards the womb, during natural stool. Violent attack of spasm of the stomach, which obliged her to sit down, at the commencement of an alvine evacuation; she passed water; while sitting, the pain increased to such an extent that she had to bend double, and was unable to speak, accompanied by nausea, gulping up and vomiting of water, with retching; the vomiting was preceded by shuddering, and accompanied by vertigo, with shaking of the hands and feet, followed by anguish and heat in the whole body; alleviation of the pain, deadly pallor of the countenance, and lastly, regular stool, (a. $\frac{1}{4}$ h.) Pinching colic, with the regular stool, followed by a constant desire for stool, until another liquid stool comes on in the afternoon, (4. d.) The pains recur after the usual stool, early in the morning, (2. d.) Tenesmus of the rectum, after the usual stool, (1. and 4. d.) *Continual burning at the anus, after stool.* Shuddering around the anus, for half an hour after stool. Pressure in the abdomen after a difficult, scanty stool. Pain in the anus, after vomiting, as if it would burst, scarcely to be endured. Tenesmus of the rectum. Tenesmus of the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

rectum and anus. *Severe lancinating tearing and cutting in the anus*, (recurring for several days.) Stitches in the rectum. Prickings in the anus. Frequent stitches in the anus, between the stools. * *Itching of the anus. Violent itching of the anus* and scrotum. Itching of the anus, after supper. *Tingling of the anus*, (even a. 6 d.) Violent itching and tingling of the anus, in the evening, continuing a long while, (1. d.) Tingling and stitches of the anus, in the evening. Prickling in the anus, previous to every stool. *Burning at the anus*, without tenesmus. Burning at the anus, during and after the dry stool. Burning at the anus, which prevented him from sleeping, (a. 21 d.) Burning at the rectum, after stool. Burning and contraction of the anus. Burning and pinching of the anus. Frequent *burning and pinching of the rectum*, (the first days.) Burning cutting in the anus. Smarting as from excoriation in the anus, in the evening. Sharp smarting and sore feeling at and above the anus, after the morning-stool. Soreness at the anus, (5. d.) Pimples at the anus. Ulcerated pimples at the anus, with stinging. Protrusion and distention of the varices of the rectum, during hard stool. Protrusion of varices, with diarrhœa, with prickings and burning, for many hours. * *Large painful varices.* Considerable protrusion of the varices during micturition, emitting, at first, blood, afterwards a white mucus. *Greatly distended varices of the rectum*, and considerable emission of blood from the same during micturition. Considerable discharge of blood from the distended varices, with good stool. Considerable discharge of blood from the rectum; afterwards, agitation of the blood and pulsations in the whole body. Burning of the varices of the rectum, with violent pain when walking. Inflammation of the varices of the rectum, (a. 24 h.) Sore pain of the varices of the rectum. Stitches in the varices of the rectum. Tingling in the varices of the rectum, as if caused by ascarides.

URINARY ORGANS.—* *Great desire to urinate.* Desire to urinate; it took a long time, however, before he succeeded in expelling the urine; it flowed very slowly. He is frequently obliged to urinate; but frequently the pressure upon the bladder lasts a good while before the urine

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

comes off; even at night, he is frequently obliged to rise for the purpose of urinating, although he drinks but little. She is frequently obliged to urinate, emitting but little urine at a time, but always experiencing a renewed and almost painful desire after every emission, (a. 48 h.) She has to make an effort before the urine makes its appearance. A little quantity of pale urine, (1. and 2. d.; greater on the third day. Increased secretion of urine, (the first days.) When eagerly busy about something, she has to emit urine very hastily. After micturition, a few drops more flow out. Discharge of a few drops of urine, two, three minutes after micturition. Turbid urine. Greenish pale urine, with burning during and after micturition, (8. d.) Dark-yellow urine, cloudy; pale urine an hour afterwards, (a. 1 h.) *Fiery, diminished urine*, (the first time.) Urine, resembling loam-water, with copious sediment when standing. Interrupted stream, without any pain, in the afternoon, (1. d.) Discharge of a milky, inodorous, flocculent fluid after micturition, (prostatic juice?) Cutting in the region of the bladder. Cutting tearing in the neck of the bladder, during micturition, (37. d.) Intermitting cutting in the urethra, between the acts of micturition, passing over into a tearing pain in the glans, and especially in the orifice of the urethra, (7. d.) Frequent drawing and sharp tearing in the forepart of the urethra, (18. 19. d.) Tearing pain in the urethra. Pinching tearings in the urethra, (a. 12 d.) *Burning in the urethra, during and after micturition*, (a. 5 d.) Burning and biting sensation in the orifice of the urethra and the upper part of the internal surface of the prepuce, early in the morning when in bed, shortly after micturition, (20. 21. d.)

GENITAL ORGANS.—Violent itching of the mons veneris and near the genital organs, on the thighs; violent itching, with fine red pimples. Straining sensation in the penis, (a. 24 d.) Sharp drawing pain through the penis, (a. 12 d.) Tearing drawing in the penis, (24. d.) Gurgling sensation in the glans, (a. 2 d.) Tearing in the glans, (a. 20 d.) Stinging itching of the glans. Straining sensation in the left testicle. Pinching in the left testicle and the os pubis. * Swelling of the testes and the spermatic cord, with heat which may be felt externally.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

The scrotum is painful, as if it were bruised. *Itching of the scrotum.* Soreness of the scrotum, (a. 17 d.) *Excited sexual instinct,* (a. 24 h. and 3 d.) Burning sensation and excitement of the sexual organs. The male organs of generation strongly smell of semen. * Deficient sexual instinct, with undiminished morning erections. Nightly erections, without exaltation of fancy, (a. 7 d.) Frequent erections, (a. 13 d.) Vehement erections, (a. 24 h.) A number of erections, even painful ones, with spasmodic contraction in the spermatic cords. * No erections for 18 days. Pollution two nights in succession, (1. 2. night.) Pollutions, with voluptuous dreams, (the first days.) Copious *pollutions with subsequent weakness,* (a. 3, 4, 7, 23 nights.) Decreased quantity of pollutions, otherwise very frequent, (a. 14 d.) Suppression of the usual pollutions for 42 days. Embrace without emission of semen, (a. 10 d.) Lewd dreams after an embrace, at night, and a pollution. Aversion to an embrace, in a female, (the first days.) Her sexual desire is easily excited, (a. 29 d.) Pinching in the vagina, during an embrace. Sore pain in the vagina, during an embrace. Tearing in the left side of the pudendum, extending through the abdomen as far as the chest. Pinching pain in the labia. Stitches transversely through the pudendum. Burning stinging of the pudendum. *Burning and itching* in the pudendum. Menses too early, by two days, immediately after a new dose, (8. d.) Menses too early by 4 days, (a. 24 h.) Menses too early by 5 days, and more profuse, and lasting longer than usual. Menses too early by 6 days. Menses too early by 6 days, scanty on the first day, on the second day more profuse than usual; again scanty on the third, and ceasing entirely on the fourth. Menses too early by 10 days, lasting 6 days; weak during the first days, more profuse during the latter; with weakness and drowsiness, colic and toothache. Reappearance of the menses, which had been suppressed for 87 days, without any other trouble, except heaviness in all her limbs the day before, (3. d.) Reappearance of the suppressed menses: they look better,† (5. d.) Menses delay one day, with pain in the

† **Kali brings back the menses, when Natrum m. did not accomplish this result.**

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

abdomen. Menses delay 13 days, (secondary effect.) The menstrual blood is very acrid, has a bad, sharp smell; it makes the thighs sore, and covers them with an eruption. Great soreness around the pudendum before, during and after the menses. Voluptuous feelings previous to the appearance of the menses, early in the morning when walking, such as are experienced during an embrace. A good deal of heat previous to the menses, great thirst, and restless nights. Impatience, one week previous to the appearance of the menses, as if they would reappear, (a. 16 d.) A good deal of chilliness previous to the menses, tremor of the limbs, spasmodic sensation in the abdomen. Headache with great heaviness during the menses, early in the morning. Repletion and nausea after a meal, during the menses, soon after vomiting. At the time when the menses ought to have made their appearance, but did not, she was attacked with sour eructations, swelling of the cheeks with stitches, but without any heat and swelling of the gums. Pain in the abdomen, during the menses, putrid taste in the mouth, rumbling in the abdomen, great weakness and drowsiness. On the second day of the menses, violent headache from morning till evening. A quantity of flatulence during the menses, fetid taste in the mouth, and frequent rising of bile. Cutting in the abdomen during the menses. During the menses, violent pressing in the small of the back and in the abdomen, as if all the contents would issue out from the genital organs. Costiveness during the menses. Pain in the small of the back, like weight, during the menses. On the second day of the menses, coryza, colic, toothache, pain in the small of the back, stitches in the ears and restless sleep. Restless sleep during the menses, with anxious dreams. During the menses she never falls asleep again after waking in the morning, but passes into a very disagreeable state between sleep and waking; while in that state she hears things which give her anguish and pain, although she knows that she dreams; she is, nevertheless, unable to open her eyes, and she has great difficulty to tear herself from that state of half waking. Violent itching of the whole body, during the menses. After the menses, in the evening, coldness in the back, and waking

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

after midnight, with spasm of the stomach and coldness in the stomach, which continued until noon, (a. 19 d.) A female in her fifth month of pregnancy is attacked at night after some vexation, with a violent hemorrhage from the vagina, with clots of coagulated blood, accompanied by dull headache and yellow complexion; however she did not miscarry. * Leucorrhœa, (3. d.) Leucorrhœa, (the first 5 days.) Mucous leucorrhœa. Yellowish leucorrhœa, with itching and burning of the pudendum.

Cold, Catarrh.—Dry coryza, (a. 26 d.) Dry coryza, becoming fluent in the afternoon during a walk, (a. 3 d.) Dry coryza with itching in the nose; she has difficulty in breathing through the nose, for several days, (a. 4 d.) Dry coryza, frequently, also in the evening, when in bed, with tingling in the throat, (a. 11 d.) Dry coryza, he found it very difficult to breathe. Dry coryza, with a quantity of yellow-green nasal mucus. Obstruction of the nose. Expulsion of purulent matter from the right nostril; followed by obstruction; when blowing the nose, one experiences a stinging and contractive pain extending as far as the occiput. Fluent coryza, with excessive sneezing, at least 30 times in one day. Fluent coryza, almost the whole day, but especially in the evening. Profuse fluent coryza, every evening, with frequent sneezing. *Profuse fluent coryza.* Profuse fluent coryza, with much sneezing, pain in the back and headache, (a 10 d.) Excessive fluent coryza, (a. 29 d.) Coryza, with bloody nasal mucus, (a. 8 d.) Frequent drawing pain about the larynx, with a raw feeling. When eating, particles of food easily get into the larynx. Rough voice.

Throat.—Rough throat, with a good deal of sneezing. *Rough throat with cough.* Rough throat when uncovering the body. Rough and hoarse throat, for several days. * Complete hoarseness and aphony, (a. 24 h.) First, hoarseness, afterwards excessive fluent coryza. Hoarseness, as if something were lodged in the throat, with irritation, inducing clearing. The throat feels plugged; the throat is cleared by coughing. Scraping sensation in the chest, caused by the wind. Grunting and slight snoring in the trachea, previous to the coughing fit. Titillation in the larynx, inducing cough, with violent

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

hoarseness. *Titillation in the throat, inducing hawking and coughing*, with sensation of tenacious mucus, morning and evening, (12. 22. 29. d.) * Cough, brought on by titillation in the throat, (20. d.) Cough, brought on by titillation in the throat, without expectoration. Scraping sensation with cough. Cough, affecting the chest, brought on by tickling in the throat. *Cough, with titillation.* Cough, when playing upon the violin. Violent cough, early in the morning, before breakfast, subsiding after breakfast. Early in the morning, hawking up of mucus. She begins to cough at three o'clock in the morning, the fits recurring every half hour. A good deal of cough, early in the morning, with expectoration, but mostly in the evening. Evening-cough, when in bed. Profuse cough every evening, after having laid in bed for some time, for several weeks. Fatiguing cough in the evening. She was obliged to cough from 9 o'clock in the evening, until morning, every 5 minutes. * *Night-cough.* Frequent cough before midnight, not in day-time. Short and hacking cough with some expectoration, mostly at night and early in the morning, with coryza. Frequent short and hacking cough, in the afternoon and the following forenoon, (a. 6 d.) Spasmodic cough, with irritation in the throat, in a few violent paroxysms, even unto choking, with a sore pain in the top of the head and subsequent great weakness. Suffocating and choking cough, at 5 o'clock in the morning, as if caused by dryness of the larynx; she was prevented from talking by a spasm in the chest, accompanied by redness in the face and sweat over the whole body. Cough, bringing on vomiting easily. *Violent cough, almost causing vomiting*, early in the morning. Fatiguing cough, so violent that she loses her senses. Dry cough, coming and going rapidly. Dry cough, almost only at night, with stitches in the throat. Dry cough, at night, waking her from sleep with intense pain in the chest, when coughing; little cough in day time, (1. d.) Cough with abundant expectoration. Discharge of small, red lumps from the throat. Sourish expectoration from coughing. Cough, with expulsion of blood-streaked mucus, three times, (17. d.) She coughs up mucus, which, however, does not get into her throat, and cannot, on that

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

account, be thrown off. Rough pain in the larynx, when coughing. Stinging in the throat, when coughing, with fluent coryza. Occasional stitches in the left half of the chest when coughing. Tearing scraping in the chest, when coughing. Sparks dart from her eyes when coughing. Nausea when coughing. Pain in the varices when coughing. Concussive pain in the abdomen when coughing.

CHEST.—Short breath, early in the morning. *Shortness of breath, early in the morning.* Inclination to breathe deeply. * Oppressed breathing. Asthma, as if caused by impeded expansion of the chest. Asthma with short breathing, while writing, (3. d.) *Oppression of the chest,* with groaning, deep breathing. Oppression of the chest, with aggravated, painful breathing, 2, 3 times, (a. 30 d.) Oppression of the chest, with distended abdomen. Sensation in the open air as if the throat were being constricted. Arrest (interception) of breathing wakes him at night. Slight rattling in the chest, at night, when lying on the back. Anguish in the chest, towards evening. *Great pain in the chest,* especially *when talking.* Pressure in the chest, when taking an inspiration. Paroxysms of pressure, in the forepart of the chest, especially on the right side, increased by inspiration, diminished by eructations. Pressure in the xiphoid cartilage, when coughing, and also when taking a strong inspiration, (a. 16 h.) Frequent pressure in the left half of the chest and the region of the heart, (8. d.) Pressure in the whole of the left half of the chest. Aching in the right border of the sternum, early in the morning, when rising; the border is painful when touched. Pressure in the middle of the chest, with a sensation as of drawing downwards. Sharp aching behind the sternum, when breathing, or when swallowing liquid substances, and during eructations, for several days. Stinging pressure in the left half of the chest, when breathing deeply. Stinging pressure in the right half of the chest, sometimes, for several days. Clawing pressure in the right half of the chest, (26. d.) Tension across the chest, during an expiration, when walking. Spasmodic pain in the chest, going off by eructations. Pinching in the muscles of the chest, several

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

times. Pinching, dull stinging in the right half of the chest. Cutting sensation in the lower part of the chest, especially in the left side of it, moving into the epigastrium and leaving a stinging sensation in the left half of the chest, (20. d.) *Cutting pain in the chest*, early in the morning, especially around the pit of the stomach, as if owing to the pressure of incarcerated flatulence in that part. Cutting pain in the chest, in the evening after lying down; she knew not how to turn herself, and felt worst when lying on the right side. Stitches in the sternum, also during an inspiration, in the evening, (1. d.) Stitches under the left mamma, and sometimes ascending deep into the chest; also, in the evening. Stitches in the right half of the chest, during breathing. A stitch in the right mamma, occasionally. Stitch in the left mamma. Stitches in the left mamma. Stitches in the sides, when taking an inspiration. Stitches in the region of the heart. Violent stitch under both breasts, after lifting a heavy burthen, afterwards colic in both hypochondriac regions, in front, in the afternoon. Single cutting stitches under the right clavicle, with pain as if a thorn were lodged in the part. Suddenly a dull stitch in the sternum, during an eructation and when swallowing liquid substances. Dull stitches, deep in the left half of the chest, under the short ribs. Dull, painful stitches into the chest, under the left clavicle, going off for a short while by pressing upon the parts, in the evening. Dull stitches, pressure and tearing below the axilla. Burning stitches in the right half of the chest, when raising the head after stooping, (a. 1 h.) Burning stitches in the left side of the chest, when sitting, going off after rising (7. d.) Boring, deep into the left half of the chest, (9. d.) Drawing pain across the chest, (a. 4 d.) Tearing in the sternum, over the pit of the stomach, towards the left side, in the evening. Tearing in the left side of the chest, in the lower short ribs. Tearing pain in the right half of the chest, (9. d.) Tearing in the left half of the chest, (17. d.) Sore and tearing pain, a little below the left axilla, increased and excited by violent breathing. Sore pain, in the upper part of the chest, when taking an inspiration, when feeling the parts, and when lifting up something heavy. Pain in the chest, as if from

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

bruises. Tickling in the right half of the chest, (a. 16 d.) Guggling, resembling a jactitation of the muscles, in the upper part of the right half of the chest, (22. d.) Tremor or twitchings in the fore part of the chest. The chest is fatigued from loud talking. Weakness of the chest. Weakness and faintness in the chest, from walking fast. Painful throbbing in the clavicle, shoulders and side of the abdomen, etc. * *Frequent and violent* palpitation of the heart, with anguish. Palpitation of the heart, when he is hungry, (a. 10 days.) Violent palpitation of the heart, in the forenoon, with obtusion of the head, and nausea, (a. 24 h.) Frequent intermission of the beats of the heart. Clawing pain in or about the heart, as if the heart were hanging by bands firmly drawn together; the pain is most felt when taking a strong inspiration, or when coughing, not during exercise, (a. some hours.) Burning in the region of the heart, (a. 2 d.) Darting pain, externally, in the region of the lowest left rib. Tearing lancinations in the mammæ. Itching of the right half of the fleshy part of the chest, with fine eruption, only coming out when rubbing the parts. Biting sensation on the chest, in alternate places, in paroxysms.

Back.—Violent gnawing in the os coccygis, both when at rest and in motion. * Pain in the small of the back, after having stood or walked for some time. Pain in the small of the back, only when bending the part backwards, not when at rest. Frequent pain close over the small of the back, when sitting. Violent pains in the small of the back, with labor-like colic and leucorrhœa. Heavy pain in the small of the back. Flatulent pain in the small of the back, early in the morning, when in bed, with sensation as if large bubbles were incarcerated in the small of the back and were pressing against it, and with desire for stool, all these symptoms disappearing after emission of flatulence. Sensation, early in the morning, as if the small of the back were pressed into from both sides. Stiffness of the small of the back. Darting pain in the small of the back, when stooping, so that he was for a long time unable to raise himself again. * *Drawing pain in the small of the back.* Violent, continual drawing in the small of the back, alternating in the part with pulsations, relieved only when lying down. Throbbing in the small of the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

back. Violent pain as from blows and bruises, in the small of the back, especially early in the morning, when rising. Violent pain in the small of the back, as if the parts were broken, during motion. Itching of the lower part of the small of the back. Titillating weary pain over the os sacrum. An occasional stitch, from the small of the back, through the left side of the abdomen, towards the chest. Violent pain in the back. Pressure in the back over the region of the right kidney, early in the morning, (20. d.) *Pressure in the region of both kidneys*, (7. 8. 15. 19. d.) Pressure in the left scapula. Sharp pressure in the upper part of the back, (34. d.) Drawing pressure in the scapulæ. Drawing pressure in the back. Tensive pressure, like great fatigue, beginning at the right scapula, extending over the side of the back down to the small of the back, when the parts are at rest, even in the morning when in bed, especially when riding in a carriage. Burning pressure in the back, worse when walking in the open air, (a. 19 d.) Sore pressure in the region of the right kidney, (6. d.) * Stiffness of the back; she is unable to stoop. * Stiffness and paralytic feeling in the back and small of the back. Tensive pain under the left scapula, when breathing. A few sharp, biting pinchings on the posterior portions of the ribs, on both sides of the back. Contractive pain in the back, at rest, after performing some physical labor. Tearing in the region of the right kidney, (13. d.) Tearing in the psoas muscles, arresting the breathing. Lancinating and oppressive tearing in the back, near the right scapula, (10. 38. d.) *Tearing* in the right *scapula*, early in the morning, (4. d.) Burning tearing, on the right side, near the dorsal spine, over the small of the back, (18. d.) *Stitches in the region of both kidneys*, (11. 29. d.) Stitches in the right side, in the back, extending through the chest, (25. d.) Stitches in the right scapula, when taking an inspiration. Lancinating pain between the scapulæ, with oppression and anguish of the chest, almost only when sitting, so that she is obliged to get up and walk about. Stitch from the apex of the left scapula to the pit of the stomach, during fatiguing labor, (a. 7 d.) Dull stitches in the left scapula. Clawing stitches in both scapulæ. Sharp, tearing stitch under the right scapula. Pressure between the scapulæ, followed

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

by a burning sensation extending from the scapulæ to the crest of the ilium, equally felt when at rest and in motion, the burning is felt when laying the hand upon the parts. Pulsative throbbing near the upper border of the left scapula. Pain in the loins, when walking, and especially when touching the parts, as if the flesh had started off from the bone. *Pain in the back as if from blows and bruises,* when at rest, not in motion. Pain as from blows and bruises, between the shoulders and near the left shoulder, going off by motion. Stinging pain as from blows and bruises, in the right scapula, when in motion; it may be felt as far as the chest. Pain in the back as if the part had been strained by lifting. Pain as from a sprain in the left scapula. Violent stinging pain as from a sprain in the left scapula, extending into the chest. Itching of the back, becoming painful after scratching.

NECK.—Pain in the nape of the neck, when bending the head backwards. Violent tension in the nape of the neck, becoming still more painful when moving the head. *Stiffness in the nape of the neck,* with elongation of the uvula in the pharynx. * *Stiffness of the nape of the neck early in the morning, when in bed,* (also a. 3 d.) Stiffness in the nape of the neck, early in the morning, disappearing in day-time; for some weeks. Drawing pain in the nape of the neck, (a. 2 h.) Tearing in the nape of the neck, sometimes very quick and slight. Tearing in the nape of the neck, on the right side, early in the morning. Itching pimples on the nape of the neck, in the evening; they pass off after 24 hours. The cervical muscles are painful when acting. Darting pain in the left side of the neck, (a. 2 d.) Pressing drawing in the right side of the neck. Tearing in the lower part of the right side of the neck, (a. 24 d.) Congestion of blood to the neck; the neck appears larger, and the neck-cloth too tight. Pain of the cervical glands, as if from a cold, (3. d.) Stitches in the cervical glands. *Swelling of the cervical* glands, (a. 5, and 14 d.) Swelling of a cervical gland below the chin, painful when catching cold. Swelling of the cervical glands on both sides, with pain when turning the head. Hard swelling of the sub-maxillary gland. Tickling in the glandular swelling of the neck; she had to press upon the part with her cold hand, in order to find relief.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

SUPERIOR EXTREMITIES.—*Sweat in the axilla.* Swelling, and pain as from sub-cutaneous ulceration, of the axillary gland, (2. d.) *Swelling of an axillary gland.* Dull cutting and tearing in the right axilla, (24. d.) Tearing in the left axilla, when at rest, (10. d.) Tearing lancinations in the right axilla, when lifting up the arm, early in the morning, when writing, (7. d.) Violent stitches in the left axilla. Sensitiveness and burning in the right axilla, (15. d.) Itching in the axilla. The left shoulder is painful, when moving the left arm violently or when pressing with it strongly. Clawing pressure in the right shoulder-joint, more painful when breathing. Violent tensive pain in the left shoulder, early in the morning, so that he was unable to lift up his arm. Tension and pressive drawing in the right shoulder, with paralytic sensation in the right arm. Drawing pain in the right shoulder, (17. d.) Tearing in the left shoulder, when at rest and in motion. Tearing in the right shoulder, when knitting; going off after motion. *Tearing in the left shoulder.* Clawing tearing in the right shoulder. Fine stitches in the left shoulder, afterwards in the tendons of the left side of the neck. Fine stitches in the shoulders, when at rest and in motion. Pain as from blows and bruises under the right shoulder-joint, especially when moving and touching it. Cracking in the shoulder-joint, when moving or raising the arm. Pimples on the shoulder, with violent itching and burning after scratching. Violent tensive pain in the arm, for 8 days, so that he was unable to raise it on account of pain; he was indeed able to move it backwards upon the back, rest upon it and touch the joint, without any pain. Paralytic tension and drawing in the left arm, from the shoulder to the lower arm, with disposition of the lower arm to go to sleep, early in the morning when waking up, (34. d.) Drawing pain in the left arm, (a. 21 h.) Jerkings in the arm, in the evening when going to sleep. Frequent jerkings of the left arm. Tearing in the left arm from the top to the wrist-joint. Violent tearing in the whole of the left arm, (7. d.) In the cold air the arms lose their warmth, become numb and feel as if they had gone to sleep. * Both arms go to sleep—and become rigid in the cold air; they go to sleep even after violent exercise.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

The arm upon which he rests at night, goes to sleep. The arms go to sleep, early in the morning when in bed; they feel rigid and paralysed, with sensation of internal pressing, and the hands are insensible for half an hour. Paralytic sensation in both arms, (the first days.) When writing, the arms become easily tired, (a. 3 d.) *Weakness and want of strength of both arms.* Weakness of the arm, with swelling of the upper arms and hands. Itching of the arms, with white pimples after scratching, looking like millet-seed. Slight twitchings in the flesh of the left upper arm. Burning tension of the right upper arm, close above the elbow. Tearing in the right upper arm, above the bend of the elbow, in the evening. Tearing in the left upper arm, sometimes as far as the shoulder, (11. 16. 20. d.) Tearing in the upper part of the right upper arm, and in the elbow, (12. 22. d.) Lancinating tearing in the left upper arm. Stitches in the right upper arm. *Pulsative pain in the left upper arm, at intervals.* Jactitation of the muscles of the left upper arm, (18. 19. 25. d.) Pimple on the upper part of the upper arm, itching and painful. Pain as from bruises and blows in the right upper arm, especially when raising the arm. Paralytic pain in both upper arms, mostly during motion. Pain in the elbow as if it were stiff, when extending the right arm, after it had been bent. *Drawing and tearing in both elbows*, sometimes with a feeling of warmth in the parts, recurring frequently. *Tearing in the bends of both elbows*, (3. 6. 22. d.) Tearing stitches in the bend of the left elbow. Violent stitches in the bends of both elbows, early in the morning, when in bed, going off after rising. Tensive pain in the lower arm. Drawing pain in the lower arm. Violent, short drawing from the arm into the hand, (a. 2 d.) Paralytic drawing with dull pain, from the left lower arm into the hand, in the evening. *Tearing in the upper part of both lower arms.* Tearing in the middle of both lower arms. Tearing in both lower arms, in the direction of the wrist-joint. Pain of the metacarpus, when grasping something. Dull aching in the dorsum of the left hand. Drawing in the inner surface of the left wrist-joint, aggravated by motion. Tearing in the wrist-joints, (11. 20. d.) Tearing in the outer process of the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

wrist-joint, (29. d.) Fine tearing in the left wrist-joint, in the direction of the ring-finger, recurring frequently, (4. d.) Tearing in the right wrist-joint, when knitting, recurring frequently. *Dull tearing, with pressure, in both hands, between the thumb and index-finger*, (16. 12. 21. d.) Violent tearing extending from the dorsum of the left hand to the fingers. Violent tearing in the dorsum of the left hand, apparently in the marrow of the bones, scarcely endurable. Stitches in the left wrist-joint, when moving it, afterwards a few sharp stitches when at rest. *Cold hands.* Burning of the dorsum of the left hand, as from red-hot coal. *The hands go to sleep*, early in the morning when waking, with dull headache, increasing after rising, with frequent empty eructations, until noon. Want of strength of both hands. * Trembling of the hands when writing, early in the morning. Itching of the wrist-joint. Violent itching of the palms of the hands, in the evening, close to the fingers, (1. d.) Itching above the right wrist-joint, going off by scratching. Itching vesicles in the palm of the hand. Red, elevated spot above the wrist-joint, of the size of a small pea. Rough, chapped skin of the hands. Drawing pain in the posterior joints of the fingers. Tearing in the middle-joint of the index-finger. Tearing under the nail of the thumb. Tearing in the articulations and the phalanges of several fingers. Tearing under the finger-nails, (34. d.) Quick and slight tearing in the fingers and the toes at the same time. Tearing in the left thumb. Tearing in the metacarpal-joint of the left middle finger. Tearing in the left little finger, towards the tip, going off by moving it, but recurring frequently. Violent tearing behind the right index-finger, as far as the tip. Dull tearing in the ball of the thumb, (9. d.) A few fine stitches in the left little finger. Pricking tearing under the nail and in the tip of the index-finger. Drawing tearing in the tip of the little finger. Burning tearing in the tip of the index-finger, (10. d.) Stitches in the fingers of the right hand. Fine painful stitches under the nail of the left middle-finger. Tearing in the left little finger, towards the tip, going off by motion, but recurring frequently. Violent tearing behind the right index-finger, as far as the tip. Dull tearing in the ball of the thumb,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

(9. d.) A few fine tearings in the left little finger. Pricking tearing under the nail and in the tip of the index-finger. Drawing tearing in the tip of the little finger. Burning tearing in the tip of the index-finger, (10. d.) Stitches in the fingers of the right hand. Fine, painful prickings under the nail of the left middle-finger. Fine, intermitting prickings in the middle-joint of the right index-finger. Painful prickings in the right index-finger, as if needle and thread were being drawn from the posterior part to the tip; relieved by bending the finger, renewed by extending it again. Fine, painful stitches in the tips of the four last fingers of the right hand, as if they were ulcerated. Ulcerative pain in the posterior joint of the left thumb, in the evening. Burning pain as from a red-hot coal, of two fingers of the left hand. Burning in the tip of the little finger. Sore pain, early in the morning, in the anterior joint of the middle-finger, especially under the nail, not increased by contact. Tingling in the tip of the right middle-finger. The fingers are drawn open and then contract again, with a jerk, early in the morning, when in bed; afterwards numbness, difficult movement and chilliness of the fingers. Numbness and insensibility of the right thumb, for some weeks. The thumb goes to sleep, towards morning, when in bed. *The tips of the fingers go to sleep*, generally early in the morning. Paralytic sensation of the thumbs, from knitting. Stiffness and paralytic weakness of the right thumb and index-finger, when writing. Itching pimple behind the left thumb, continuing to itch even after being scratched. Vesicle on the little finger. Corrosive vesicle on the left index-finger, which, when opened for several days in succession, poured forth a watery, not purulent humor. Burning pain around the border of the nail of the middle-finger. Inflammation round the border of the nail of the index-finger, resembling a panaris; when pressing upon it a thin pus oozes out, for seven days.

INFERIOR EXTREMITIES.—Tearing to and fro in the left lumbar region, when sitting, going off by motion. Twitchings of the glutei muscles. Dull pain in the glutei muscles. Tearing about and in the glutei muscles, not far from the hip-joint. Attacks of clawing tearing in the glutei muscles.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Ulcerative pain in the glutei muscles and thighs, when sitting. Itching between the nates. Soreness between the lower limbs. Clawing tearing in the hip-joint. Tearing pain in the left hip, from time to time. *Tearing in the hips and knees*, even when sitting. Tearing in the hips or the integuments of the hips; occasionally with a sensation as of tingling, (19. 21. 30. d.) Darting pain in the left hip-joint, when turning the thigh. Fine, but intensely painful stitches in the left hip-joint, when standing; after sitting down, one experiences tearing stitches in the thigh from above downwards, apparently in the marrow, going off after rising from one's seat. Contusive pain, as from a shock, in the upper part of the left os innominatum, when walking, or when feeling the bone. Pain in the hip-joint as from bruises; with pain during motion and when sneezing. * Pressure in the lower extremities, apparently in the bones, at times in one, at times in another place. * Tearing in the left thigh and tibia (during the menses.) *Burning pain in both lower limbs*, sometimes with intensely painful stitches. Uneasiness in the lower extremities, in the evening; he had to extend them frequently. Sudden attack of heaviness in the left lower limb, so that he had great trouble in moving it; it felt paralysed, at night when in bed and next day when sitting. Heaviness of the lower limbs. Numb feeling in the whole of the right leg, as if it would go to sleep, with subsequent feeling of great heaviness, early in the morning after waking up. *Numb feeling of the whole of the right lower limb, with disposition to go to sleep*, especially in the leg. *The lower limbs frequently go to sleep.* Weakness of the lower limbs; they bend involuntarily. Stiffness of the lower limb, (a. 10 d.) *Jactitation of the muscles of the thigh.* Drawing pain in the left thigh, (a. 11 d.) Drawing pain in left thigh down to the knee. Drawing pain in the thigh, when stepping up as if it would break. *Paralytic drawing* in the whole *thigh*, which frequently passes over into tearing, worse when standing and in the warmth of the bed, frequently only in the evening, and at night. Tearing in the upper and internal part of the thigh. Tearing in the posterior part of the thigh, close to the glutei muscles, in the direction of

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

the genital organs. Sore pain of a spot in the middle of the thigh, when touched. Weary sensation in the thighs, as if he had fatigued himself by walking, for many days, mostly in the afternoon. Feeling of both limbs, over the knees, as if they had been knocked away, in the evening. Violent pain as from bruises, one hand's breath over the right knee, as if the thigh would fall off when standing; worse when sitting, and afterwards continuing both when at rest and in motion. Twitchings in the right thigh, on the anterior surface. Paralytic feeling of the whole thigh, with sensation as if it would go to sleep. Two litle pimples on the left thigh. Itching spot on the thigh, near the hip, ulcerating when scratched. Nodosity over the knee, aching and tearing. Dull pain in the side of the knee, when walking, and especially when extending the limb. Tensive pain in the right knee, and afterwards tingling in the part; it is felt only when walking, and goes off when sitting. Stiffness in the knees, (a. 2 d.) Stiffness, tension and weakness in the right knee, as if something were tied round the knee very firmly. Drawing pain in the knee, when walking, reaching into the thigh. *Frequent tearing in the knees.* Tearing in both knees, (1. d.) Tearing in the bend of the knee. Tearing in the knee and bend of the knee, in the evening, with warmth in the part. Ripping-up pain in the knees, when walking or sitting. Throbbing and beating in the left knee, frequently during the day. Slight twitchings in the left knee, (2. d.) Pain, as from a sprain, in the knee, when rising from one's seat, for some minutes. When walking considerably the knees go to sleep and feel a pain; he found it difficult to bend them. Paralytic sensation in the knee, when sitting. Paralytic pain in the right knee, when walking, (6. d.) Itching of the knee. Eruption in the bend of the knee. Drawing pain in the right leg, extending down to the foot, for several days. Paralytic drawing in the legs. *Drawing and tearing in the tibiæ.* Tearing in the upper part of the tibia, below the knee. *Tearing in both tibiæ, with pain of the periosteum when touching it, and tension of the same when walking.* Tearing in the left tibia. Tearing in the upper part of the calf, (20. d.). Tearing in the right

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

calf, receding towards the interior of the calf when pressing and rubbing it, and disappearing afterwards. Straining sensation in the left calf, as if the tendons were too short, when rising from a seat, in day-time. Cramp in the right calf, (a. 20 h.) Stitches in the tibia. Twitchings in the anterior surface of the left leg, when standing. Itching of the tibia. Violent itching of the legs, in the evening. Itching nodosity and three vesicles with inflamed areolæ, upon the tibia. Herpes upon the lower limb. Tension in the feet, almost without any swelling. Aching in the heel. Aching in the sole of the foot, when setting the foot down and walking. Clawing drawing in the foot, as if caused by great weariness. Spasmodic tearing in the region of the ankle, with pulsations all around, along the tibia as high up as the knee. Tearing around the ankles, when the feet are cold; it ceases when they become warm. *Frequent tearing in the tarsal joints.* Tearing close above the ankle, (20. d.) *Tearing in the inner side of the foot and in the sole.* Tearing in the dorsum of the foot, extending to the toes. Tearing from the internal malleolus transversely to the tendo Achillis. Drawing tearing in the foot, to the toes. Stitches in the left foot, towards the inner side. Stitches in the dorsum of the foot. Stitches under the ankle. When walking, she experienced a darting into the internal malleolus, as if the foot would break; she had to stand still. Violent drawing and stinging, early in the morning, a few hours after rising, in the left tarsal joint when setting the foot down, most violent and painful in the evening, with jerking pullings in the joint, and stitches in the heel; she dares not touch the foot, has to keep it balancing in the air all the time; the foot feels too heavy. the place is swollen and hot to the touch. Prickings under the heel. *Stitches in the feet.* Violent stitches in the tendons behind the outer malleolus of the right foot, when running; going off when at rest. Stitches and burning in the feet, after a walk. Visible twitchings in the dorsum of the left foot, in the evening. Tingling in the soles of the feet, towards the toes. Tingling burning in the soles of the feet, and painful sensitiveness of the same. Heaviness and stiffness of the feet. *The feet go to sleep after a meal.* The left

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

foot goes to sleep, while at dinner. * *Cold feet,* in bed. Cold feet, with heat in the face. * Sweaty soles. Profuse sweat on the feet, for many days. Swelling of the feet. * Considerable swelling of the feet, as high up as the ankles. Swelling and redness of the soles, with burning in the soles even while lying down, but much more when setting the foot down. Violent itching around the tarsal joint, early in the morning, while in bed. Violent itching and heat of the feet, in the evening, as if they had been frozen. The tips of the toes are very painful when walking. Spasm in the big toe of the left foot; he was unable to extend it, when sitting, in the evening. *Tearing in the toes,* (4. 7. 11. 20. d.) *Tearing in the posterior joints of the big toes,* (11. 16. 19. 34. d.) *Tearing in the tips of the big toes,* (17. 11. 36. d.) Gnawing sensation in the outer border of the big toe of the right foot. * Prickings in the ball of the big toe. Titillating stinging in the tips of the toes. Fine stitches, with itching, in the tip of the big toe. Pain, as from a sprain, in the posterior joint of the big toe, while walking, but most intense when raising the foot upon the toes, (11. 33. d.) Titillating tingling in the toe and sole. Itching in the lower surface of the toes. Violent itching of the big toe, below the nail, with pain when touching the part. Inflamed, red chilblains on the toes, aching. Chilblain on the ball of the big toe, with lancinating pain, red and large. Inflamed, blue-reddish chilblains on the ball of the big toe, with cutting and pricking pain, especially when having on shoes or boots. The corns are painful and sensitive. Stitches in the corn. One side of the nail of the big toe is painful, as if the nail would grow into the flesh, (a. 14 d.)

GENERAL SYMPTOMS.—*The limbs are painful in those parts upon which they rest,* (a. 4 d.) Her whole body felt very sensitive; she felt a pain wherever she touched herself, or in whatever part was set in motion, (3. 4. d.) *Aching in the knee-joints,* and the joints of the feet and hands, only when at rest. Aching in the joints and drawing pain in the tibiæ. A sort of tension in the interior of the body, extending as far as the head and eyes. Pinching and feeling of contraction ir the anus, in the region

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

of the stomach and towards the pharynx. Tingling in the limbs, especially the lower limbs, when sitting, with drawing in the legs, which makes him restless. Drawing pain in the whole body, in various parts, nape of the neck, shoulder-blades, hands, and knees, (a. 10 d.) * Drawing pain in all the limbs, with sensation as if he had been sick for a long while, with great paleness of face and emaciation. Violent drawing in the abdomen, arms and lower limbs, with pain as from bruises in the upper arms; worst when at rest, (the first days.) *Stitches in the joints and tendons. Stitches are the characteristic symptoms of Kali.* Ulcerative pain when pressing upon any part of the body. Pain as from bruises of all the muscles of the body. The limb, upon which she is resting, goes to sleep. The pains appear at two or three o'clock in the morning, and are so violent that she is unable to lie still on account of the pain; they are then more violent than they are in day-time during motion. Chilliness immediately after the cessation of the pains. She thinks she feels more easy in the open air than in the room. The symptoms become more intense in the open air, especially the fever. Drowsiness with yawning, when walking in the open air. Violent tearing in the outer side of one part of the head, when walking in the open air. Violent headache, for some hours, from walking in the open air. Great dread of the open air. Liability to cold when walking in the open air, with sweat and restlessness at night, and weight in the nape of the neck as from a load, (4. d.) *Liability to catching cold.* Catches cold easily after having heated himself by exercise; he loses his appetite, is attacked with feverish chills, diarrhœa with colic, restless sleep, etc. After cold, headache on the right side, and heat in the eyes. Every breath of air causes catarrhal ailments. After having caught cold by a draft of air, she feels hot for a moment in the room, is then attacked with heaviness in the limbs, tearings in the whole body and about the head, with whizzing before the ears, and cold all over, with sour-smelling sweat in the night, (a. 31 d.) After catching cold, he is attacked with fever in the evening, sweat towards morning, with violent headache and a dreary feeling in the head after rising. Itching in different parts

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

of the body, especially the legs; the parts become easily ecchymosed when scratching them. Violent itching on the abdomen and thighs. Itching of the whole body, at times in one, at times in another part; scratching brings on little blotches. Itching of the whole body, in the evening before going to bed, disappearing in bed. Violent itching of the whole body, morning and evening, especially on the back, where small pimples make their appearance, (a. 3 d.) Violent, almost prickling itching over the whole body, at night when in bed. Itching stitches, in various parts of the body. Stitches and gnawing in the skin of the whole body. Burning itching in the face, back, and on the head. Burning itching of the os sacrum and below the right patella. Burning itching of the whole body, in the face, on the hands, tibia, etc., at times in one, at times in another place. *Burning in many places of the skin*, even below the axillæ, *as if from a drawing plaster*. Pimples, in different parts of the body, also in the face; nettle-rash, for a fortnight. Nettle-rash, with itching, so violent that she did not know what to do to obtain relief, for 15 days. Yellow, scaly, violently itching spots over the whole abdomen and around the nipples; when scratching the parts, they become moist. * Herpes, (on the thigh.) Violent bleeding of the ulcer, almost without any cause. Fissure in the cicatrix of an old issue. Pressure and tension at the place which had been formerly occupied by an ulcer, (on the leg.) An old wart (in the face) begins to itch. Orgasm of the blood, and heat in the head. Orgasm of the blood, in the evening, before going to bed, with oppressed breathing. He feels the pulsations of all the arteries down to the tips of the toes. Sensible throbbing of all the arteries. Feeling of emptiness in the whole body, as if the body were hollow. The whole body feels heavy and broken down. The limbs feel so heavy that she finds it difficult to move a foot. Heaviness of the body, early in the morning, when in bed; going off after rising. Heaviness, especially in the feet; walking gives him much trouble. Laziness, (a. 2 d.)

WEAKNESS, FITS.—Frequent weariness and depression of strength. Weakness in the evening, almost unto nausea. **Lassitude, weakness and nausea, after the siesta.**

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Great lassitude, in the evening. Great weariness, early in the morning, when waking, abating after rising, but returning in the afternoon increased. Relaxation and weakness in the limbs. Titillating weakness in the limbs. The child (of 4 years) wants to be carried all the time. Going up a few steps is very troublesome; walking on a level is not. She is wearied by much talking. She feels weary, weak, and broken down in the hips and lower limbs, and especially in the calves, when walking, (13. d.) Weak feeling, every morning, as if he would faint, or be attacked with vertigo, (the first 6 days.) Fainting fit, as soon as he set himself in motion. Attack of weakness upon returning from a short walk, so violent that she finds it difficult to reach her home; at the same time she feels a warmth in the region of the stomach, drops of sweat are standing on her forehead, (in the winter season,) and her lower limbs are trembling; the weakness passes off after resting herself for a short while. Fit of weakness in the whole body, especially the small of the back; the cervical muscles feel relaxed, arms and legs are relaxed as if he would sink down; at the same time he feels a fainting weakness about the heart, (in a few h.) Sudden attack of weakness, in the evening, when lying down, with sick feeling, nausea, warmth and faintishness in the pit of the stomach, vertigo and vanishing of thoughts; he has two similar attacks early in the morning, leaving great weakness behind. Attack of nausea, early in the morning, with violent yawning, eructations, writhing around the stomach, violent heat and anguish, (3. d.) Attack of nausea and vomiting, with pain in the abdomen as from bruises, worse when pressing upon the part; weakness unto falling, obtusion of the head, drowsiness, and a few watery stools with subsequent costiveness. Contractive pain in the back, while resting from bodily labour; he has to lie down; afterwards profuse sweat through the night, and stool early in the morning with blood and mucus, but without any pain. Spasmodic fit; tearing sensation between the shoulders; afterwards the nape of the neck became stiff; when he attempted to move the head, it was several times drawn backwards with a jerk. The spasmodic fit was succeeded by alleviating eructations and

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

extreme depression of strength, weakness and uncomfortableness ; she was not able to talk except with a low voice. Jerkings in the limbs, (8. d.) *Slight twitchings in the muscles*, in different places, for several days. Sudden tremulousness. Violent trembling. Tremor of the arms and lower limbs ; he is easily tired by walking. * Tremulous weariness when walking, first in the knees, afterwards in the abdominal muscles and arms.

SLEEP.—Frequent yawning. Feeling as if one had not slept enough, early in the morning, looking as if he had not slept enough, whether he wakes up late or early. He sleeps too long, which causes a dreary feeling in the head; great lassitude, sensation as if one had a cold and pressure in the eyes. Early in the morning, after a sound sleep, he is obliged to lie down again soon, and feels well again after having slept three hours. **Great drowsiness by day ;* she falls asleep immediately after sitting down. Drowsiness with yawning, in the forenoon, until noon. Drowsiness after dinner, going off in the open air. Continual drowsiness in the afternoon, with yawning and wretched complexion. Great drowsiness ; she was near falling asleep again while at breakfast. Drowsiness in the afternoon, (3 d.) Unconquerable drowsiness, afternoon and evening. * *Timely drowsiness in the evening*, (a. 10 d.) Timely drowsiness in the evening, and peevish silence. Difficulty in falling asleep, in the evening, after walking in the open air. He is unable to fall asleep till late in the evening, for several days. He is unable to fall asleep before midnight, after having performed some mental labor. She went to bed at 11 o'clock at night, but she was unable to fall asleep on account of a violently stinging itching over the whole body ; she slept from 11 to 1 o'clock. At night he is unable to fall asleep till 1 or 2 o'clock in the morning, without any particular cause. Sleeplessness at night ; he has anxious dreams after he falls asleep ; in the morning, his senses are dull, and his hands hot. Restless sleep, (4. d.) Restless night, she woke up at least 20 times without any particular cause. She wakes at 2 o'clock, and remains wide awake the whole night. He wakes at 4 o'clock in the morning, and a number of times after this hour until morning. At night she wakes up, and is then prevented

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

from falling asleep again by ideas which crowd upon her mind. Half waking night-sleep; mere slumbering at night. Inclines to wake early, without being wide awake. At night, after lying down, the recollection of sad events prevents him from falling asleep. * At night, when in bed, he is delirious for three hours in a waking state, with heat in the brain, and external heat over the whole body, afterwards some sweat, coldness of the limbs and shuddering, with great tendency to start. At night she raises herself in her bed, while asleep, says all sorts of absurd things to her husband, is for a long time unable to collect her senses, but she knew that she was talking to her husband. * *Sleep full of fanciful ravings, with talking.* *Talking while asleep*, loud talking while asleep. Violent weeping during a nightly dream. At night the child tosses about and cries. * Restless sleep full of dreams. A number of dreams with restlessness and tossing about while asleep. At night he falls from one dream into another. He falls asleep immediately, but he likewise begins to dream immediately. Sleep full of dreams and interrupted by frequent waking, (a. 10 d.) He dreams of the business of the day the whole night. Voluptuous dreams, frequently, (the first 14 d.) * Anxious dream, she cries for help. Dreams about robbers, (a. 11 d.) Dreams about the sick part of the body. He dreams that one announces to him his near death. He dreams that his father is about to beat him. She dreams that she is falling down from a high mountain. Her sleep is disturbed by horrid dreams. She dreams that dangerous figures are passing near her and that some of them attempt to lay themselves upon her. She has all sorts of horrid figures before her eyes, when asleep. Dreams about serpents, illness and dead persons. Dreams about dead persons, as if they were living; he quarrels with them. Dreams about masks, ghosts and devils. * *Starting when asleep.* Starting when falling asleep. In the evening when falling asleep, he has a jerk through the whole body, which started him. In the evening after lying down, he started with his whole body, while waking, with shuddering over the whole body. He starts and trembles several times while asleep. While asleep, he twitches with his limbs and snores. For two nights in succession

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

his whole body started as if an epileptic fit would come on, with twitchings in the arms and short kickings with the feet, but without any rattling; after waking, he knew nothing of all this. In the evening, after falling asleep and waking up again, her head felt confused, she had no ideas, knew not where she was, and was then attacked with a horrible anguish, after which she came to her senses again. Congestion of blood to the head, at night, when lying in the bed, sometimes so violent that he would almost lose his senses. Aching in the head, for several nights, going off after wrapping something around the head. Gnashing of the teeth, at night, while asleep. Acidity in the mouth, at night. Gulping up of the ingesta, at night. Dryness in the mouth prevents him from falling asleep at night. At night, one hour after falling asleep, violent, jerking spasm of the stomach, with anguish, moaning, coldness of the tip of the nose, hands and feet, followed by vomiting of food and acid substances, with much rising of air; the same attack occurred in the night following, but slighter Pressure and burning at the stomach at night. Pressure below the pit of the stomach, at night, with cough. At night, she has to spit up a good deal of mucus, frequently for half hours. Colic at night, for two hours, without any subsequent stool; afterwards pain in the small of the back and chest, early in the morning; in the afternoon all the limbs feel bruised. Pinching in the abdomen, at night, in paroxysms, with nausea and constant eructations. Colic at night, from making the slightest movement in the bed, not felt when at rest, dull stitches and pressure, as from an internal induration. Three nights in succession he is tormented by flatulence. He emits a quantity of flatulence at night. Colic and diarrhœa, at night. Diarrhœa every night from 3 to 4 o'clock, (the first week.) Burning at the anus, at night, which prevents him from falling asleep. Burning itching of the perinæum at night. Sweat about the perinæum at night. Itching of the scrotum at night, which prevents him from falling asleep. A number of erections at night, during a restless sleep full of dreams. Violent erections after midnight, disturbing his sleep, exhausting him and foreboding a pollution which, however, does not take place. **In the evening, when falling asleep her throat becomes con-**

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

stricted, which causes her to start up from her sleep with fright; afterwards stinging dryness in the throat, (a. 12 d.) Arrest of breath rouses him from sleep, at night. Nightmare at night, dreaming that a stone was lying upon him, and as if he was choaked with a cord; he endeavors in vain to wake up. She wakes up at 2 o'clock at night, with anguish about the heart, and is then not able to fall asleep again. At night, when lying upon the right side, she is attacked with oppression and anguish, and has to sit up in bed, until relieved by eructations. At night, the child is uneasy and restless, cries a good deal and asks for one thing and another without taking anything. Tension in the right or left side at night. Stitches in the right or left side at night. Violent stitches in the left side of the chest the region of the heart, and sometimes extending to the back, tolerable only when lying upon the right side; they become intolerable whenever he attempts to rest upon the left side; the second night he wakes up very early, with violent stitches in the chest, and asthma when lying upon the left side; they are insufferable when in a state of perfect rest and go off when lying upon the right side; another attack in the third night when lying upon the back. Stiffness of the nape of the neck at night. At night he is roused from sleep by a throbbing pain in the upper arm. Uneasiness in the hands at night. Burning pain in the lower limbs at night, when in bed. At night when waking, tearing and drawing, with sensation of great fatigue in the legs, especially in the tarsal joints. At night he wakes up twice from a cramp in the right thigh and calf. Cramp in the calf and sole, at night, when in bed and drawing the leg up towards the thigh. At night the left leg and the right arm go to sleep. Uneasy sleep at night on account of an aching pain in the side upon which he was resting. Excessive pain in the whole body at night, like strokes of a hammer. In the evening, when in bed, she feels such an uneasiness in the limbs that she was unable to lie quietly in any one place. At night, the ulcer bleeds considerably.

FEVER.—The night-sleep is interrupted by heat over the whole body, especially the hands, and by too vivid dreams. Very hot hands in the evening, when in bed,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

with momentary shuddering, and sleep being prevented for a long time. Frequent shuddering in a room, without any subsequent heat. Frequent shuddering, with yawning, which may be relieved by the warmth of the stove, in the forenoon, (1. d.) Shuddering in the back, early in the morning, when in bed. *Very chilly*, especially after a meal and towards evening. Chilliness, in the forenoon; hot hands in the evening. Chilliness with shuddering, in the evening, before going to bed. Chilliness at every motion, even when in bed, (the first days.) Chilliness, at 9 o'clock in the evening, disappearing after lying down, without any subsequent heat or thirst, (1. d.) Chilliness, in the evening, on the top of the head and over the whole body. Violent chilliness in the evening, like fever, without thirst; for several evenings. Chilliness, as is felt when a cold sets in. Chilliness for two hours, without thirst, with obtusion of the head; when driving upon a stony road. Continual, internal chilliness, for four days, without heat or thirst, with icy cold feet, obtusion of the head, and exhaustion even unto fainting; swelling of the lower jaw and gums, and burning toothache, with stitches, (a. 32 d.) Violent chilliness with fever, towards evening, for some minutes; he is obliged to lie down; afterwards nausea and vomiting, and spasmodic pain in the chest, the whole night, with short breathing, internal anxiety, and a good deal of sweat about the head, (a. 6 d.) Internal chilliness, at noon, with heat of the hands and subsequent heat over the whole body, without thirst. Sensation of chilliness and coldness, early in the morning, when in bed; in a quarter of an hour, heat; in a few hours the chilliness returns, without, however, being followed by heat. At 6 o'clock in the evening, every day, feverish chilliness for one hour, with thirst; afterwards, heat without thirst, with violent fluent coryza; afterwards, slight perspiration with sound sleep; in the morning, scraping sensation in the throat, bad taste in the mouth, want of appetite, and agglutination of the left eyelids. Continual chilliness, with violent thirst and internal heat, hot hands, and aversion to eating, (a. 14 d.) First, chilliness; afterwards, heat in the face, (a. 2 d.) Fever, consequent upon having been heated so as to sweat, and catching cold afterwards in a

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

cold bed, with burning headache, heat in the face, and violent shuddering over the whole body, which can scarcely be endured; afterwards, sweat for three days, with such a violent fluent coryza, that it is almost impossible for him to breathe, (a. 38 d.) Feverish heat in the afternoon, and burning sensation of the eyes; (with open windows;) immediately afterwards, chilliness in the open air. Dry heat of the cheeks and hands, with short breathing. Increased warmth, at night, with violent pains in the zygoma. External and internal heat, towards morning, without any thirst, (2. d.) Heat in the evening, when in bed, without any subsequent sweat. Dry heat over the whole body, in the evening. Feverish heat, at 9 o'clock every forenoon, and at 5 o'clock every afternoon, from one half to one hour, with wide yawning, violent thirst, headache, and pulsations in the abdomen. In the evening, alternate chilliness and heat, with sweat in the night following, (3. d.) Sweat breaks out easily in day-time. Profuse sweat, when walking. Profuse sweat at every mental exertion, when reading, writing, etc. *Night-sweat*, (the first 3 nights, after 2 hours, and likewise after 6 days.) Morning-sweat, when in bed. The whole body feels moist, early in the morning when in bed, (6. d.) Sweat about the head, neck, and upper part of the body, during sleep. The pulse is slower than usual, (29. d.)

LYCOPODII POLLEN.

This dust-like powder, which is yellowish, smooth to the touch, is obtained from the ears of a moss, known under the name of *Lycopodium Clavatum*, which grows in the forests of Russia and Finland. Towards the end of summer, the ears are dried, and afterwards beaten.

When thrown into a flame, it flashes up, and has been used to cover pills which easily adhere to one another, and to protect sore places in folds against the friction occasioned by walking, and otherwise. It floats upon liquids, without being dissolved, has neither taste nor smell, and, **when in its natural crude condition, has almost no effect**

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

upon the health of man; at any rate, that which the ancients tell of its medicinal powers, has not been confirmed by moderns, but has, on the contrary, been doubted.

This drug has wonderful medicinal properties, which can only be disclosed by trituration and succussion. It is especially indicated when one or more of the following symptoms characterise the disease:—

Melancholy; *grief;* anxiousness, with soft grief and weeping mood; fear of being alone; nervousness; irritability; obstinacy; sensitiveness; *ill humor;* effects of vexations; disagreeable thoughts; impeded mental activity; ailments from mental labor; vertigo, especially when stooping; headache from vexation; aching and tension in the head; tearings in the top of the head, forehead, temples, eyes and nose, extending as far as a certain tooth, and forcing one to lie down; tearing in the forehead, to and fro, every afternoon; aching in the outer parts of the head every night, tearing, boring, scraping; heaviness of the head; congestion of blood to the head; congestion of blood to the head, early in the morning, when raising one's self in the bed, with subsequent headache; baldness, *pressure in the eyes; smarting of the eyes;* biting and burning pain in the eyes, in the evening; itching burning in the upper eyelid; stitches in the eyes, in the evening, by candle-light; *inflammation of the eyes, with nightly agglutination,* and *lachrymation* in day-time; lachrymation in the open air; viscid humor in the eye, hindering sight; *agglutination of the eyelids;* short-sightedness; far-sightedness; *dim-sightedness, as if there were feathers before the eyes;* twinkling before the eyes, and *obscuration of sight;* fiery sparks before the eyes; the eyes are irritated by the candle-light; excessive sensitiveness of hearing; music, sounds, an organ affects one's nerves; *hardness of hearing;* tingling in the ears; roaring in the ears; scurfs in the nose; nightly closing of the nostril by pus; ulcerated nostrils; bleeding at the nose; swelling and tension in the face; frequent attacks of heat in the face; itching eruption in the face; freckles in the face; toothache, with swelling of the cheeks; grumbling toothache after a meal; *dryness about and in the mouth,* causing a tension of these parts, the motion of the tongue becoming difficult,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

and the speech indistinct, with absence of thirst; coated, unclean tongue; chronic sore throat; ulcers in the throat from abuse of mercury; dryness in the fauces; burning in the fauces, with nightly thirst; hawking up of mucus; slimy taste, early in the morning; sensation as if the breath were fetid; loss of taste; *bitterness of the mouth*, early in the morning, with nausea; excessive hunger; *canine hunger; want of appetite;* vanishing of the appetite when swallowing the first morsel; aversion to boiled, warm aliments; aversion to brown bread or meat; excessive craving of sweet things; milk brings on diarrhœa; inability to digest heavy food; palpitation of the heart during the period of digestion; violent eructations, in the afternoon; greasy eructations; sour eructations; heartburn; nausea when riding in a carriage; frequent, continual nausea; *globulus hystericus;* (as if a hard ball were rising from the pit of the stomach into the pharynx;) qualmishness in the stomach, early in the morning; water-brash; *pressure of the stomach,* also after a meal; swelling of the pit of the stomach, and pain when touching it; *fulness in the stomach and abdomen;* tension around the hypochondria, as if caused by a hoop; pain in the liver, after eating to satiety; pain above the umbilicus, when touching the part; *troublesome distension of the abdomen;* indurations in the abdomen; griping, pinching, and clutching pain in the abdomen, with arrest of breathing; stitching-pinching pain in the abdomen, apparently in the bladder, extending into the urethra in the evening, when in bed; *pinching in the abdomen. pinching in the right side of the abdomen;* colic; colic in the epigastrium; tearing in both iliac regions and groins, extending into the thighs; burning in the abdomen; incarceration of flatulence; *deficient emission of flatulence;* rumbling in the left side of the abdomen; *gurgling in the abdomen;* unsuccessful desire for stool, and very hard stool; difficult stool, it can only be pressed out with difficulty; *constipation for several days; costiveness;* ascarides; pains in the anus after a meal and after stool; itching of the anus; tension in the anus; cutting in the rectum and bladder; *urgent desire to urinate;* excessive frequency of micturition, with urgent desire; *gravel;* hemorrhage from the urethra; itching in

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

the urethra, during and after micturition; weak erections; want of erections; want of pollutions; old swelling of the testes; *excessive pollutions;* want of sexual desire; impotence of several years' standing; aversion to an embrace; a desire for an embrace is kindled even by merely thinking of it; excessive desire for an embrace, every night; too quick emission of semen; *the menses are too long and too profuse;* the menses have been long suppressed by fright; *sadness and melancholy before the menses;* itching, burning and gnawing sensation of the pudendum; pressing outwards above the pudendum, into the vagina, when stooping; stitching pain in the labia when lying down; emission of flatulence from the vagina; *leucorrhœa;* leucorrhœa, preceded by cutting in the hypogastrium. *Catarrh of all kinds;* fluent coryza; coryza and cough; *dry coryza;* obstruction of both nostrils; cough after drinking; dry cough, day and night; dry morning-cough of several years' standing; difficulty in coughing something up; cough which convulses the chest; cough with expectoration; cough with discharge of pus; ulceration of the lungs; shortness of breathing in children; continual oppression of the chest, with shortness of breathing when performing some work; constant pressure on the left lower rib; stitches in the left side of the chest; pain as from bruises in the chest; sort of heartburn; anxious palpitation of the heart; stitches in the small of the back when raising one's-self after stooping; nightly pain in the small of the back; tearing in the shoulders; drawing and sensation in the nape of the neck as of being grasped into a heap, extending to the occiput, day and night; stiffness in the nape of the neck; stiffness of one side of the neck; hard swelling on one side of the neck; glandular swellings below the jaw; *drawing pain* in the arms; jerkings of the arms during the siesta; nightly pain in the bone of the arm; the arms go to sleep, even when being raised; nightly crampy, going to sleep of the arms; *weakness of the arms;* nightly pain in the bones of the elbow; arthritic stiffness of the wrist-joint; numbness of the hands; dryness of the skin of the hands; tearing in the joints of the fingers; redness, swelling and arthritic tearing of the finger-joints; stiffness of the fingers from ar-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

thritic nodosities; the fingers become rigid, while at work; the little finger goes to sleep; nightly tearing in the lower limbs; tearing in the knee; tearing in the bend of the knee, in the evening; tearing in the knees as far as above the tibia, and in the dorsum of the foot; *stiffness of the knee;* swelling of the knee; burning of the legs; burning-biting, itching of the bends of the knees; *contractive pain of the calves, when walking;* old ulcers upon the legs, with nightly tearing, itching and burning; swelling of the malleolus; cold feet; cramp of the feet; cold, sweaty feet; profuse sweat of the feet; swelling of the soles of the feet; pain of the soles when walking; the toes bend suddenly when walking; cramp of the toes; corns; pain of the corns; *dryness of the skin;* the skin cracks in different places; itching in day-time, when being heated; itching in the evening previous to lying down; painful eruption on the neck and chest; itching and corrosion of the arms and legs; *boils;* cramp in the fingers and calves; spasmodic contraction of the fingers and toes; tearing in the upper and lower extremities; tearing in the knees, feet and fingers; pain in the bones from abuse of mercury; drawing pain in the limbs; uneasiness in the feet, in the evening; the arms, hands, lower limbs, go to sleep, both in day-time, and during the night; insensibility of the arm and foot; varices; *varices of pregnant females;* parts become strained easily, bringing on a painful stiffness in the nape of the neck; jerkings and startings of single limbs, or of the whole body, when asleep or waking; troublesome to rest on the left side, on account of palpitation of the heart and stitches; liability to cold; weariness of the feet, and burning of the soles, after a little walk; *internal weakness;* faintness of the limbs; weariness when waking up: frequent yawning and drowsiness; drowsiness by day; restless night-sleep, with *frequent waking;* sleep full of dreams; sleep full of ravings; fearful, *anxious* dreams; *falling asleep late at night;* thoughts crowd upon him and prevent him from falling asleep; jerkings and impatience in the feet at night; nightly headache; starting up when asleep; want of animal heat; thrills of heat; tertian fever with sour vomiting after the chilliness, and bloat-

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

edness of the hands and face ; feverish sweat by day ; *day-sweat*, from moderate work or little exercise, especially in the face.

Camphor generally moderates the violent effects of lycopodium ; Pulsatilla subdues the violent feverish feelings from lycopodium ; the ill-humor, readiness to find fault, diffidence, tendency to reproach, &c., are subdued by Causticum. A cup of coffee prevents and completely neutralizes the action of lycopodium.

A moderate dose acts from 40 to 50 days. It may be repeated after the intermediate use of another anti-psoric, but a second dose acts less favorably than the first.

It acts with especial benefit, when it is homœopathically indicated after the action of Calc. shall have passed over. This drug has been tried by me and my colleagues Drs. Gersdorff, Goullon, Hartlaub, Rummel, Schreter, Wahle, Rückert.

From Noack and Trinks : This drug has an especial action upon the sensorium, the splanchnic nerves and the organs over which they are distributed, the intestinal canal, the liver, the uropoëtic and sexual organs of both sexes, the lungs, nervous and fibrous tissues, the skin, and perhaps also the periosteum and the bones.

Lycopodium is especially adapted to individuals of a soft, mild, melancholy disposition, to lymphatic and leucophlegmatic constitutions, and when they are moreover subject to mucous discharges, coryza, catarrh, &c.—Diseases with aggravation of the symptoms in the afternoon and in a room, amelioration and mitigation of the same in the open air.—Great sensitiveness to open and cold air.—Typical, periodical diseases.—Diseases with paroxysms of pain, causing great uneasiness, walking about and weeping.—Disease with difficulty of animal heat as a prevailing symptom.—Diseases based upon obstructions, stases, and plethora of the portal system.—Diseases with inclination to dropsy and affections of the urinary organs.—Bad consequences from vexation : great weakness unto falling, palpitation of the heart, trembling, heaviness in the limbs, as if they were made of lead.—Diseases from dyscrasia.—Scrofulosis.—Scrofula of the bones.—Rickets.—Secondary syphilis.—Condylomata.—Liability to infection after being

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

cured from syphilis.—Periostitis and ostitis with nightly pains, even when brought on by mercury.—Spina ventosa. Scrofulous caries.—Mercurial caries.—Fistulous ulcers in the bones.—Gout.—Arthritic affections, even nodosities.—Rheumatic fevers.—Drawing and tearing in the limbs, in periodical attacks, when the weather is rainy and stormy.—*Great liability to cold.*—Parts are easily strained.—Stiffness in the joints and limbs, with numbness and insensibility.—Paralysis.—Uneasiness in the body.—Orgasm of the blood.—Spasmodic conditions: convulsions in alternate parts; involuntary turning and twisting of the whole body.—*Spasms of the muscles.*—Attacks of spasmodic extension and contraction of the limbs and of single muscles, without pain and with entire consciousness.—Contraction of the fingers and toes.—Cramp in the fingers and calves.—Epilepsy.—Rhagades.—Itching and painful little nodosities on the occiput, os sacrum, and nates.—Scabies vesicularis humida.—Large, red spot on the epigastrium, pit of the stomach and joint of the thumb, itching and burning. Hepatic spots.—Impetigo figurata.—Lichen agrius.—Freckles.—Urticaria febrilis chronica.—Psoriasis.—Psoriasis simplex.—Psoriasis palmaris.—Lupus vorax.—Herpes; small, herpetic, itching spots, itching spots on both sides of the neck and on the back.—Violently itching herpes. Insensible, yellow-brown, withering herpes.—*Humid suppurating herpes, full* of rhagades and covered with thick crusts.—Herpes squamosus.—Herpes ecthyma.—The herpes becomes worse when Lycop. has been given improperly; it becomes ichorous and spreading; in this case Sepia affords relief.—Periodically recurring boils on the nates, scapulæ, arms, &c. Boils in and around the axilla.—Furuncles after typhus.—Intertrigo of infants and full-grown persons.—Warts.—Chilblains.—Old, malignant ulcers on the feet, with hard, shining, elevated edges, copious, watery, ichorous, extremely fœtid pus, blue-red, hard, hot, painful areola, big-swollen, shining, red leg and foot, and tearing, lancinating pains, increased at night and by bad weather.—Phagedenic ulcers.—Salt rheum.—Dropsical conditions.—Dropsical conditions with œdema of the feet and diminished secretion of urine.—Chlorosis.—Chlorotic colour of the skin.—Intermittent fe-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

vers.—Intermittent fevers with constipation.—Quotidian and tertian fevers.—Consumptive fever with viscid night-sweats.—*Nervous fevers*, characterised by ill-humor, when waking, by coryza, screaming, improper manners and constipation.—*Exhausting night-sweats.*—Hysteria and hypochondria.— Melancholy.—*Absence of mind.* — Vertigo, with the loss of thoughts, confused talking, rheumatic pains in the face, teeth, back, small of the back, and limbs. Headache from vexation, with great uneasiness and faintishness.—*Nervous, chronic headache.*—Megrim.—Hemicrania neuralgica.—Rheumatic headache.—The hair turns white.—Falling off of the hair, after a heavy illness during confinement.—Pityriasis ; scald-head.—Tinea capitis favosa.—Eruptions on the head, with violent suppuration, covered with coherent, thick crusts, forming one mass with the hair, violently itching, with swelling of the glands in the neck and nape of the neck, humid tinea behind the ears and pustules in the nape of the neck.—Plica Polonica.—Spasmodic twitchings and jerkings of the eyelids.—Hordeoli.—Inflammation of the eyelids, with swelling and redness of the same ; profuse secretion of acrid tears and mucus ; nightly agglutination of the eyelids and the canthi.—Inflammation of the eyelids.—Ophthalmia neonatorum.—Catarrhal, rheumatic, and scrofulous ophthalmia.—Arthritic ophthalmia, with formation of pannus.—Fungus medullaris cornea. — Synechia anterior. — Weakness of sight.—Obscuration of the crystalline lens from amenorrhœa. — Cataract ? — Muscæ volitantes. — Amaurosis. — Purulent otorrhœa, with hardness of hearing and humid eruption on the head.—Caries of the ossicula aurium, and of the internal ear, after scarlatina.—*Chronic* wheezing in both ears, with diminution of hearing, in the case of a woman, who had suffered with it for 5 years past. Scrofulous, mercurial ozæna. Inflammation and suppuration of the frontal cavities and the antrum Highmori.—Catarrhal inflammations of the Schneiderian membrane, of the mucous membrane of the frontal cavities, with tearing pains in the head and secretion of lemon-yellow mucus.—Chronic blennorrhœa of the Schneiderian membrane.—Polypi in the nose.—Wretched, sunken, small, pale countenance.—Blue margins around the eyes, with clay-colour-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

ed countenance, deep wrinkles, and blue lips.—Circumscribed redness of the cheeks.—Contractive and tearing pains in the face, cheek, and bones of the face.—Spasmodic distortion of the muscles of the face.—Corrosive, biting, and stinging pains in the right half of the face.—Nervous prosopalgia.—Rheumatic swelling of the cheek and face.—Itching in the face.—*Itching, suppurating eruptions in the face, covered with elevated crusts.*—Pimples in the face.—Unclean skin of the face.—Itching herpes near the nose.—Scaly herpes in the face and corners of the mouth.—Herpes farinosus in the face.—Herpes furfuraceus, with a yellow base.—Crusta lactea.—Crusta serpiginosa.—Pale, bluish lips.—Spasmodic movements of the lips and corners of the mouth—first, distensive—then contractive.—Swelling of the lips.—Sore and ulcerative pain of the corners of the mouth.—Eruption around the mouth.—Ulcers in the vermilion-border of the lower lip.—Cancer of the lips.—Itching pimples on the chin.—Sycosis menti.—Lock-jaw.—Spasmodic movement of the lower jaw.—Swelling of the sub-maxillary glands, with a boring pain.—Toothache, with swelling of the cheeks.—Drawing, stinging, *grumbling*, boring, *burrowing*, beating, throbbing pains in sound and hollow teeth.—Pains in the teeth, as if there were sub-cutaneous ulceration, or as if they were dashed to pieces, with or without swelling of the cheek and gums.—Looseness and elongation of the teeth.—The teeth become yellow.—Swelling of the gums.—Fistula dentalis.—Bleeding of the gums.—Fetid odour from the mouth.—Ptyalism.—Spasmodic movements of the tongue.—Heaviness of the tongue.—Soreness and rawness of the tongue.—Vesicles on the tongue.—Aphthæ.—Coated tongue.—Exudations of the mouth and fauces.—Chronic inflammation, and induration of the amygdalæ.—Ulceration of the tonsillæ.—Angina gangrænosa with stupor, in scarlatina.—Chancre-like ulcers on the tonsils and in the pharynx, also from abuse of mercury.—Syphilitic and non-syphilitic ulcers of the palate and the tonsils.—Troublesome feeling of dryness, and actual dryness of the throat.—Syphilitic herpes in mouth and fauces.—Syphilitic affections of the throat—Bitter, sour, slimy taste.—Loss of taste.—Want of appetite.—Desire to eat.—Canine hun-

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

ger.—Gastric affections.—Difficult, slow digestion.—Violent, empty, and sour eructations.—Heartburn.—Waterbrash.—Vomiting of food and bile.—Chronic vomiting.—Chronic gastritis.—Gastrodynia; enterodynia.—Chronic affections of the stomach and abdomen.—Pressure of the stomach.—Contractive spasm of the stomach.—Spasm of the stomach and abdomen.—Cancer of the stomach.—Chronic inflammation of the liver and spleen.—Chronic irritation of the mucous membrane of the intestines.—Peritonitis.—Cutting, pinching, drawing, constrictive pains in the abdomen; also after a cold.—Atrophia miseraica infantum.—Spasms in the abdomen.—Flatulence.—Flatulent colic, with cold feet.—Ascites.—Inguinal hernia.—Incarcerated hernia.—Swelling of the inguinal glands.—Constipation.—Dysentery.—Prolapsus recti.—Hemorrhage from the rectum.—Protrusion of the varices of the rectum, with itching, aching and burning pains in the rectum and anus.—Hæmorrhoidal affections.—Herpes at the anus.—Hemorrhage from the kidneys.—*Diseases of the urinary organs.*—It effects the discharge of gravel and stone, and appears to remove the disposition for their formation.—Colica renalis from gravel.—Ischury, disury, and strangury.—Retention of urine in children.—Polypus of the bladder.—White fig-warts, fissured at the top and on stems.—Suppuration of the kidneys.—Gravel.—Hematuria, with obstinate constipation.—Itching, cutting, and burning in the urethra.—Blennorrhœa from the urethra.—Secondary gonorrhœa.—Suppurating ulcers on the glans and prepuce, not syphilitic, in conjunction with hæmorrhoidal herpes.—Soreness between the scrotum and thigh.—Impotence, with pollutions brought on by onanism.—Impotence caused by the vapour of chlorine.—*Exhausting and excessive pollutions.*—Chronic dryness of the vagina.—Suppressed pollutions.—Promotes fecundity.—Prolapsus uteri.—Inclination to miscarriage, with varices of the pudendum.—Nodosities and indurations in the mammæ.—*Soreness of the nipples*, or corrosive scurf on the nipples.—*Intertrigo neonatorum.*—Constipation of new-born children.—Weak, hollow voice.—Hoarseness in croup.—Rattling and râling in the chest.—Pneumonia.—Neglected pneumonia, threatening to become fatal.—Chronic catarrh.—Influenza.—

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Chronic, dry cough.—Cough, with bloody expectoration.—Cough, with discharge of salt mucus, or green purulent substances.— Incipient phthisis tuberculosa.-- Hæmoptysis.—Cough, with yellow purulent discharge.—Cough, with expectoration of mucus.—Pituitous phthisis.--Phthisis purulenta, cavernosa.—Spasm of the chest.—Asthmatic affections.—Palpitation of the heart.—Aneurism.—Aneurism of the external iliac artery.—Painful eruption on the chest.--Hepatic spots on the chest.—Spasmodic movements of the head: involuntary nodding and shaking, stretching it forwards and backwards; extension and contraction of the muscles of the neck.—Painful eruption on the neck.—Large nodosities, formed of red pimples, around the neck, itching violently. — Swelling of the cervical glands.— *Struma*.— Caput obstipum.— Neck spasmodically drawn to one side, with spasmodic distortion of the facial muscles.—Drawing and stiffness of the neck.—Rheumatic, *tearing* pains in the upper limbs.—Spasmodic jerking upwards at times of one, at times of the other shoulder, and of the arms, during the siesta.—Spasmodic shaking of the hands.—Cramp of the fingers.—The arms go to sleep and feel paralytic.—Chiragra.—Dryness of the skin of the hands.—Sweaty palms of the hands.--Rheumatic, *tearing*, drawing, and darting pains in the lower limbs.—Ischias?—Phlegmasia alba dolens?—Gonagra?—Stiffness of the knee.—White swelling of the knee.—The lower limbs are involuntarily spread asunder, and then pressed against one another.—Cramp of the calves and toes.—Spasmodic contraction of the fingers and toes.—Podagra.—Varices.—Œdema of the feet.—*Icy coldness of the feet.*—Sweaty feet until they become sore.—Restores a suppressed sweat of the feet.—Corns.

Moral Symptoms.—Hypochondriac, tormenting mood; he feels unhappy, (the first 2 d.) Desponding, grieving mood, (a. 17 d.) The child loses its cheerfulness, becomes silent and discouraged. Desires to be alone. *Dread of men*, (1. d.) When men approach her, she feels a sort of anguish in the pit of the stomach. She avoids the society of her own children. * Melancholy, in the evening. Want of cheerfulness, sad thoughts. Sad mood: she had to weep the whole day, and found it impossible to

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

calm herself, without any particular cause. Sad, desponding, finally weeping mood. Extremely sad and low-spirited. *Weeping mood*, with chilliness. He weeps first on account of the past, and then on account of future evils. *Great apprehensiveness in the pit of the stomach, from vexation. Great anxiousness, apparently in the pit of the stomach*, without any particular thoughts, (a. 24 h.) Internal anguish, in the forenoon, and internal chilliness, resembling an internal trembling. *Anxiousness, in the evening*, her sight is half confused. Anxious, fearful, timid. Great fearfulness, (10. d.) She is very much afraid of phantoms, crowding upon her fancy in the evening, with weeping mood during the day. In the evening, when attempting to open a door which is rather tight, in the dark, he is assailed by fear. In the evening, when stepping into the room, he imagines he sees somebody, which gives him fear; also in day-time, he sometimes imagines he hears somebody in the room. * She is afraid of being alone. Internal uneasiness, (a. 24 h.) Impatience. Discouraged and faint. Want of confidence in his strength. *Pusillanimous, sad*, fanciful. Distrustful, suspicious, inclining to find fault. Extreme suspiciousness and distrust. Despairing and disconsolate. * Extremely sensitive; she weeps when people thank her for a kindness, (a. 20 h.) Extremely irritable, frightful, and out of humor. *Great tendency to start.* Every noise hurts her. *Dissatisfied*, (a. 72 h.) Very irritable, and inclining to be melancholy. *Ill humor.* Out of humor and desponding, (15. d.) A number of past disagreeable occurrences crowd upon the mind, even at night, when waking; they put her out of humor. He finds it difficult to conceal his internal obstinacy and vexed mood. Excessive irritability, with apprehensiveness. He smiles without being cheerful; it is mere caprice. The child becomes less docile, although not ill humored. Insolent, self-willed, * headstrong,—vehement, angry. Very vehement. Vehement mood, without ill humor, (in a few hours.) She cannot bear the slightest contradiction, and is immediately beside herself with ill humor. Angry rage, partly against herself, partly against others. One is easily irritated and angry. He disputes with absent per-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

sons. Frenzy and madness, breaking forth in envy, reproaches, pretensions, and commanding spirit, (a. 12 d.) She seeks disputes as if she were mad, scolds one in the most vehement manner, and strikes the person whom she scolds, (a. 2 h.) Ennui, (a. 2 d.) The mind is insensible to external impressions. Irritated mood, but indifferent to external impressions. Extremely indifferent. Want of desire to talk. Inclines at the same time to weep and laugh. Anxiousness, followed by great desire to laugh at trifles, for some hours, after which she weeps for half an hour without knowing why. Excessively merry, with turning vertigo. Wantonly merry. She laughs when one looks at her, while telling something serious. Involuntary whistling and humming with the lips. Excessive mirth, as if he were obliged to distort the muscles of the face, succeeded by want of cheerfulness and by impatience. Weakness of memory, (a. 3 d.) Acting as if he were absent. Living as it were beside one's-self, as at the commencement of a fever. When thinking, his head feels empty ; he is unable to fix one idea. He is unable to do any thing, or to think ; he spends his time in doing insignificant things, without being able to do that which he has to do. * He is unable to retain his idea ; it is difficult for him to express himself and to find suitable words, especially in the evening. She is unable to conceive an idea, owing to obtusion of the head with internal tension. His thoughts feel as if they were arrested, the mind feels clumsy and rigid, there is obtusion without gloominess. He is capable of talking logically about higher, abstract things, but he becomes confused when talking about common-place things : for instance, he says plums instead of pears. *Using wrong words and syllables.* He is unable to read, because he mistakes and confounds the letters ; when seeing them, he is able to copy them, but he does not recollect their meaning ; he knows, for instance, that Z is the last letter in the alphabet, but he has forgotten its name ; he is able to write any thing he chooses ; he writes the proper letters, but he is unable to read that which he has written.

Head.—Obtusion of the head, as if he could not collect his senses. Obtusion of the head, as if from spoiled

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

stomach. Obtusion of the head, with pressure over and in the eyes, in the evening. Gloominess of the head, with dull pressure in the fore part of the head, as if a cold had been repelled, with dryness of the lips and mouth, and with thirst. Dizzy, early in the morning, as if one were intoxicated. Dizziness, she could not tell where she was. Dizziness, indolence and relaxation of her limbs, the whole of the second day. Stupor, towards evening, with heat of the temples and ears. She felt as if every thing before her would pass away, (3. d.) * Reeling sensation, early in the morning, and sensation of heaviness in the eyes. As soon as she sees something turn about, she feels as if the contents of her body were likewise turning, for one hour. Vertigo like obtusion of the head, in the forenoon, with sensation as if her eyes were deep in the orbits; thinking and comprehending are likewise very difficult. Vertigo, in the forenoon; every thing turned with her, accompanied by violent nausea. Vertigo when rising from a seat. Vertigo when drinking. Vertigo in a hot room, (a. 23 d.) Vertigo, early in the morning, while getting out of bed and afterwards, so that he staggered to and fro, (a. 30 d.) Simple headache, for several days, more violent when at rest, less when walking in the open air. Headache, especially when shaking and turning the head. Violent headache, at night, as if caused by a wrong position. Headache on the left side, apparently on the outside, reaching also the ear and teeth, especially violent in the evening, and becoming insupportable from writing and reading, or by the least pressure on the temples. *Headache over the eyes, immediately after breakfast*, (the first two days.) Headache between both eyes. Pain in both temples, at every step, not when at rest. She feels every step which she makes, in her head; at every motion she feels a sort of concussion in her brain. Dull pain in the forehead, as if the head were being compressed from both sides. Pressure as from a nail, in the middle over the hairy part of the forehead. Pressure in the forehead and down the nose, early in the morning. Pressure, at times in the right, at times in the left temple. Pressure in the right half of the occiput, towards the ear. *Pressure in the nape of the neck, at a small spot.* Violent pressure in the nape of the neck, for

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

many days. Paralytic aching in the temple. Aching in the top of the head, as if a cold would come on, (a. 12 h.) Aching in the whole head, in the afternoon, especially when stooping. Aching, with contractive sensation in the head, the former prevailing early in the morning, when rising. The headache increases when lying down. Headache, as if the bones of the skull were being driven asunder, and as if the brain were vacillating to and fro, especially when walking, going up stairs, and raising one's self after stooping. Sense as of being pressed asunder, in the top of the head, with nausea, as if she would throw up every thing, with trembling of the limbs, (a. 1 h.) * Heaviness in the head. Heaviness of the occiput. Dull feeling of heaviness in the occiput, with confused pain in the forehead, increased by motion. Pain in the head, at night, moving about and making the head feel heavy; it felt so painful that she did not know where to place it, in order to obtain some rest; the whole night, (a. 9 d.) Pinching pain about the head, behind the ear, (a. 48 h.) * Tearing in the occiput. Tearing in (about) the head, for 48 hours, ceasing after the appearance of a painless swelling of the cheeks. Tearing headache, in the evening, on the top and on both sides of the head. Tearing, in various places, in (about) the head; afterwards in other parts of the body. Tearing in the head, (4. d.) Tearings through the left side of the head as far as the ear. Sharp irradiating tearing in and above the left frontal eminence, towards the left side. Irradiating tearing, in jerks, in the right half of the head, commencing at the temple. Tearing with pressure, in jerks, in the right half of the forehead, extending close to the root of the nose, and the right eye-brow, apparently in the bone. Tearing with pressure in (about) the left side of the occiput, at a small place near the nape of the neck. Aching, with tearing close over the eyes and into them, early in the morning, (3. d.) *Tearing* and lancinating headache, *at night*, over the right eye, in the temple, and about the occiput. Stitching headache in the region of the eye. Stitching pain in the occiput. Stitches in the forehead, from within outwards, coming with a jerk, frequently during the day. Violent stitching pain in the left side of the forehead. Tearing headache from after-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

noon till evening: toothache in the subsequent night. Single stitches in the head, worst at night, causing one to start. Stitches and pressure in the head (in a few hours.) Stitches and pressure in the top of the head, at night, (a. 7 d.) Stitching headache with pressing and pushing in the eyes, with violent fluent coryza. Sore pain in the forehead, almost every day, aggravated when stooping. Violent shock from the back to the top of the head, which obliged him to hold his head, when sitting, (after a copious meal.) Jerking sensation in the head, with sensation as if something heavy were balancing to and fro in the head. Darting headache, apparently in the bones of the skull. Throbbing pain near the orbits, from within outwards. *Throbbing in the head*, in the evening, after lying down. Throbbing headache, after every attack of cough. Throbbing in the brain, when leaning the head backwards, in the day. Violent beating in the head, resembling a hacking (with sour eructations.) Violent throbbing in the forepart of the head, in the evening, afterwards extending over the occiput, down to the nape of the neck, with a feeling of tension, (a. 4 d.) Throbbing in the top of the head. Continual, throbbing headache, in the centre of the forehead, from 3 o'clock in the morning until evening. Beating and pressure in the occiput. Pulsations and pressure in the head, when reading, while sitting. Beating in the brain, with heat about the head. * *Rush of blood to the head, early in the morning, when waking.* *Fulness of blood in the* occiput, after stooping. Headache, like a sound through the head, as from the breaking of a musical string. *Groaning in the head, when setting the foot down with force.* Whizzing in the head, with a feeling of heat. Excessive sensitiveness of the head externally. Painfulness of the forepart of the head, on the outside, when touching it. Superficial pain above the forehead, on the top of the head, in the region of the malar bones, the ear, the jaw-bones; intermitting in the afternoon, returning in the evening. Quickly passing tearing in the outer head, when walking in the open air. Tearing in the hairy scalp, above the right half of the forehead. Fine, burning, stinging tearing in the skin of the right temple, after itching and rubbing; afterwards

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

clawing headache. Drawing pain in the right side of the head, down the nape of the neck. Sensation on the left side of the hairy scalp, as if a single hair were being pulled at. Contractive sensation in the hairy scalp, with sensation as if the hair were being pulled out. Spasmodic contraction of the scalp. Drawing up of the skin of the forehead, with consequent receding of the eye-lids one from the other, and subsequent drawing downwards of the same skin, with closing of the eyes. Cutting pain transversely across the hairy scalp, between forehead and vertex. The bones of the head are painful. Burning pain in both occipital protuberances. * *Excessive falling off of the hair.* Falling off of the hair of the head, whereas the hair on other parts of the body grows more abundantly. *She has a number of grey hairs.* Great liability of the head to cold; a cold little breeze causes a cutting sensation in the hairy scalp. *Itching of the hairy scalp.* Corroding sensation of the hairy scalp; he is obliged to scratch himself; * Eruption on the head, with swelling of the cervical glands; large pustule on the occiput, and a scurf over the hairy scalp, which is scratched open by the child, and then bleeds. Broad tubercle under the skin of of the forehead, without the colour of the skin being changed at that place. Tumour on the occiput, of the size of a walnut, (7. d.) * Eruptions on the head, suppurating profusely. His head is turned involuntarily to the left side.

Eyes.—Pain in the eyes, in the evening; she finds it difficult to open them. The eyelids are painful when touched. Pain of the eye, as if it had been bruised. Pressure in the internal canthi. * *Aching in the eyes*, as if dust had got in. Pressure on the eyes, with drowsiness, in the forenoon. Pressure on the right upper eyelid. Pressure in the right eye, as if something had got into it. Pain in the eyes as if they had been bruised, and as if they would fall out; the pain was so violent that he was unable to look at any thing fixedly, from 1 o'clock in the afternoon, but mostly in the evening. He is unable to raise the eyelids; the lids are too heavy. Heaviness of the lids, also in day-time, especially when the light is very dazzling. The eyes feel tired and heavy; they look

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

sleepy. Tensive pain in the left eye. Pressing together of the eyes, the skin over the malar bones feeling tense. Tearing around the eyes, extending into the forehead and the cheeks. Tearing in the right eyeball. Stitches in the left eye. * *Stitches in both eyes*, (a. 12 d.) *Stitches in the eyes without any redness*, the whole day, but especially early in the morning, (a. 34 d.) Prickling, at times in one, at times in another eyeball. Itching in the eyes, (a. 30 d.) *Itching in the canthi.* Biting in the right eye, as if caused by smoke, with closing of the eyelids. Biting sensation in the external canthi, with lachrymation as if caused by smoke, in the evening at twilight. Itching around the eye. Cold feeling in the eyes, in the evening. *Burning in the eyes.* Burning in the eyes, when he was about to close them. Violent burning and itching in the eye. *Redness of the eyes*, with pressure in the same. Redness of the white of the eye, with pain. Red, inflamed eyes, with stitching pain, from 5 o'clock in the afternoon until 10 in the evening. Inflammation of the white of the eye. * Inflammation of the eyelids, with aching, and nightly agglutination in the outer canthi. * Inflammation of the eyes; with redness and dimness of the white, redness and swelling of the lids, burning, pressure and secretion of mucus in the eye. Inflammation of the eyes with redness of the white and swelling of the lids, stitches, photophobia, profuse lachrymation and nightly agglutination. Inflammation of the eyes, with itching in both canthi, redness and swelling of the lids of the right eye; tensive pain when they had become dry, and nightly agglutination. Swelling and painfulness of the eyelids, with nightly agglutination in the canthi. A number of pustules in the eye-lids. *Styes on the eye-lids*, near the internal canthus. Ulceration and redness of the eye-lids; the water which oozes out, causes a corroding and smarting sensation on the cheeks. Suppurating stye on the eye-lid. Red pimples on the upper lid of the right eye, combining into a sort of scurf. * *Agglutination of the eyes*, especially at night, and more particularly in the outer canthi. In the morning the eye-lids are agglutinated. *A quantity of purulent gum in the eyes*, with smarting pain, (a. 32 d.) Eye-gum in the internal canthus, early in the morn-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

ing. A corrosive humor flows out of the eye, the white being very red. *Mucus in the eyes ; he has to wipe them, in order to see more brightly.* Lachrymation and a quantity of gum in the eyes, with pressure and pale face. Profuse lachrymation of the right eye, in the afternoon. Water runs out of the eye, when exposed to a rough wind. * Dryness of the eyes, in the evening. Dryness of the eyes; he has to close the lids. Dryness under the eyelids, as if there were dust, early in the morning, when waking. * Dim, hot eyes. Faint eyes, in the evening, by candle-light, painful when turning them. Spasmodic twitching of the lower eye-lid, (a. 90 d.) Spasmodic twitchings of the lower lid of the left eye, towards the internal canthus, (a. 35 d.) Slight twitching of the left eyelids. Weakness of sight; she is unable to see or to sew long ; she is obliged to close her eyes from pain ; in the morning they become somewhat agglutinated. The letters become indistinct when writing. The letters become blurred when reading. Uncertain sight and frequent twinkling before the eyes. His eyes become dim, even at a very short distance ; sensation, as if he looked through a fine wire-gauze. Dim-sightedness, as if there were a viscid humour in the eye, which cannot be wiped away, and which increases and decreases in paroxysms. Farsightedness ; when reading or writing every thing seems indistinct to her, as if it were covered with a gauze ; things at a distance appear bright and distinct. He only sees the left half of the objects, the right is either wanting, or else it is obscured ; when using but one eye he sees the same as with both, except that this peculiar symptom is more strongly marked on the right than the left eye. Dimness of sight, he is obliged to hold the writing at times near to, at times far from, his eye, in order to be able to read. *Black spots hover before his eyes at a short distance,* (in 41 h.) Gauze and twinkling before the eyes, after the siesta, (a. 16 d.) * Twinkling before the eyes, when going to bed. Sensation of vibrations before the eyes, resembling those of the hot air of summer. Tremor of the objects at which she looks attentively, in the evening, by candle-light, the light trembling most, when looking at it. * Sparks before the eyes, in the dark, (a. 5 h.) His eyes

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

are very much dazzled by the evening-light, he is then unable to see any thing upon the table.

Ears.—*Otalgia* in the open air. Sensation as of pressing towards the ears. Sensation as if something had been forced into the internal ear. Pressure on the posterior surface of the right concha. *Tearing in both the right and left meatus auditorius.* Tearing behind the left ear. Tearing in the left concha. Tearing about the right ear, (14. d.) Darting in the internal ear. Stitches in the ear, when blowing the nose, the speech being more difficult. Biting sensation and sore feeling behind the right ear and by the side of it. Stitches in the ear. Continuous, tearing, pinching stitches in the ear, which feels as if it were too narrow and as if it would burst. Throbbing and tension in the ears, with spasmodic tension of the skin behind the ears, in an oblique direction towards the muscles of the nape of the neck. *Rush of blood to the ears.* Sensation as if hot blood rushed into the ears. Itching in the ear. Ulceration and running of the ears. Pain behind both ears, which obliged him to walk crooked. * *Sensitiveness to noise* when walking. * Diminution of hearing, (a. 24 h.) The sound of speech seems hollow, although as strong as usual. His ears become closed, with whizzing and hardness of hearing, (in 10 d.) Her ear became closed, with a fluttering sensation. In the evening she hears the music in her ears, which had been played to her during the day. Sounds in the ears. Roaring, humming, and whizzing in the ears. * *Violent whizzing in the ears.* Sense as of seething in the ears, (2. d.) Whistling in the ear, when blowing the nose. Chirping in the ears, for several evenings. Beating in the ears, morning and evening. Guggling in the ears, by day. Gurgling as of bubbles of air in the ears.

Nose.—The muscles of the nose first appear distended, then again contracted and shortened, like the brim of a hat turned over. Corrosive pain in the right nostril. Corrosive pain in the left nostril, when moving the nose or inserting the finger into it. Itching of the nostrils. Itching of the nose, (a. 5 d.) Pressure on the nasal bone, near the right eye. Drawing, with pressure, on the right side of the nose. Tearing, commencing on the right side of the nose, and coming out at the corner of the eye. Cut-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

ting, sore feeling of the upper part of the septum, in the right nasal fossa; in the evening, when in bed. Swelling of the tip of the nose, with pain to the touch. Heat in the nose, and burning of the eyes. *The smell is excessively sensitive;* even the smell of hyacinths makes him feel nauseated. Increased sense of smell. Total absence of smell, (a. 2 d.) Smell of lobster, when spitting. Expulsion of a bloody mucus from the nose, (a. 6 d.) Expulsion of coagulated blood from the nose, (a. 11 d.) Considerable bleeding from a little wound in the nose, in the evening, when walking, (a. 32 d.) Bleeding at the nose, three days in succession, at 2 o'clock in the afternoon. Two bleedings from the nose in one day, (a. 26 d.) Considerable bleeding from the nose, frequently followed by expulsion of blood from the nose when blowing it, (a. 20 d.)

Face.—Paleness of face, with drowsiness in day-time and ill humour. The paleness of the face increases towards evening, (a. 8 d.) Very pale, sunken countenance, early in the morning. * *Pale, wretched complexion.* Paler and narrower countenance. The countenance is changed, with sunken eyes. Sunken eyes, (a. 7 d.) * Blue margins around the eyes, (a. 12 d.) * *Yellow countenance.* * Yellow-gray complexion. The whole countenance first extended in length, afterwards in breadth. Heat in the face, eyes and palms of the hands. Heat in the face, with hypochondriac mood. Flushes of heat in the face, early in the morning, shortly after rising. * *Frequent flushes of heat,* (the first days.) Violent heat in the face, without redness. Striking redness in the face, early in the morning. Burning in the face, (a. 26 d.) Red, bloated face, full of dark-red spots covered with pustules. * Swelling of the cheeks. * Eruption in the face, (a. 12 d.) Single pimples on the face. Itching of the face, head, and nose. Itching all over the face, and pimples with pus in the tips, upon the cheeks, forehead, and especially on the temples, (a. 12 d.) A number of pimples and freckles over the whole face. There is a larger number of freckles on the left side of the face and over the nose. The face looks as if it were covered with a fine eruption. Itching herpes on the side of the nose, near the eye. * Itching, scaly herpes in the face and in the corners of the mouth,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

with bleeding. Simple pain in the left side of the face, when touching it. Contractive pain in the muscles of the face and the occipito-frontalis muscle, (a. 4 d.) Sensation of swelling on the forehead. Tearing in the cheek. *Tearing in the malar bone*, below the left eye.

JAWS AND TEETH.—*Tearing in the upper jaw bone*, (2. d.) Tearing in the right upper jaw. Spasmodic twitchings in the muscles of the cheeks. The muscles of the lips and cheeks contracted, and made the mouth look pointed; this was followed by extension of the mouth. The left corner of the mouth was drawn upwards; afterwards the right corner became distorted. Paleness of the lips. Swelling of the right half of the lower lip. Swelling of the lips, early in the morning. Swelling of the upper lip, increasing for several days, lastly accompanied with evening-fever—first, chilliness, afterwards heat in the face, in the hands and feet, restless sleep and night-sweat. A sore place on the lower lip. Soreness of the corners of the mouth. Ulcerative pain of the corners of the mouth. Eruption around the mouth. Fine eruption around the mouth, (11. d.) Itching pimple on the upper lip, (a. 14 d.) Eruption on the margin of the vermilion border of the lip, with cutting pain when moving the lips, or when touching them, (a. 12 d.) A white blister on the inner side of the upper lip, with burning pain when at rest—not when eating, (a. 30 h.) A large ulcer *on the vermilion border of the lower lip*. *Itching pimples around the chin*. Pressure on the right lower jaw, towards the back part. Drawing pain in the right lower jaw; the sub-maxillary glands feel heavy, as if they were swollen, with throbbing in the glands, increasing after walking or eating. * Drawing in the jaws. Darting pain in the lower law, in the evening. Jerking tearing, at times in the right, at times in the left lower jaw. The lower jaw is involuntarily pushed backwards and forwards. Hard swelling in the corner of the lower jaw, with a feeling of heat in the head. Boring pain in the swollen sub-maxillary glands, (a. 4 d.) Disagreeable toothache; she would like to bite her teeth together continually. *Dull ache* of the upper and lower teeth, with swelling of the gums, (neither throbbing, nor stinging, nor drawing, (a. 15 d.) The teeth only ache when chewing.

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

The teeth are excessively painful when touching them, or when chewing, as if there were sub-cutaneous ulceration. Toothache only at night; when it ceased early in the morning, she felt nervous and agitated, and was not able to fall asleep again. Toothache when touching the tooth ever so little, or when coughing. Spasmodic pain in the teeth. * Drawing, cramp-pain in the teeth, relieved by warm drinks. Drawing ache in the right lower molares. Tearing-drawing toothache in the left lower molares. Tearing in a hollow tooth. Stinging and drawing in teeth which cannot be distinctly indicated, alternately in the upper and lower row; this prevented her from falling asleep in the evening, (a. 9 d.) Single, violent stitches in a hollow tooth, slowly succeeding one another, and ceasing after she got warm in bed. Frequent stitches in an upper molar tooth of the right side. Stitches, gurgling and boring in a hollow tooth, (a. 12 h.) Boring pain in the crown of the tooth. Single jerks in the right upper molares. Burrowing toothache, with stitches, in an upper molar tooth; after a meal, Throbbing and clawing toothache. Throbbing toothache (after a meal.) Beating in a tooth, with *swelling of the gums.* Pain in a hollow tooth, as if it were being dashed to pieces, extending as far as the temples, (3. d.) Pain in a lower molar tooth, as if it were being dashed to pieces, very painful when eating, (7. d.) A sound tooth is painful when eating, as if it were too long. All the teeth ache, as if they were too dull. Looseness of some of the incisores. Great looseness of the teeth. The teeth become yellow. Heat and pain in the gums. Tearing in the gums, and in the region of the roots of the left lower incisores. Darting pain in the gums of the lower row, in the afternoon, (a. 10 d.) Prickling and stinging pains in the left side of the gums and cheek. *Swelling* of the gums over the anterior teeth, with swelling of the upper lip. Swelling of the gums hinders opening the mouth. Swelling between the upper gums and the zygoma, with some swelling of the cheeks and prickling pains. Fistula dentalis in an old socket, in which one of the roots is yet lodged, with swelling of the gums. Ulcer of the gums. The gums bleed violently when touched. Involuntary

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

clashing and grinding of the teeth. Small tumors in the mouth, in various places.

Mouth.—Numbness of the inner mouth and tongue. Involuntary movement of the tongue, causing the sound A to be changed to O. The tongue feels swollen. The tongue is painful and swollen in different places; this hindered talking. The tongue, at times, gets involuntarily between the upper lip and upper teeth, at times between the lower lip and lower teeth. The tongue darts involuntarily out of the mouth, and moves between the lips to and fro. Nasal twang. Relaxed condition of the mouth, and heaviness of the tongue. Soreness of the tongue. *A number of vesicles on the tip of the tongue, which feels painful, as if it were raw and burnt.* Tubercles on the tongue. Ulcer under the tongue, being very troublesome when talking or eating.

Throat.—Swelling and elongation of the uvula, (a. 6 d.) Pain in the throat during deglutition, frequently; she feels as if she swallowed too much at once, (a. 9 d.) *Sore throat.* Sore throat, when swallowing or coughing. Sore throat, as if the throat were swollen, only during empty deglutition. Inaction of the pharynx when swallowing, the food does not go down. * The pharynx feels contracted, nothing will go down. The throat feels too narrow, during deglutition; the food and drink comes out again at the nose. When eating the soup very hot, he is unable to swallow it. The throat feels sore and swollen; this is not perceived when talking or swallowing. Glandular swellings inside and outside of the throat, with stitching pain in the glands when swallowing, and in the ear. Sensation as if a ball were rising into the pharynx from below. Sensation in the throat, as if a stone were pressing into it from without and closed the throat, rather painful during deglutition, without hindering breathing. Tearing pains in the pharynx from below upwards. Tearing in the left side of the neck. Tearing on the left side of the pharynx and in the throat. Tingling tearing, with pressure, in the posterior and upper part of the palate. Stinging and dryness in the throat, (a. 5 d.) *Continual stinging in the throat.* Inflammation of the throat, with hoarseness and stitches, which make it difficult, either to swallow

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

solids or liquids, for nine days, (a. 12 d.) Inflammation of the whole fauces, with aching and stinging pain. Suppuration of the tonsils, with stinging pain when swallowing. * Ulcers of the tonsils, resembling chancre. Ulceration of the tonsils. Roughness of the pharynx, with sensation of swelling when swallowing. * *Dryness in the mouth and throat.* * *Great dryness in the mouth*, early in the morning, (a. 3 d.) Sore feeling of dryness in the pharynx, when swallowing, early in the morning. Feeling of dryness in the mouth, with much saliva. The saliva becomes dry on the palate and lips, and is converted into tough mucus. Feeling of dryness in the throat and mouth, without thirst, only in the evening, immediately after lying down, and during the night. Continual dryness in the throat. Troublesome dryness in the throat; she would like to drink, but is unable to swallow anything on account of the pain. In the morning the throat always feels very dry. Dryness of the throat, with much thirst. *Dry* and bitter mouth Mouth dry and sourish. Desire to throw up a sourish, watery, sometimes bloody fluid. Continual accumulation of water in the mouth, and spitting. Flow of saliva, tasting salt. The posterior part of the mouth is covered with tough mucus. Hawking up of bloody mucus, when taking his usual ride. Coated tongue. Scraping sensation in the mouth. Fœtid smell from the mouth, early in the morning, he perceives it himself.

Taste and Appetite.—* Bitter taste in the mouth, early in the morning. A bitter substance rises into her mouth, early in the morning, as if from acidity in the stomach. *Very bitter taste in the mouth*, at night; she is obliged to rise and to rinse her mouth. Continual bitter taste in the mouth, but not of the nourishment she takes. Bitter taste of all her food. Bitter, sour taste in the mouth, before and after breakfast. *Sour taste* in the mouth, especially early on waking up. Sour taste of everything she eats, even sweet things. Sour taste in the mouth while drinking cocoa. Mouldy taste in the mouth, from morning till noon. Cheesy taste in the mouth, (a 13 d.) Very sweetish taste in the mouth, (a. 48 h.) Water tastes as sweet as sugar, early in the morning. *Absence of thirst.* Continual thirst with dry lips and dry

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

mouth; but, when trying to swallow some liquid, it became repulsive to her and she was unable to get it down; at the same time she felt sick, faint and tired. Violent thirst, with saliva in the mouth, resembling a fine foam, (a. 1 h.) A good deal of thirst, she would like to drink all the time. Great desire to eat and eating in a hurry. Hunger immediately, after a meal, although stomach and abdomen felt replete and distended. * Excessive hunger; the more he eats, the more the stomach craves, and as long as he eats, he is well; but after eating, he constantly has a sourish taste on the tongue, and even the saliva appears sour; he does not perceive this however, when eating. * Canine hunger, at noon, when eating, with sensation as if she could not satiate herself. Constant canine hunger, in the afternoon, with sensation as if heavy substances were lying in the stomach. If she does not eat while she is attacked with the canine hunger, she has headache, which goes off after a meal. Desire to eat, without any real hunger. * Want of appetite, (a. 3 d.) He does not relish what he eats, he has no appetite. Want of appetite, but much thirst, (a. 30 d.) *Aversion to solid food, especially meat, with thirst, (1. d.) She is unable to eat, is continually replete, has no appetite; and what she eats is repulsive to her, even to nausea. Aversion even to eating. Sometimes he loathes the best nourishment for moments, before having tasted it; but afterwards he cannot eat enough of it. Aversion to coffee and smoking. * Bread is repulsive to him, he is so much more fond of warm food. Sour taste, after taking milk. Bad taste in the mouth, shortly after dinner. After supper the forepart of the mouth feels watery, the backpart dry. Sour taste in the mouth after eating or drinking, with inclination to sour eructations. Acidity in the mouth after a meal. Oppressive bitter taste in the mouth after a meal.

Gastric Symptoms:—At dinner, excessive nausea unto fainting, sweat on the forehead and complete loss of appetite. Nausea in the pharynx and stomach after a meal, unto vomiting; with accumulation of water in the mouth. A good deal of thirst, after a meal. Hiccough after supper, for half an hour. Frequent eructations after a meal. Retching after a meal, rising of water, and discharge of water from the mouth, (water brash.) Sensation

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

as if the stomach had been spoiled. Digestion appears to be very slow. She dares not eat to satiety, because this makes her feel uneasy and bloated in the region of the liver. When eating to satiety, one feels uncomfortable and distended. *Immediately after a meal,* the abdomen *is full, bloated, distended,* until evening, when sitting, walking or lying down; when this symptom occurs, he is averse to walking and remains sitting. Distension of the abdomen, and tension of the head after dinner. Bloated abdomen, after a meal. Fulness and heaviness after a meal. After a meal, sensation in the stomach as of fasting, but without any hunger. Colic, after a meal. Pressure in the stomach, after every meal. Pinching in the abdomen after a meal. Pinching in the abdomen after breakfast, as if one had taken a cathartic. Pressure in the forehead, when eating. Continued pricking in the forehead, when eating; followed by violent single stitches during motion, (a. 36 h.) *Heat in the head after a meal, and a red spot upon the left cheek.* Black spots before the eyes after a meal, with pain, especially of the left eye, increased by shaking the head. Distortion of the features, especially after supper. High, burning redness over the whole face after dinner. Redness of the cheeks, followed by frightful pallor, after a meal. Urgent but almost unsuccessful desire to urinate after dinner. Tremor through the whole body, after a meal. Throbbing through the whole body, after a meal. After a meal, she feels more nervous and weary, her pulse beating faster, (a. 10 d.) Unconquerable sleep after a meal. Unconquerable sleep after dinner, followed by exhaustion. Shuddering during a meal, shaking the whole body, but without chilliness. Hot hands after a meal. After a meal, she feels as if the food touched a sore place; this sensation is followed by pressure in that part. *A number of eructations, alternating with yawning. Frequent empty eructations,* (the first days.) Eructations tasting of the ingesta, (1. d.) Continual bilious eructations, in the afternoon. * *Sour* eructations immediately, with colic. Sour eructations not leaving any taste in the mouth, but the acidity corroding the stomach. Rising of a sour fluid, with sour taste in the mouth. Sour eructations after every meal, with regurgitation of digested food, leaving a sort of

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

fetid taste in the mouth for one hour, with obtusion of the head, (a. 11 d.) Gulping up of the milk which had been taken early in the morning, with scraping, raw taste in the throat. Imperfect, burning eructations, reaching only the pharynx, where it causes a burning sensation for several hours, (a. 4 h.) Burning eructations, like a sort of heartburn. *Heartburn* from the stomach, acidity rising into the mouth. Heartburn, half an hour after every meal, with sour eructations and burning in the pit of the stomach for many hours; it almost takes away his breath, and makes him feel very weak. Heartburn after eating cold mutton, with pressure on the chest as if a stone were lying upon it, (a. 3:3 d.) Heartburn three hours after a meal, increased by smoking. *Hiccough.* Frequent hiccough, three days in succession, (a. 4 d.) Hiccough after every meal, (a. 19 d.) *Nausea every morning, before breakfast.* * Nausea in the afternoon, with sour eructations. Nausea, with oppression of the chest and pit of the stomach, with nausea in the afternoon, with some risings. Nausea, with oppression of the chest and pit of the stomach, and weakness in the lower limbs, relieved for a short while by eructations, returning afterwards, with tingling in the pharynx and pit of the stomach. Nausea rises into his head, it feels oppressed and obtuse, the pain extending to the nape of the neck; accompanied by trembling of the hands, better in the open air. *Nausea in a room*, going off in the open air; and then again nausea in the open air going off in a room. Nausea, with heat in the abdomen, and icy coldness in the face, (a. 2 d.) Loathing at the sight of food, with conflux of saliva, and flat, insipid taste in the mouth. * *Waterbrash* every other day, griping in the pit of the stomach, nausea, she is obliged to keep her mouth open, and to discharge a quantity of salt water which seems to come out of the stomach. Accumulation of water in the mouth, as takes place in canine hunger, in the forenoon, when writing, (a. 12 d.) Accumulation of water in the mouth with nausea, she had to spit a good deal, (the first 2 mornings.) A sort of bitter water rises every morning into her mouth from the stomach, she is obliged to bend over the edge of the bed and to spit it out. Qualms of sickness in the region of the stomach, early in the morning after rising, and especially when walk-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

ing out of the room, as if sweat would break out. Inclination to vomit, with hawking up of phlegm. * Nightly vomiting of food and bile preceded by nausea and anguish about the heart, (a. 9 d.) The child vomited mucus, five times after the siesta. Vomiting of coagulated blood, and acrid acidity.

Stomach.—Pain in the stomach, increased by stooping. * Intense pain of the pit of the stomach, when pressing upon it. Emptiness of the stomach previous to dinner; it obliged him to gape all the time. Violent pain in the stomach after a meal, and after catching a slight cold, with chilliness so that she cannot get warm, and deadness of the hands, (a. 23 d.) Violent pain across the stomach; she cannot bear lacing, nor tying any thing firmly over the part, (a. 8 d.) Cramp in the pit of the stomach, early in the morning when waking up, for three quarters of an hour, (a. 3 d.) Weight in the stomach, two hours after breakfast. * Pressure over the stomach, in the region of the orifice, in the evening. * Violent pressure in the stomach and abdomen, the whole forenoon, with pain to the touch and when breathing. Continued pressure in the stomach, with tension in the abdomen. *Pressure in the pit of the stomach,* (1. d.) Pressure in the pit of the stomach and the lower part of the chest, after lifting a heavy weight. Pressure in the pit of the stomach, especially in the afternoon and after lifting too heavy a weight, with pain in the pit when touching it. Aching from the pit of the stomach to the umbilicus, with guggling in the epigastrium. Spasm of the stomach previous to a meal, with unsuccessful eructations. Contusive pain in the stomach, going off by eructations; the stomach is also very painful to the touch: *Contraction and spasm of the stomach, as far as the chest,* from morning till evening. The spasm in the stomach is brought on again by wine. Whirling sensation in the pit of the stomach, with flushes of dry heat in the face. Tearing and drawing pain in the stomach, with nausea and colic, resembling the prick of a pin in the intestines. Gnawing and griping sensation in the region of the stomach; it feels replete. Stitching tension around the pit of the stomach, when breathing. Throbbing in the pit of the stomach, when raising the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

upper part of the body. Anxious feeling about the pit of the stomach, as one feels when swinging.

ABDOMEN.—Pain of the liver to the touch. Violent pain in the liver, with regular stools, (a. 8 d.) Pressure in the region of the liver. Aching in that region when taking an inspiration, (a. 13 d.) Pressure in the right side of the abdomen. Pressure in the region of the liver, from within outwards. Dull pressure in the region of the liver. Sharp pressure below the last right rib, during a deep inspiration and when bending to one side; also when pressing upon the right iliac region. Sore aching in the right hypochondrium, as if caused by a shock, increased by contact. *Tension in the lower part of the region of the liver*, with pressure. Grasping, as if with the hand, in the region of the liver. Violent cramp-pain of the diaphragm in the region of the liver, when stooping, and brought on by other slight causes, as if the liver were sprained. Pinching in the region of the liver, (a. 11 d.) Stitches in that region. Pinching stitch in that region. Stitches in the liver, in the evening, for an hour, (a. 6 d.) Painless twitchings in the surface of the liver, when coughing. Raw pain in the liver. Itching in the interior of the liver. Painful tension in the left hypochondrium. Colic, early in the morning, after rising. Pressure in the middle of the left side of the abdomen. Pressure in the abdomen, near the hips, at times on the right, at times on the left side. Oppressive colic, early in the morning, (5. d.) * Itching in the right iliac region, the whole day; he was obliged to walk crooked, from pain, and had short breath, (a. 6 d.) * Aching in the epigastrium, as if caused by flatulence, increased when drawing the abdomen in, relieved by eructations. Pressure in the abdomen, with drawing pain. * Pressure and cutting in the abdomen, before dinner. Sharp pressure at a small spot in the middle of the epigastrium. * Frequent pinching, pressing from within outwards, on the right side of the umbilicus, towards the hip, and somewhat below it. Weight in the abdomen. Weight in the left side of the abdomen, independent of breathing; he feels it in the same degree when walking, sitting, or lying, (a. 24 h.) Fulness in the abdomen, with pressing towards the rectum. * *Full, distended abdomen*, and cold feet, (a. 6 d.) Big abdomen, and colic

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

every day, (a. 2 d.) Distension of the abdomen, especially immediately before the menses. Distension of the abdomen from flatulence, (a. 4 d.) Distension of the abdomen towards evening, with incarceration of flatulence. Distension of the abdomen for several afternoons, beginning at 4 o'clock. *Tension in the hypogastrium*, (a. 6 h.) Tension in the abdomen, with accumulation of much flatulence. Tension of the abdomen, with incarceration of flatulence. Tension and grunting in the abdomen. Distended abdomen, with desire for stool, mostly in the evening. Spasms in the abdomen, which is very much distended. Spasms in the abdomen. Spasmodic contraction in the abdomen. Intermittent oppressive clawing in the left iliac region. Griping and pinching around the umbilicus, beginning early in the morning, when in bed. Pinching in the abdomen, relieved by emission of flatulence, (a. 4 h.) Pinching in the abdomen, in the afternoon, (after a natural stool,) from 3 to 10 o'clock, with inclination to vomit. Cutting colic, previous to stool, (a. 17 d.) Cutting in the abdomen, about midnight, with vomiting and diarrhœa. Cutting in the hypogastrium, after dinner, followed by a stitch extending to the tip of the glans, twice in succession. Cutting in the epigastrium, every forenoon, and even early in the morning when in bed, without diarrhœa, until afternoon, going off by walking. Quickly passing cutting in the intestines, into the sides and hips, towards evening, (a. 11 d.) Intermittent tearing, with a sensation as of gurgling, in a small spot of the middle of the epigastrium, towards the left side. * *Drawing pain in the abdomen.* Drawing pain in the abdomen, with pressure. Drawing pain in the abdomen, down into the calves. Drawing colic. Clawing drawing, deep in the hypogastrium. Stitches, in the right iliac region, extending into the pelvis, at every breath, and every turning of the body; worse in the evening and at night, (a. 10 d.) Burning stitches, on the right side, near the umbilicus. Jerks in the abdomen, (a. 4 d.) Pulsations in the abdomen, with anxious feeling, as of cramp. *Intense sensitiveness to pain of the skin of the abdomen.* Stinging sore pain of the skin of the abdomen, even when merely touched by the clothes. Pain in the groins when walking, and pain in the back, (a. 6 d.) Pains in the region of the hernial sac. Pressure and frequent dull stitches

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

in the right inguinal region. Pressing from within outwards, in the right inguinal region. Pressing from within outwards, in the right inguinal region. Pressing from within outwards in the left inguinal region, followed by a gurgling sensation in the abdominal ring. Pulsative tearing pressing from within outwards in the right groin, near the thigh. Pulsations, deep in the right abdominal ring. Stitches in both groins, late in the evening, (2. d.) Painful, boring stitches on the left side, just above the groin, when walking or when at rest. * *Tearing stitches in the hernial region*, (a. 24 h.) Red swelling in the right groin; during motion, and when touching it, it is painful as from sub-cutaneous ulceration, (a. 16 d.) Small glandular swellings in the groins, (a. 21 d.) The hernial sac, in the groin, protrudes immediately after the menses, and is affected with a tearing pain. Accumulation of flatulence, in the evening, going off only in part, with pressure in the region of the umbilicus. He is tormented by flatulence, even early in the morning before breakfast. Incarceration of flatulence, after having been seated for two hours. *Incarceration of flatulence*, causing a decline of health, (a. 6 d.) A quantity of flatulence in the abdomen, the hypochondria, back, region of the ribs and chest, causing tension and gurgling, relieved by empty eructations. A chief symptom of lycopodium is to bring on flatulence. This flatulence seems to be the cause of a large portion of the symptoms with which the patient is affected. Violent colic, in the evening, as if from incarceration of flatulence, followed by rumbling in the abdomen, and discharge of flatus, (a. 10 d.) A quantity of flatulence moving about towards evening, causing some colic, with noiseless emission of inodorous flatulence, and distended abdomen. Grunting and tension of the abdomen. *Grunting and gurgling in the abdomen.* * *Audible and tangible gurgling in the left hypochondrium.* Violent rumbling in the abdomen, (a. 16 d.) Emission of flatulence, preceded by colic, (a. 4 d.) Desire for stool, followed by cramp-pain in the rectum, which prevents the expulsion of fæces. Desire for stool, extending only as far as the rectum, (in a few hours.) Troublesome stool, owing to contraction of the rectum. Painful closing of the anus.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

STOOL.—* Stool not daily, sluggish, and no desire. Stool every other day. No stool for the first two or three days; at the end of this period he has a copious, natural evacuation of the bowels. No desire for stool, early in the morning; he has a desire in the evening; however, he passes only little, and with great exertions, although the desire is very strong. * Desire for stool only in the evening, with distended abdomen. * Little stool, with sensation as if much remained behind, followed by excessive and painful accumulation of flatulence, (a. 24 h.) Urgent desire during stool, as if he would have a copious evacuation, but it amounts to very little. Inaction of the rectum at stool. He had to make great efforts in passing the stools; he had, however, a daily evacuation, although slight, and with burning at the rectum. For many days in succession the first part of the stools is knotty, the second soft, (a. 16 d.) Stool comes in small fragments. Thin stool mixed with hard pieces. Papescent stool, once or twice a day, from the fifth day, for several weeks. Soft stool, several times a day; he has to press it out with much force; no emission of flatulence. Tenesmus in the morning; diarrhœa in the afternoon. Diarrhœa, with colic, generally early in the morning, (at 2, 3 o'clock.) Pale stool. Fetid stool. Copious expulsion of thin mucus, but little stool, at noon. Blood with stool. *Hæmorrhage from the rectum*, even with soft stool, (a. 14 d.) Biting in the anus, with thin stool. Stinging in the rectum, with good stool. Fine stinging in the rectum, with knotty stool. Burning at the rectum during stool. *Burning at the anus during the frequent stools*, (a. 48 h.) Hard stool, with pain in the small of the back as if it would break, with colic as if the intestines would burst, (a. 40 d.) Pain in the upper part of the head and buzzing in the ears, during stool. Shock in the temples, during hard stool. Desire continues after a good operation, but it is unsuccessful. Burning itching of the rectum after soft stool. Burning at the rectum during stool which is not hard. Feeling of fulness in the rectum continues after a copious stool. Violent contractive pain in the perinæum for many hours, after scanty, hard stool. Stool is followed by abdominal and uterine spasms, transversely across the abdomen, mostly after soft stool.

Digitized by UNIVERSITY OF MICHIGAN
Original from UNIVERSITY OF MICHIGAN

Distension of the whole abdomen from flatulence, after stool. A good deal of rumbling in the abdomen, after stool. After stool, heat and pressing sensation in the head, and lassitude of the thighs. Great fatigue after stool. Distension of the varices of the rectum. Protrusion of the varices. When sitting, the varices of the rectum are painful. The rectum is frequently so narrow, that it protrudes during hard stool. Pressure in the rectum, at night, (after 24 d.) Aching, with pressure upon the rectum, with cramp-pain in the abdomen, which induced her to believe that her confinement would take place immediately, (it was to take place in 16 days.) Spasms, bearing down upon the rectum and the small of the back, like labour-pains. Clawing, cutting in the anus and perinæum early in the morning. Pinching and stinging in the margin of the anus. *Stitches in the rectum,* (a. 2 d.) Stitch in the rectum, from the small of the back. Stitches and sore pain in the rectum. Tearing in the rectum, arresting breath, (a. 40 h.) Itching of the rectum and anus, (even a. 28 d.) (or a. 12 d.) Itching of the anus and mons veneris. * Itching eruption of the anus, painful to the touch.

Urinary Organs. — Diminution of urine the first 8 days, so much more abundant in 14. During emission, the flow of urine is suddenly interrupted; only a few drops of turbid and slimy urine come out, with pains in the urethra, followed by aching in the groins, (a. 24 d.) Frequent, copious micturition. Frequent micturition, at night, (a. 9 d.) Frequent, foamy urine. Urine white-turbid, immediately after emission. Urine with yellow sediment, (a. 6 d.) Dark urine, with burning. Urine red-brown. Dark urine, with sediment, (a. 18 d.) Dark urine, with red sediment, (a. 32 d.) Red sand in the urine. Red sand in the urine, which remains pretty clear. Red-yellow sand in the urine. A little red sediment in the urine. Bright-red sediment in the urine. Strong smell of the urine, (the first days.) Hemorrhage from the urethra, without pain, (a. 6 d.) Burning in the female urethra during micturition. Smarting sensation during micturition. Smarting sensation in the female urethra, in the evening, during micturition. Clawing sensation in the perinæum, close to the anus, during micturition; it continues and sometimes recurs be-

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

tween the acts of micturition. Tingling burning in the urethra, in the evening when going to bed, after micturition. Violent, but short drawing pain in the forepart of the urethra. Jerking drawing in the posterior part of the urethra. Tearing in the orifice of the urethra, some time after urinating. Quickly-passing cutting in the forepart of the urethra. Sharp cutting from the posterior termination of the urethra obliquely into the abdomen. Violently cutting stitch obliquely through the penis close to the abdomen, at night, after emission of a quantity of flatulence. Stitches in the bladder. Stitches in the neck of the bladder and in the anus at the same time.

Genital Organs. — Quickly passing cutting in the genital organs, from the abdomen. Violently darting pain in the penis. Titillation through the genital organs. Titillating drawing in the tip of the glans. Stitches in the tip of the glans. Drawing and cutting in the glans. Tearing, with pressure, in the region of the corona glandis. A quantity of yellowish humour behind the corona glandis, with dark-red, soft elevations, smarting and itching for several days. Itching of the internal surface of the prepuce. Itching of the frænulum, under the prepuce. Itching of the scrotum. Prickling itching, especially of the scrotum. Stitches in the scrotum. Lancinating tearing in the side of the scrotum, in the evening when in bed. Darting sensation in the left testicle, (a. 29 d.) Great weakness in the genital organs and the neighbouring parts, with pain in the perinæum when sitting, (a. 3 d.) * *Diminution of the sexual instinct* for 10 days, (a. 7 d.) Extinction of the sexual desire, (secondary effect?) (a. 30 d.) Diminished sexual instinct; even voluptuous fancies do not bring on an erection, although the desire for an embrace is not wanting. Rare erections, (the first days.) * *The penis is small, cold, and remains relaxed*, (the first 14 d.) Excessive sexual desire, (a. 6 and 14 d.) Erections with relaxed scrotum, (a. 5 d.) Several erections during the day, (a. 7 weeks.) Pollution, (the first night.) Exhausting pollution, (2 d.) Discharge of prostatic juice, without any cause. Discharge of prostatic juice, without erection, with great lewdness. Even during an embrace the scrotum is relaxed, and the emission of semen is very slow,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

(a. 4 d.) He falls asleep during an embrace, without emission of semen, (a. 12 d.) An embrace is succeeded by weakness the whole next day, (a. 48 h.) Weakness with trembling, early in the morning, after a pollution. Tearing stitches in the organs of generation. Violent burning in the vagina, during and after an embrace. Drawing in the groin, as if the menses would come on, in an aged female. Bearing down in the abdomen, as if the menses would come on, 16 days after the former, (12. d.) Reappearance of the menses, which had ceased for two days past, (a. 16 h) Taking the drug three days after the cessation of the menses, blood flowed again in a fortnight. Menses two days too early, and too scanty, (a. 41 d.) *Menses four days too early*, (a. 12 d., also after 2 d.) Menses seven days too early, (4. d.) Brought back the menses, at the period of new moon, in a girl of 17 years, which she had lost for five months past; the usually accompanying pain did not return, (a. 16 d.) Delays the appearance of the menses by four days, (secondary effect ?) by three days; by five days. The menses continue until the sixth day; they generally lasted four days. Bloatedness of the abdomen previous to the appearance of the menses. Great heaviness of the lower limbs previous to the appearance of the menses. Cold feet previous to the appearance of the menses. Violent chilliness the day previous to the appearance of the menses, (a. 13 d.) *Uncomfortableness and chilliness previous to the appearance of the menses*, the whole day. Chilliness, about midnight, previous to the appearance of the menses; afterwards heat, especially in the face, with restlessness. Low-spirited, desponding and melancholy previous to the menses. Delirium, with weeping, as if she would become crazy, the day previous to the menses, and on the first day of the menses, (a. 7 d.) Dilatation of the pupils, some days previous and during the menses. Violent itching in the pudendum, which appeared to be swollen, during the menses, (a. 12 d.) During the menses, pain in the temples, as if they were being screwed towards one another, as if the forehead would burst. Dull headache, almost like tearing, during the menses. Acidity in the mouth, with coated tongue, during the menses. Nausea, during the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

menses. Violent pain in the back during the menses, early in the morning, when rising, she was unable to move. Swelling of the feet, during the menses. Sort of swoon during the menses, while standing in the church; she heard and saw nothing, with feeling of great heat in the interior, especially in the head, with great paleness of face; she had to sit down immediately, remained the whole evening in a sort of stupor, and complained of obtusion of the head the next day, (a. 3 d.) Stitches in the head, after the menses, recurring at short intervals. * Profuse leucorrhœa, with fits and starts, (a. 5 d.) Milky leucorrhœa. Discharge of blood-red leucorrhœa, several times, before full moon, (a. 7 d.)

Cold, Catarrh.—Sneezing, without coryza. Sneezing, every morning, for half an hour. Sneezing, fifteen times a day, without coryza, (a. 5 d.) She is unable to sneeze on account of a prickling pain in the throat. Violent tickling in the nose, without being able to sneeze. Obstruction of the nose, high up. Obstruction of the nose, towards morning. Obstruction of the nose, he can only breathe with his mouth open. Entire obstruction of the nose; the child's breath was frequently arrested for fifteen seconds during sleep, even when the mouth was open. *Dry coryza*, (a. 10 d.) Dry coryza; he cannot breathe at night. Dry coryza, with burning in the forehead and obtusion of the head, her eyes becoming entirely closed, with much thirst and nightly heat, which hindered sleeping. *Dryness of the nose*, and obstruction about the root of the nose. Dry feeling in the region of the posterior nares. Coryza, in a person who never suffered with it, (a. 21 d.) *Violent coryza*, with swelling of the nose. Violent coryza, with catarrhal headache, (a. 10 d.) Coryza with acrid discharge, making the upper lip sore, (a. 28 d.) Frequent coryza, with fetid discharge from the left nostril, ulcerating within. Fluent coryza in a few hours. Profuse fluent coryza, (a. 3 d.) Profuse fluent coryza, with pressure upon the chest. Renewed, profuse fluent coryza, (immediately.) Indurated mucus in the nose.

Throat.—Frequent aching in the larynx, when swallowing. A violent, titillating scraping in the trachea, just below the larynx, rouses him from a deep sleep, at two

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

o'clock at night. Dry feeling in the larynx. *Hoarseness*, (a. 25, 48 d.) Hoarseness, the chest becomes rough and sore from talking, especially in the afternoon. The chest feels oppressed. Sensation as if there were too much mucus in the chest; whizzing breathing, in day time, (a. 18 d.) * Rattling in the chest. Desire to clear the throat, with rough feeling in the throat, as if mucus were adhering to it, with titillation in the throat inducing cough. Irritation in the throat, inducing dry cough. Short and hacking cough, from titillation in the throat. Titillation in the larynx, followed by a few coughing fits, terminating in sneezing. Intolerably itching titillation in the larynx, forcing one to cough violently, (a. ¾ h.) Exhausting cough, in the evening before going to bed, as if the larynx were tickled with a feather, with little discharge, (a. 3 d.) *Titillation with cough and grey expectoration.* Titillation with cough, unto retching. Titillation with cough, as if caused by the fumes of sulphur in the throat, with grey, salt expectoration. * *Irritation with cough, caused by deep breathing*, extending the neck, and sometimes by empty deglutition. Short and hacking cough, with sore pain along the larynx. In the evening from four to eight, she has to cough, and to drink a good deal. In the evening, when in bed, he has to clear his throat, and suffers with a short and hacking cough. Night-cough and hoarseness; sore pain in the chest in those places where the expectoration is loosened. Nightly cough, affecting the stomach and the diaphragm, mostly previous to the rising of the sun. Nightly cough, almost without any intermission, bringing on a pain in the head, and in both sides of the abdomen. Nightly cough, with some expectoration, (a. 6 d.) * Dry cough, with wheezing, and sense as of snapping in the throat as of an electric spark. Dry, short cough, every morning, with sensation as if the throat were hoarse, although there is no hoarseness. Dry, wheezing cough, as in drunkards. Dry, rough cough, incommoding one especially at night. * The expectoration is salt. * Expulsion of salt mucus, early in the morning, evening and night. * Cough, *with grey, salt expectoration.* Cough, with expectoration of blackish mucus, day and night. Morning-cough, with green expectoration, after violent

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

pain in the chest. Expectoration of white mucus. Expectoration of pus, first thin, then thick, with troublesome cough with titillation. * Thick, white, yellowish expectoration, with violent cough. * Expectoration of *yellowish* pus, with raw and sore feeling in the chest, after long dry cough. Cough, with discharge of pus, for eight days, almost uninterrupted, with fever and violent night-sweats, as if in the last stage of tubercular phthisis. * Cough, with bloody expectoration. Hemorrhage, in a female affected with pulmonary phthisis, (a. 10 d.) The breathing becomes too short previous to the cough. Smarting, as from excoriation, in the chest, with yellow-grey expectoration. The breath is very short when coughing, not otherwise. * Concussion, as from a shock, in the temples, and at the same time in the chest. When coughing, she feels a violent beating in her head. Pain in the head, and in both sides of the abdomen, when coughing. A stitch-like jerk in the head, with a feeling of pressure, when coughing. Stitches in the throat, when coughing, not otherwise, nor during deglutition. She suffers with oppression of the chest, and stitches in the throat; these bring on a cough with sensation as of scraping, (5. d.) *The region of the stomach has become painful from the coughing.*

CHEST.—*The breathing is united with violent oppression of the chest. Arrest of breathing when going up stairs. The chest feels oppressed and full, when staying in the open air. Tightness of the chest, especially during motion, for several days, with aching in the pit of the stomach. * Oppression of the chest, (a. 24 h.) Asthma, as if the chest had been spasmodically constricted, (a. 8 d.) Asthma and shortness of breath, with congestion of blood to the chest, (a. 20 d.) A stitch in the chest, here and there, when breathing. Stitches in and below the chest, when breathing, for two hours, (after supper.) Stitches in the sternum, when taking a deep inspiration. When breathing, dartings and stitches in the left side. Sensation as if a quantity of air were ascending the trachea with a wave-like motion, and rushed out at the mouth. Pain in the chest, with cough. during a deep inspiration. Pain in the chest, the first 6 days, so violent, that it was impossible for him to lie on the left side; afterwards cough with green

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

morning-expectoration. From time to time he experiences painful dartings into the chest. Tension of the chest (in a few h.) Sensation of tension on the left side of the chest. Tension in the chest, especially the right side, when taking an inspiration. Violent tension and pressure in the right half of the chest. Tension and pressure on the chest, oppressing the breathing, alternating with distension of the abdomen, in the evening, (a. 4 d.) Pressure in the chest, (a. 10 d.) Pressure in the left side of the chest. Pressure on a small spot of the true ribs, below the left axilla. Pressure as from a button, on the right, true ribs. Feeling of pressure and sore pain in the chest. Dull pressure in the left side of the chest. Oppressive, rheumatic feeling in the chest, relieved by empty eructations. Pressure and anguish in the sub-præcordial region, passing over into a tendency to be sad; when extending the trunk after violent exercise. Pressure in the chest; it feels full and oppressed, (7. d.) Fulness in the chest (and stomach,) after a meal. Fulness in the chest, at noon, as if it were oppressed. Oppression of the chest, as if it were too full. *The chest feels oppressed and raw internally.* Heaviness on the chest. Slight anguish in the chest. Cutting pain in the right side of the chest. * *Stitches in the left side of the chest, also during an inspiration,* (a. 7 d.) Stitches in the left side of the chest, extending to the back, they almost hinder breathing. A number of stitches in the left side of the chest, from time to time. Tearing stitch in the lower part of the sternum, independent of breathing, when at rest. Pulsative stitches in the left side of the chest. Pulsative tearing below the right axilla. Pulsative tearing in the præcordial region. Tearing in the region of the left clavicle. Pain, as from a sprain, in the left side, with occasional jerks. Pulsative or gurgling sensation in the præcordial region, beside the beats of the heart. Violent palpitation of the heart, early in the morning, from 4 to 5 o'clock, (a. 48 h.) * Sudden, violent palpitation of the heart, after having become wearied, with yawning. Tremulous palpitation of the heart, (3. d.) *Itching on the chest,* (a. 3 and 7 d.) Stitches in the nipples. Swelling of one breast, which was painful when touched. Hard burning nodosity in the left mamma and under the arm.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Blood and viscid water ooze out of one niple, especially when touching it.

Back.—*Pain in the small of the back*, so violent that it contracted his chest, with pressure of the stomach and constriction of the abdomen, (in 3 d.) Pain in the small of the back, when lying upon the back, with great weakness, (3. d.) Pain in the small of the back, extending down to the feet. Violent pain in the small of the back; he dares not sit straight, (a. 5 d.) * Stiffness in the small of the back. Aching in the small of the back, (a. 4 d.) * Drawing pain in the small of the back, for 17 days. Tearing in the small of the back, in a transverse direction, when sitting straight. Stitches in the small of the back. Gurgling sensation, a little on the left side of the small of the back. Pain in the small of the back, as if the flesh were loose. Chilliness in the small of the back. Large swelling in the psoas-muscle, very painful when moving the body; (Silicea cured it.) Back and small of the back become stiff and inflexible from riding on horseback, walking and stooping; it gives him much trouble to raise himself. Stiffness, from the scapulæ towards the back. The spinal column had become curbed in a child of two years, for several weeks. At times the scapulæ behind are violently and involuntarily drawn to one another; at times the pectoral muscles in front. Pain in the back, extending to the shoulders and the small of the back. Pressure in the back, below the scapulæ (5. d.) Pressure in the region of the left kidney. Pressure in the region of the right kidney. Pressure in the back, above both hips. Tension, with pressure, in the left scapula, as if caused by a vesicatory. Rheumatic tension in the back, and the right side of the chest, more violent during an inspiration. Pinching in the back. Pinching and aching pains in the right side of the back. * *Drawing pain in the back*, for several hours, (4. d.) Drawing pain in the back, when sitting. Drawing in the back, between the scapulæ, (a. 11 d.) Drawing in and near the right scapula, in the evening, (a. 10 d.) Rheumatic pain in the left scapula, which made him unable to raise his arm to the head. Tearing near the dorsal spine, below the scapulæ. * Tearing on the right side of the dorsal spine. Tearing in the region of the right kidney.

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

Stitches in the back, towards the small of the back, when sitting. Stitches in the region of the left kidney. Stitching pain in the back, extending to the right scapula. Stitches between the scapulæ. Fine stitches in the middle of the back. Repeated stitches in the back, over the region of the right kidney. Stitches in the left side of the back, when taking an inspiration. Spasmodic stitches, in paroxysms, in the middle of the back, making motion impossible for a few minutes. Pain, as from a sprain, in the left side of the back, extending to the left hypochondrium. Guggling below the left scapula. Continual beating in the back. Chilliness in the back, for several days. Burning as of red-hot coal between the scapulæ. Burning in the skin, below the left shoulder. Pressure, afterwards burning in the upper region of the right scapula. Burning in the back. Itching of the back, (a. 3 d.) Violent itching of the back, towards the neck. Violent itching of the back in the evening, (a. 15 d.) Itching in the upper part of the back, with eruption, (a. 37 d.)

Neck.—* Large pimples between the scapulæ and in the nape of the neck, with a burning sensation. Pain in the nape of the neck, when bending the head backwards. When stooping, the nape of the neck feels too short. The posterior cervical muscles feel stretched. Stiffness of the neck, with gloominess of the head, (a. 5 d.) Stiff neck. Painful stiffness of the left side of the neck. Pressure with tension, posteriorly and on both sides of the neck. Drawing pain in the left cervical muscles. Drawing-clawing pinching along both sides of the neck from below upwards. Darting pain in the right cervical muscles from below upwards. Drawing pain in the external cervical muscles, extending to the shoulder and the elbow. Tearing through the right side of the neck, from the face, and down the arm, as far as the fingers. A sort of paralysis of the cervical muscles, the head fell forwards as if it would fall off, with sensation of vertigo for six hours, but without any disposition to lie down. Involuntary nodding of the head, alternately to the right and left side. Involuntary nodding of the head, first slow, then quicker. Involuntary stretching of the head now forwards, now backwards. Involuntary and alternate stretching and contraction of the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

cervical muscles. Swelling of the external and internal cervical muscles. Hard swelling of the glands on both sides of the neck. Stitches in the cervical glands, during deglutition, extending to the ears. The glands increase in size and hardness from the cold feet. * *Swelling of the cervical glands.* Throbbing and darting in the goitre, (in a few hours.) Large clusters of red pimples round the neck, with violent itching, (a. 28 d.)

SUPERIOR EXTREMITIES.—*Swelling of the axillary glands.* Large boil in the left axilla. Sharp pressure on a small spot of the left shoulder, posteriorly, close to the neck. Rheumatic tension in the right shoulder-joint. Tearing in the right shoulder, beginning at the neck, only in the evening, after lying down, and at night. * *Tearing in the shoulder and elbow-joints*, when at rest, not during motion. Violent tearing in the shoulder-joint, beginning at the neck, when in perfect rest, day or night; the pain may be relieved by lying upon the affected side; the pain is worse in day-time when that part of her becomes cold, and goes off by motion, even when merely knitting or sewing. Stitches in the shoulder and tearing in the arm, (a. 27 d.) Stitches in the shoulders and the left lower arm, (a. 8 d.) Paralytic pain in the shoulder-joint, which prevented him from lifting the arm high. Pain, as from bruises, of the right shoulder-joint, scapula, and upper arm. Involuntary starting upwards alternately of one or the other shoulder. * Painless jerks in the weak arm, at night. Spasmodic jerking of the arms. * Drawing in the left arm, apparently in the nerve. Drawing pain in the bones of the arm, as far as the fingers. The elbows of the child become bent; he dares not either to stretch or to touch them, on account of the pain. The arm, where the axillary glands are swollen, goes to sleep. Weakness of the arms when at work. The arms feel paralysed; he has to let them hang down; they feel strong when at work or during motion. Sudden paralysis of the right arm, in the evening, as if caused by apoplexy, (a. 5 d.) Twitchings in the left upper arm. Twitchings of the muscles of the upper arms. Drawing in the left upper arm. Tearing in the right upper arm. Itching of the upper arms, (5. d.) Tearing in the elbow-joint, only

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

during motion. Tearing in the region of the right olecranon process. Tearing in the left elbow, down to the wrist-joint. Tearing, with pressure, in and about the right elbow. Rheumatic drawing in the right lower arm, early in the morning. Tearing in the fore-arms, down to the hands, from washing. Tearing in the left lower arm near the bend of the elbow. Tearing in the ulnar nerve, down to the hand. Hot feeling in the lower part of the fore-arm. Large, inflamed swelling, like erysipelas, on the fore-arm below the elbow, passing into suppuration like a boil. Biting-itching pimples on the lower arms, filled with pus. Cramp in the hand, the whole day. Tearing in the right hand and the two middle-fingers, only at night when holding them below the feather-bed cover; the pain ceases when laying those parts upon the cover, (a. 13 d.) Tearing between the right wrist and the knuckle of the thumb. Tearing in the right hand, between thumb and index. Tearing in the hands, towards the fingers. Tearing in the outer side of the left hand and in the knuckle of the little finger, towards the wrist. Tearing in the right palm, below the middle fingers. Tearing in the palm of the right hand, with burning and itching in the skin, close below the fingers. Dull tearing in the wrist-joints. Stitches in the dorsum of the hand, (a. 21 d.) Violent dartings in the right hand. Involuntary shaking of the hands. Pain, as from a sprain, in the right wrist-joint. Continually cold hands. The hands go to sleep, early in the morning, when in bed. *The hands go to sleep after a long talk. Her hands are hot all the time; this is very disagreeable to her. Swelling and heat of the right hand, in the evening. Feeling of heat in the left hand, with anxiousness. Red swelling of the right hand, extending to the finger-joints, without pain, for several days. Sweaty palms of the hands. Great dryness of the skin of the hands. *Itching pimples on the hands*, (a. 7 d.) Small boils on the hands, with stinging pains when touching them. Warts are forming on the hands. At times the fingers set themselves apart one from the other involuntarily, at times they become involuntarily clenched. The middle finger becomes contracted and is drawn to one side, without any pain. Involuntary twitching of the fingers, while

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

asleep. Involuntary twitching of the left index-finger. The knuckles are painful when pressed upon, without either redness or swelling. Tearing in the joint of the thumb, he cannot bend it. Tearing in the left thumb. Tearing in the tip of the right thumb. Tearing in the ball of the left thumb. Tearing in the middle-fingers of the right hand. Tearing in the joint of the right middle-finger towards the tip. Tearing in the tip of the right middle-finger. Violent lancinating tearing in the tip and under the nail of the left middle-finger. Pain, as from a sprain in the posterior joint of the fourth finger, when clenching the hand. The two last fingers go to sleep, early in the morning when waking up. Deadness, coldness, and numbness of the two last fingers, early in the morning when waking up; nevertheless he is able to move them. Deadness of two fingers, early in the morning, for half an hour, with blue nails, (a. 31 d.) Hot feeling in the fingers which appear cold, externally. *Redness, inflammation and swelling of all the joints of the fingers.* Redness, inflammation and swelling of the middle joints of the fingers, with some swelling of the hands. Inflammation of a scratched spot on the finger. Inflammation and pain of the right-middle finger, owing to a small panaris. Itching of the fingers. Violent, almost painful itching of both the anterior joints of the right index-finger, such as is felt in an ulcerating wound, with some redness, and not going off by friction. Itching and stinging in some frozen fingers. Itching of the fingers which had been frozen formerly. Burning in hands and fingers, with redness of the fingers, as if they were frozen. Chilblain on the little finger, with redness and violent itching. An ulcer on the left index-finger, becoming larger, with violent pains which prevent him from sleeping in the evening. Pimple on the right thumb. Itching pimples between the fingers. Wart-shaped tubercles on the index-finger, going off soon.

INFERIOR EXTREMITIES.—Itching, with violent stitches on the right half of the nates. In the left nates, burning as in a wound. Tearing in the left nates, with slight pressure. Tearing in the upper part of the nates, under the right hip. Pressure on the hips, beginning in the small

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

of the back. Pain in the muscles surrounding the hip-joint, when pressing upon them, when sitting or lying down; does not hinder walking. Pressing in the region of the left hip. Tearing in the left hip-joint. Rheumatic tension in the left hip. Tension and tearing in the left hip. Paralytic pain in the hip-joint, posteriorly, when stooping and when rising from a chair after having been seated for a time. Pain, as from a sprain, in the hip, towards the small of the back, when rising, so that he had to limp, for two days. Boil on the nates. In the right limb he felt a pain every four days, from the hip-joint down to the foot, which obliged him to limp when walking. Drawing in the limbs from top to bottom, when at rest, better when taking exercise. The limbs go to sleep in day-time, when sitting down, (a. 6, 7 d.) Uneasiness in the thighs and legs, when lying down, (a. 9 d.) A good deal of uneasiness in the lower limbs, in the evening; he had to move them frequently. Great uneasiness in the lower limbs, in the evening, previous to going to bed, less so when in bed. Jerking, and jerking sort of trembling in the lower limbs. Involuntary, violent shaking, first of the right, afterwards of the left limb. Cold and heavy limbs. While walking, the lower limbs appeared to become insensible; (although they were warm;) the upper part of the body appeared to be in danger of falling over. The lower limbs felt tired and powerless, as if they had been bruised. Soreness between the limbs, high up; she is scarcely able to walk. The inner side of the left thigh feels sore, with some biting itching extending as far as the genital organs. Pain of the left side of the thigh, as if it had been wounded; burning afterwards. Continual jactitation of the muscles on the posterior surface of the right thigh. Cramp in the right thigh, extending to the knee; he is scarcely able to go up stairs. Involuntary spreading of the thighs, followed by pressing them together, with an erection afterwards. Tension in the femora and calves, mostly when sitting. Drawing pain in the posterior surface of the thigh. Drawing and burning in the thigh, (13. d.) Drawing and aching pain in the fore part of the left thigh. Tearing in the upper part of the left thigh. **Tearing, downwards, in the left thigh, mostly when sitting,**

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

especially when bending the knee. *Tearing in the middle of the right thigh.* Tearing in the thigh, in the evening, accompanied with a sensation as if the parts were excoriated, (9. d.) Pulsative tearing, with a paralytic feeling, in the outer muscles of the left thigh, when walking, (1. d.) Stitches in the left thigh, when setting the foot down. Pain, as from a bruise, in the right thigh, just above the knee-joint, increased by contact and motion. Pain in the left hip-joint, during motion, as if he had made a mis-step. Cold thrills along the left thigh, by day. After walking, the skin of the thighs is painful, as if it were excoriated; this pain brought on convulsions of the limb, for one hour. A large boil on the thigh, above the knee. Early in the morning, when rising from the bed, the knees feel painful, as if they would break; also during motion. * Stiffness in the knee-joint, early in the morning, when rising, as after a long journey on foot. The left knee is curbed, the child is unable to stretch it, from pain. Tension around the knees, as if the parts around were too short; she was unable to set her foot down. Fine dartings in the knee, several evenings. Drawing in the bend of the left knee, (a. 22 d.) Great uneasiness in both knees, at night, when lying in the bed, (a. 8 d.) * Tearing in the knees and ankles, with pain, even when touching them. Unusual lassitude in the knees. Sore pain in the knees and other parts of the limbs. Pain, as from a sprain, in the knee-joint. Swelling of the knees. Sweat of the swelling about the knees. * Itching in the bend of the right knee, (6. d.) Pain in the side of the tibia, as if it were in the bone, when touching the part, (a. 13 d.) Sensation in the leg, as if something were firmly tied around it. Cramp in the left calf, when sitting. * *Cramp in the calf*, at night, also in day-time, extorting cries from him; when sitting with his knees bent. Frequent attacks of a darting pain in the leg, below the knee. Drawing in the legs, at night. Drawing in the leg from the ankle to the knee, at 5 or 6 o'clock in the afternoon, for two hours Drawing in the right leg, in the evening, and occasional starting. Drawing and tearing in the left leg, (a. 90 d.) Tearing in the left tibia. Tearing in the left tibia, below the calf. Violent tearing, before midnight, extending from the knee,

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

through the calf, down to the feet; she had to sit up, and was unable to go to sleep. Tearing in the legs and toes. Rheumatic drawing in the left leg, at night, when waking up. Sharp, darting tearing, in the lower part of the left tibia, in the evening when in bed Violent tearing in the left leg, darting with a jerk. Lancinating tearing in the leg below the knee, which is at the same time felt in the thigh. Sensation as if the legs were swollen and heavy. Great heaviness of the legs, with uneasiness in the same. Swelling of the legs extending beyond the knees, with large red, hot patches, painful and burning, especially on the knee and ankle; she is unable to set her foot down, from pain and stitches; in the afternoon these symptoms are accompanied by shuddering and constipation. Red spots on the legs, resembling mosquito-bites, going and coming. Violent itching of the calves, down to the ankles. The ankles are painful, at night, (a. 10 d.) Pain in the ball of the right foot, when beginning to walk. Pain in the heel, when setting the foot down, as if a little stone were underneath it. Straining sensation around the ankles, (a. some d.) Burning tension in the dorsum of the foot, near the big toe. Drawing in the foot, below the ankle, with heat in the parts. Pressing in the foot, which had been affected formerly, (as if it would break open again, (a. 9 d.) Tearing under the left heel. Tearing near the heel, (6. d.) Tearing in the heels and in the ball, (12. d.) Pricking in the ball of the foot, when stepping upon it, and when pressing upon it. Stitches in the dorsum of the foot, (a. 20 d.) Prickings in both heels. Prickings in the feet, when walking in the open air. Violent cutting stitches in the left side of the heel. Pain, as from a sprain, in the right tarsal joint. Pain, as from a sprain in the outer malleolus, also when at rest. Sensation of stiffness in the left tarsal joint, (a. 4 d.) Pain as from sub-cutaneous ulceration, in the ball of the right foot. Pain as from sub-cutaneous ulceration, in the soles of the feet, when setting the foot down, or when sitting, with a burning sensation Burning in the feet, (a 28 d.) Nightly burning in the soles of the feet. Great weight of the feet, (a. 6 d.) Swelling around the ankles, (a 6 d.) *Swelling of the feet*, also during the menses *Considerable swelling of the right foot*, (the first days.) Swelling of the

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

left foot, with stitches in the toes, when stepping upon it. Swelling of the feet, with stitches in the ankles, mostly when waking. Swelling of the dorsa of the feet, (the first days.) The swelling of the feet increases into ascites, with swelling of the genital organs, oppressed breathing, and scanty micturition with pressing, (a. 10 d.) Both feet are numb and have gone to sleep, as far as the calves, at night. The left heel feels as if it had gone to sleep. His feet feel cold easily. Cold feet, constantly. Coldness, first, of the right, afterwards also, of the left foot, in the evening, when in bed, for one hour. Coldness of the right foot, and heat of the left, (a. 2 d.) * *Cold, sweaty feet.* Sweaty feet. * *Profuse sweat on the feet*, until they become sore. Itching around the tarsal joint. Tumours on the borders of the foot, painful when walking. Callosities in the heel, with soreness. Rhagades in the heel. The toes are involuntarily extended and contracted again. Pressure on the ball of the big toe. Tearing in the first three toes of the right foot. Tearing and drawing in a corn on the little toe, painful when touched. Stitches in the big toe of the right foot, in the evening. Stitches in the big toe, in the evening. Stitches in the big toe, and afterwards in the sole. Stitches, mostly early in the morning, in the little toe of the right foot, looking red and frozen. Sore pain in the ball of the big toe when walking; it feels as if it were excoriated. Sore pain between the toes, as from excoriation, (a. 28 d.) Burning sore feeling of the toes, as if sand were upon them. Burning-stinging sore feeling of the toes. Inflammatory pain around the nail of the big toe. Corns come on in a fortnight. Stitches in the corns, (a. 13 d.) Stitches in the corns, with sore feeling.

General Symptoms.—All the limbs are painful when touched. All the soft parts of the body are painful, when touched or pressed against. Every thing, on which she is sitting or lying, feels too hard for her. Pain, in various parts of the trunk, as if some muscles were spasmodically contracted, and then extended again. Intermitting, cramp-like drawing in the knees, fore-arms, hands and fingers. Drawing, with pressure, in all the joints, especially the knees. Drawing in the left hand and tarsus, early in the morning. Drawing and tension in the carpal and tarsal

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

joints, early in the morning, when in bed. Drawing in the limbs, every other afternoon, and also across the face. Drawing at times between the scapulæ, at times in the right limb, at times in the chest. Quickly passing tearing in various parts of the body. Pinching pains in different parts of the body. Violent stitches in the thoracic cavity and the umbilical region, arresting the breath, (10. d.) Stiffness of all the joints. Stiffness of the limbs and the small of the back; audible cracking in the joints, when bending them. Stiffness of the arms and lower limbs with insensibility and numbness; he is unable to walk without falling, or to eat alone, since he is no more able to use his hands. Stiffness of all the muscles of the upper part of the body, and the whole trunk; he is so stiff that he is unable to stir. During the paroxysm of the pains she is obliged to walk about and to weep; she is unable to remain quiet. Her ailments increase in the afternoon at 4 o'clock; but at 8 o'clock in the evening, she feels better, with the exception of her weakness. He feels better in the open air than in a room, where he is frequently unable to endure from heat and uneasiness. *Desire to go into the open air.* Aversion to stay in the room. She is extremely sensitive to the cold, open air. Feverish sensitiveness to cold air, especially after a meal. Sort of anguish and attack of giddiness, when walking in the open air. Severe obtusion of the head, after having been long in the open air. Heaviness of the lower limbs, when walking in the open air. Constrictive pressure in the middle of the chest, from walking in the open air. The oppression of the chest is very much increased by walking in the open air, and accompanied by loud palpitation of the heart. Heat in the eyes and palms of the hands after walking in the open air. When walking in the open air but a little, he perspires excessively and becomes quite weak. * Disposed to catch cold. * The skin of the whole body is hot and dry; hot hands. Early in the morning, itching of the head and back. * Itching, as if caused by fleas, of different places of the skin and of the herpetic eruptions. Prickling itching of different parts of the skin. Stitches in different parts of the body. Insufferable tingling stitches in the small of the back and in other places. Darting stitches from

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

the neck to the right foot. Smarting and burning itching over the whole body. Burning sensation in different parts of the skin, on the back, arms, etc. Violent itching of the lower limbs, the back, the nates, in the evening, when in bed, with blotches after the scratching, going off again very speedily. Pimples, partly itching, partly painful, on the occiput, small of the back and nates. Large, red spots on the legs, neither painful nor itching. Large, bright-red spots on the epigastrium, around the pit of the stomach, and on the joint of the thumb, with itching and burning. *Itching hepatic spots.* Small, herpetic, itching spots on both sides of the neck, and upon the back. * Violent itching of a herpetic eruption on the tibia. * Large boil on the left lower arm, making the whole arm stiff, and another boil on the left half of the nates, (in a few days.) * The painless ulcers bleed when bandaging them, and are then affected with painful stitches. Large boil appears on the left scapula, with inflammation all around, and burning stitches, with alternate chilliness and heat of the body. Appears to favor the ramollissement and curvature of the bones. Sensation in the bones, as if they contained no marrow. She feels a heaviness in all her limbs. * The whole body feels bruised, especially in the evening. He feels a heaviness in all his limbs, he is not disposed to work, he is out of humor, a good deal of heat rises to his face from time to time. * Drawing and stretching in all his limbs. Uncomfortableness in the whole body. He feels, early in the morning, as if he had spent a sleepless night. Desire to move about. Disagreeable sensation of uneasiness in the body, when sitting, which does not permit him to continue writing; he has to jump up and to breathe deeply; his chest feels oppressed. Great agitation of the blood, in the evening, increasing until it becomes a sensation of trembling. *Considerable orgasm of the blood*, towards evening. Orgasm of the blood; agitation in the whole circulatory apparatus. Disagreeable feeling of heat over the whole body, everything he wears is too heavy and hot; he has frequently to breathe deeply, he feels oppressed; his hair stands upon end, and feels as if it were drawn together into one cluster, (a. 24 h.) Internal restlessness, she feels as if she must

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN

kick about with hands and feet, accompanied by a feeling of faintish weakness, (during her headache.) Sensation as if the circulation of the blood were arrested. Frequent attacks of a very painful sensation. as if he became quite cold internally, as if the blood gradually ceased to be warm. Pain in the chest, with nausea, as if she would vomit, she then lost her voice, was only able to speak in a low tone; all this disappeared again after an eructation.

FITS, SPASMS.—Several daily attacks of half an hour each, first, of griping and grasping in the back, after which the side becomes affected with a sort of stitches; this is accompanied with obscuration of sight, and she has to lie down wherever she may be. After a vexation he becomes quite weak, as if he would fall over, with palpitation of the heart and trembling, the whole forenoon, (a. 14 d.) He has a vexation, after which the pit of the stomach becomes suddenly affected, and his limbs feel heavy as if they were made of lead. Involuntary jerkings in various places, which affected her a good deal Involuntary turning and twisting of the whole body; this makes him pant, and makes his face hot and red. Involuntary alternate extension and contraction of the muscles in different parts of the body, without pain, and with full consciousness, in paroxysms, one every seven days, for the period of eight weeks. Spasmodic contraction and extension of the limbs, almost without any pain. Epileptic fit; screaming, foam at the mouth, loss of consciousness, he kicks about with arms and limbs, he then imagined he would have to die, and complained of great anguish about the heart. (a. 39 d) Epileptic fit; his left arm was bent upwards, his fingers clenched for a couple of minutes; he then was without his senses, tore things, kicked about with arms and legs, screamed, had foam at the mouth for a quarter of an hour; he then lay like a dead being, motionless; then he began to mutter. Epileptic fit; visible twitching of the muscles of the whole of the right lower limb; the pit of the stomach became affected, he began to scream, without any consciousness, kicked about with arms and legs, had foam at the mouth; then he was motionless for half an hour; upon cold water being introduced into his mouth, he blew it out

Digitized by UNIVERSITY OF MICHIGAN Original from UNIVERSITY OF MICHIGAN