

Valorization of carrot, cabbage, and red beet by-products – A review

Marja Lehto¹, Hanna-Riitta Kymäläinen², Minna M. Kahala¹ and Eila P. Järvenpää¹

¹Natural Resources Institute Finland (Luke), Production systems, FI-31600 Jokioinen, Finland

²Department of Agricultural Sciences, University of Helsinki, P.O. Box 28 (Koetilantie 5), FI-00014 University of Helsinki, Finland
email: hanna-riitta.kymalainen@helsinki.fi

Approximately 30% of vegetables end up as by-products during processing. Vegetable by-products have traditionally been used as animal feed, soil improvers, or raw material for biogas. In this literature review, we examine and discuss studies concerning the utilization of cabbage, carrot, and red beet by-products in a wide variety of applications, including technical applications. These vegetables are widely cultivated in northern climates as well as around the world, and there is a need to outline potential uses for their by-products. Numerous product ideas and research results exist concerning the use of by-products in food, feed, pharmaceuticals and cosmetics, biosorbents, composites and films, energy products, soil amendments and pesticides, enzymes, dyes and biosurfactants, and electrical products. Food and food additives were the most examined targets of carrot, cabbage, and red beet by-product valorization. These studies could be used, for example, in creating a company's by-product utilization strategy. High-value biochemicals, tailored nutraceuticals, feed additives, bioplastics, or biofuels may be realistic options in large-scale production. Environmental, safety, and economic aspects must be taken into account when utilizing by-products.

Key words: side-stream, *Daucus carota* L., *Brassica oleracea* L. var. *capitata*, *Beta vulgaris* L. var. *conditiva*, utilization

Introduction

Roughly one-third of all food, approximately 1.3 billion tonnes, produced for human consumption is lost globally per year (FAO 2011). Both globally (FAO 2011, Sagar et al. 2018, FAO 2019) and in Europe (Caldeira et al. 2019), fruits and vegetables are the largest groups of food waste. Although the largest share of food is wasted at the consumption stage, the processing stage is considered a close second, and it can result in about 30% of vegetables going to waste (Sagar et al. 2018, EU 2020). At the primary production and processing stages of the vegetable value chain, vegetable loss is due to managerial, financial, and technical issues, including insufficient harvesting facilities, inadequate harvesting techniques, and a lack of information on climatic conditions and marketing systems (FAO 2011). For example, in Sweden, one third of the carrots (*Daucus carota* L.) produced are used for purposes other than food (Olsson 2023). Data on the exact waste quantities produced through food processing are very limited, and no official data are issued annually by the EU.

Waste is defined as any substance or object that the holder discards, or intends to discard, or is required to discard. A by-product, on the other hand, is a substance or object resulting from a production process whose primary aim is not the production of that item. The term “side stream” broadly refers to by-products or waste generated during industrial processes (EU Directive 2008/98/EC). Residue refers to a small amount of matter that remains after the main part has been used, removed, or has otherwise gone. By-products of vegetable processing include solid residues of peels, skins, seeds, stems, and pulp. Pomace is the by-product of carrot juicing, and mash or pulp is the by-product of the carrot peeling and cutting process (Duval 2020). For example, up to 40% of the outer leaves and core of cabbages (*Brassica oleracea* L. var. *capitata*) are considered by-products, and up to 30% of the total production is estimated not to be utilized as food (Zhang et al. 2022). Large amounts of waste are also generated during the trimming process (Mago et al. 2022). Vegetable waste thus involves a loss of valuable biomass and nutrients (Plazzotta et al. 2020, Marcillo-Parra et al. 2021).

Many original study articles exist concerning the by-products of carrots, cabbage, and red beets (*Beta vulgaris* L. var. *conditiva*), but their utilization as a whole has not been widely examined, although reviews concerning other vegetables are available. These three vegetables, particularly important in northern climates, are the focus of this review.

By-products of carrot, cabbage, and red beet are currently often used as such as animal feed (Nilnakara et al. 2009, Tumbas Šaponjac et al. 2016, Battistella Lasta et al. 2019, Kaur et al. 2022b), in landfills (Kaur et al. 2022b), as fertilizer (Nilnakara et al. 2009) or composted (Battistella Lasta et al. 2019). Their nutritional and biological

potential could offer benefits for health promotion (Galanakis 2012, Ben-Othman et al. 2020), endorsing business (Coman et al. 2020, Galanakis 2020), and decreasing negative environmental impacts (Lai et al. 2017, Majerska et al. 2019). The compounds in carrot, cabbage, and red beet by-products (Table 1) are the basis for several valorization options presented in this review. Utilization of the compounds in different valorization targets is discussed in the sections in question. Most valorization options require the use of processing technologies. These include, e.g., drying, blanching, fermentation, extraction, extrusion, encapsulation, thermochemical technologies, anaerobic digestion, and composting. In some cases, also the fractionation technique for separating the desired component from the bulk by-product, e.g., protein from cabbage leaves (Nynäs et al. 2021), needs development. Although important, technical feasibility, including pre-treatment, extraction and production methods, is outside the scope of this review.

Table 1. Compounds of carrot, cabbage and red beet as a basis for valorization of by-products of these vegetables

Vegetable	Beneficial compounds	Reference
Carrot (<i>Daucus carota</i> L.)	Carotenoids (carotene, lutein), dietary fibers, phytochemicals (polyphenols)	Sharma et al. 2012, Anal 2018, Amin et al. 2021, Sepúlveda et al. 2021
Cabbage (<i>Brassica oleracea</i> L. var. <i>capitata</i>)	Dietary fibers, protein, bioactive compounds, vitamins, minerals, sulphur-containing amino acids, antioxidants (ascorbic acid, phenolic compounds), carotenoids, glucosinolates, phytochemicals, tocopherols	Wennberg et al. 2003, Kim et al. 2004, Liang 2016, Singh et al. 2006, Jongaroontaprangsee et al. 2007, Kusznierevicz et al. 2008, Nilnakara et al. 2009, Singh et al. 2010, Samec et al. 2011, Gül et al. 2013, Šedlar et al. 2021
Red beet (<i>Beta vulgaris</i> L. var. <i>conditiva</i>)	Protein, fiber, lipids, phenolics, nitrates, flavonoids, carotenoids, betalains (betacyanin, beta-xanthins), vitamins, minerals, dietary fiber, polyphenols	Vulić et al. 2013, 2014, Porto Dalla Costa et al. 2017, Chauhan and Rajput 2018, Kohajdová et al. 2018, Kaur et al. 2021, Nutter et al. 2021, Petrovic et al. 2021, Sedlar et al. 2021, Zin and Bánvölgyi 2021, 2022

The aim of this literature review is to examine potential applications for by-products of carrot, cabbage, and red beet. Valorization of the by-products is discussed in terms of potential product sectors, as well as environmental and economic aspects.

Material and methods

For a narrative literature review (Green et al. 2006), publications about the valorization of the by-products of the vegetables in question were searched using the database of Helka libraries (UH 2025). For the valorization options, combinations of two search terms such as carrot and by-product were used. Carrot and cabbage were clear search terms, while for red beet, also the terms beetroot and red beetroot were used. As presented in the introduction, the by-product terminology used in literature is vague. Thus, for by-product, also alternative search terms byproduct, waste, reject, and residue were used.

Experimental studies were selected for further examination. The focus of this study is on by-products, which was a selection criterion. However, the experimental studies focusing on by-products included studies made with by-products from agriculture (e.g., leaves of carrots) or the processing industry (e.g., mash from juice processing), vegetable rejects, and normal vegetables as models for by-products. Since there was plenty of literature, the selection was made first by focusing on peer-reviewed articles and secondly by obtaining a versatile collection of valorization options, products, and utilization ideas. The second author conducted the main literature search and analysis of the articles, but all authors participated in the discussion and further processing of the data.

Valorization options for the by-products of carrot, cabbage and red beet

Valorization options for the focus vegetables were divided into nine product sectors. Food and nutraceutical applications, as the most examined sector, are presented first, followed by feed, pharmaceuticals, and cosmetics. Technical uses include biosorbents, composites and films, energy products, soil amendments and pesticides, enzymes, dyes and biosurfactants, and electrical applications.

Food ingredients and additives

Plenty of research with an interest in utilizing the nutritional components of the vegetables (Table 1) has been conducted in the food sector, including nutraceuticals. More studies on carrots and red beets than on cabbage were available for the food sector (Table 2). Most studies showed that the by-products of these vegetables had potential uses, including a variety of possible applications in food, for example, as a colorant, antioxidant, and dietary fiber. However, unwanted side effects on sensory properties, e.g., taste, are sometimes a challenge (Kohajdová et al. 2018, Petrovic et al. 2021, Abdo et al. 2022). In addition, studies on an industrial production scale are called for.

In the case of carrots, the most examined by-product was the pomace, or pulp, resulting from juice pressing (Table 2), both from fresh and fermented raw materials (Janiszewska-Turak et al. 2021). Carrot leaves and tops showed potential for producing food additives (Leite et al. 2011, de Oliveira et al. 2015), while the peels were suitable for extracting carotenoids (de Andrade Lima et al. 2018), and for preparing dietary fiber (Chantaro et al. 2008) and jam (Hussein et al. 2015).

Some studies showed potential for cabbage waste as a raw material for functional foods and beverages (Table 2). Rashad and Abdou (2001) found cabbage whey to be a suitable substrate for cultivating edible fungi. Proteins derived from, e.g., red beet and cabbage leaves had good functional properties, as noted in Sedlar et al. (2021). Red beet leaves have been used, e.g., as an emulsifier (Ralla et al. 2019) and as an ingredient for bakery products (Asadi and Khan 2021). Red beet peel could be used as a preservative agent due to its antioxidant properties (El-Beltagi et al. 2022), but these properties of a beetroot peel extract have also been questioned (Šeremet et al. 2020). Red beet pomace has been found to have antibacterial (Vulić et al. 2013, Salamatullah et al. 2021), antiradical and hepatoprotective (Vulić et al. 2014), and anticandidal properties (Salamatullah et al. 2021).

Table 2. Studies of carrot, cabbage, and red beet by-products in food products and nutraceuticals as food additives or functional food ingredients

By-products as raw materials	Main components or compounds utilized	Reference
Carrot leaves	E.g., antioxidants, chlorophyll, fatty acids	Leite et al. 2011
	Ingredient of flour: omega-3 fatty acids	de Oliveira et al. 2015
Carrot peel	Dietary fiber, pectin, β -carotene, phenolics, antioxidants	Chantaro et al. 2008
	Dietary fiber, β -carotene, anthocyanins	Amany et al. 2010
	Proteins, dietary fibers, carbohydrates, sensory properties	Baljeet et al. 2014
	Physical properties of extrudates	Alam et al. 2015
	β -carotene, phenolics, antioxidants, antioxidant dietary fiber powder	Hussein et al. 2015
	Carotenoids	de Andrade Lima et al. 2018
	Inulinase enzyme, high-fructose syrup and fructo-oligosaccharides	Singh et al. 2018
Carrot pulp and mash	Single-cell protein source for noodles or bread	Razzaq et al. 2022, Khan et al. 2022
	Carotenoids, sensory properties	Šeregelj et al. 2022
	Carotenoids, polyphenols, oligosaccharides,	Duval 2020
	E.g., carotenoids, polyphenols	Amin et al. 2021
Carrot discards	Phenolic compounds, carotenoids, tocopherols	Araújo-Rodrigues et al. 2021
	Gelling agent pectin	Christiaens et al. 2015
	Pigments, fermentation products: carotenoids, sugars, acids	Ramos-Andrés et al. 2021
Cabbage outer leaves	Colour, carotenes	Kaur et al. 2022a, b
	Substrate for the cultivation of edible fungi for human food and supplements	Rashad and Abdou 2001
	Dietary fiber powder	Jongaroontaprangsee et al. 2007
	Antioxidants, vitamin C, colour	Nilnakara et al. 2009
	Beverages, nutraceuticals: polyphenols, glucosinolates	Gonzales et al. 2015

Red cabbage leaves	Dietary fiber, minerals	Brito et al. 2020
	Antioxidants, acid binding, dialysis retardation	Liang et al. 2019
	Anthocyanins	Patras 2019
Red beet leaves	Polyphenols, antioxidants, food colourant	Zanoni et al. 2020
	E.g., proteins, dietary fiber, fats, carbohydrates, phenolics, antioxidants, colour	Asadi and Khan 2021
Red beet peel	Polyphenols, betaxanthin, betacyanin	Nutter et al. 2021
	Flavourant, colourant, antioxidant: betalains, phenolics	Kujala et al. 2001
	Cookies: e.g., protein, fiber, sensory parameters	Chauhan and Rajput 2018
	Emulsifier for food and beverage products	Ralla et al. 2019
	Phenolic compounds, e.g., betanin, and betacyanins	Kujala et al. 2000
	Fiber, protein, phenolics, antioxidants, betacyanin, biological effects	Šeremet et al. 2020
	Betalains, phenolics, antioxidants	Zin and Bánvölgyi 2021
	Antioxidant, preservatives phenolic compounds: flavonoids, betalains, antiradical scavenging	El-Beltagi et al. 2022
Red beet pomace	Betalain, phenolics, flavonoids	Zin and Bánvölgyi 2022
	Phytochemical profile, antiradical, antimicrobial, and cytotoxic activities	Vulić et al. 2013
	Antioxidant effects	Vulić et al. 2014
	Ingredient in biscuits: betacyanins, polyphenols, antioxidant capacity	Hidalgo et al. 2018
	Source of dietary fiber in bakery products: proteins, dietary fiber	Kohajdová et al. 2018
	Candy enrichment: phenolics, betacyanin, betaxanthin, antioxidant capacity	Kumar et al. 2018
	Ingredient in yogurt: protein, fiber	Jovanović et al. 2021
	Fortification for functional dairy dessert: betalains	Kaur et al. 2021
Red beet peels, parings, stalks	Liqueur: phenolics, antioxidants, colour	Petrović et al. 2021
	Colour, antioxidants, dietary fiber, phenolics, betalain	Porto Dalla Costa et al. 2017
Red beet leaves and stems	Betacyanin, polyphenols	Aguirre Calvo et al. 2019
	Fortification of functional orange juice: flavonoids, betalain, radicals scavenging activity	Abdo et al. 2022
Red beet pulp and peel	Polyphenols, flavonoids, antioxidant and antimicrobial activity	Salamatullah et al. 2021

Feed ingredients and additives

Similar to food uses, many components of vegetable by-products (Table 1) are beneficial in animal feed. For example, dried cabbage waste is an excellent source of protein (Mahgoub et al. 2018). Animal feed is a basic utilization target for carrot (Aimaretti and Yibalo 2012, Bakshi et al. 2016) and cabbage (Nilnakara et al. 2009) discards, but the need for other valorization alternatives has been raised (Aimaretti and Yibalo 2012). As an example of a new type of utilization, Rashad and Abdou (2001) found cabbage whey to be a suitable substrate for cultivating edible fungi, also for feed use. Castrica et al. (2018) estimated pet food to be the most concrete strategy for using food waste as animal feed within the European context, but most of the studies focus on animals other than pets (Table 3).

Table 3. Studies of carrot, cabbage, and red beet by-products in feed and pet food ingredients and additives

By-products as raw materials	Product or application examples	Reference
Carrot pomace	Insect feed, e.g., proteins and carotenoids	Rovai et al. 2022
Carrot extract	Fermented for feed and food additives	Stockhammer et al. 2009
Carrot discards	Supplement for prawn feed vinasse	Garces and Heinen 1993
	Forage for egg-laying hens	Hammershøj et al. 2010
	Co-fermented carrot and brewer's yeast, animal feed	Aimaretti et al. 2012
Cabbage outer leaves	Fermented cabbage for calf feed	Mukodiningasih et al. 2019a, b
	Ruminant (sheep) feed, goat feed	Ngu and Ledin 2005, Wadhwa et al. 2006, de Evan et al. 2019
	Fermented extract as a starter for rice bran fermentation	Setya Utama et al. 2013
	Poultry feed	Mustafa and Baurhoo 2018
	Rabbit feed	Nguen et al. 2009
	Snail feed, e.g., proteins and fibers	Omolara and Olaleye 2010, Babalola 2018
	Livestock feed	Mahgoub et al. 2018
Red beet discards	House cricket feed, e.g., proteins and carbohydrates	Morales-Ramos et al. 2020
	Substrate for cultivation of fungi for feed	Rashad and Abdou 2001
	Fermented red beets for lamb feed	Nkosi and Ratsaka 2010
Red beet pomace	Dog food	Jovanović et al. 2021

Ruminants have the capacity to utilize fibrous materials because of their rumen microbiota, which enables the use of many by-products in their feeds (Mirzaei-Aghsaghali and Maheri-Sis 2008), either fresh (whole or chopped), as dried and ground meal, or as silage (de Rezende et al. 2015, Bakshi et al. 2016, Du et al. 2021, Ren et al. 2021). Rust and Buskirk (2008) estimated 40% (DB, dry basis) carrot being suitable for beef cattle, and carrot was also mentioned as being suitable for cows (15.9 kg/cow/day), while a carrot addition of 5 kg/cow/day was reported to have beneficial effects on milk quality (Nalecz–Tarwacka et al. 2003). As a disadvantage for ruminant feeds, carrot and red beet pomaces increased the gas formation related to digestible organic matter (Giller et al. 2021). Cabbage waste leaves are potential in goat feeds (Ngu and Ledin 2005, Wadhwa et al. 2006), but sulfur toxicity in cabbage wastes and its effect on the feed flavor should be taken into account (De Evan et al. 2019). Ensiling and fermentation of vegetable by-products may enhance the acceptance, performance, and growth of animals, such as in carrot feed for steers (Laflamme 1992) and outer leaves of cabbage for calves (Mukodiningasih et al. 2019a).

Concerning poultry feed, carrot by-products had positive effects on the quality of eggs (Sidker et al. 1998, Hammershøj et al. 2010) and hen welfare (Steenfeldt et al. 2007). Cabbage leaf residues could improve nutrient utilization and egg quality (Mustafa and Baurhoo 2018). Cabbage has also been used in rabbit feed (Tsutsumi et al. 1967), with beneficial effects on vitamin A (Pirie and Wood 1946), but the effect of cabbage on growth rate varied from positive (Nguen et al. 2009) to not beneficial (Hang et al. 2011).

Carrot feed as a source of carotenoids improved the colour and sensory properties of catla, a carp fish (Weerakkody and Cumaranatunga 2016), the color of cichlid (Kop et al. 2010), and the colour (Lili et al. 2018) and growth (Wagde et al. 2018) of the swordtail, an ornamental aquarium fish, while gilthead seabream colour was not improved by carrot feed (Wassef et al. 2010). Carrot and carrot tops increased prawn pigmentation, although this was not considered an important feature (Garces and Heinen 1993).

Insects can utilize vegetable by-products as feed and convert the organic waste material of, e.g., carrots (Oonincx et al. 2015, Varelak 2019, Aristi et al. 2020, Rovai et al. 2022) or cabbage (Aristi et al. 2020, Morales-Ramos et al. 2020, Pinotti and Ottoboni 2021) into biomass that can be used as a raw material for human food and/or animal feed or technical applications (van Peer et al. 2021). A diet enriched with, e.g., shredded carrots or cabbages improved the growth of mealworm larvae that were considered as human food ingredients (Liu et al. 2020). The initial interest being in human food (Oonincx and Boer 2012, Van Broekhoven et al. 2015, Rovai et al. 2022) or food and feed (Oonincx et al. 2015), carrot was used as a basic feed substance for mealworms, e.g., to provide moisture. Carrot waste was promising for the fat content of black soldier fly larvae (Aristi et al. 2020). Black soldier

flies (Ooninx et al. 2015, Aristi et al. 2020) and cockroaches (Ooninx et al. 2015) fed by vegetables were intended to be feed for other animals. Syahrizal et al. (2022) observed that a mixture of palm kernel meal and cabbage waste was among the most favorable feed for black soldier flies that could be used for fish feed. Omolara and Olaleye (2010) and Babalola (2018) observed that snails can utilize cabbage waste as their sole feed. Carrots are also used as feed for insects grown for pest management, such as flies (Mainali et al. 2019).

Pharmaceuticals and cosmetics

Case studies of carrot, cabbage, or red beet by-products as pharmaceuticals were often marginal, and the potential for food, nutraceutical, and pharmaceutical uses was often combined (Table 4). Examples of promising results include red beet stalks and leaves by-products in obesity control (Micheletti Lorizola et al. 2021) and red beet pomace having cytotoxic effects against carcinoma cells (Vulic et al. 2013). Vegetable pigments could be used in pharmaceutical products (de Andrade Lima et al. 2018, Ramos-Andrés et al. 2021) and pigments and odorous substances in creams and perfumes (Naviglio et al. 2019). Components of vegetable by-products shown to have potential as health products include β -carotene (Roohinejad et al. 2014, Purohit and Gogate 2015), insoluble fiber (Ma et al. 2016), and phenolic compounds (El-Sawi et al. 2022) in carrot waste, organic acids (Selder et al. 2021), phytochemicals (Šamec et al. 2011), and phenolics (Gonzales et al. 2015) in cabbage waste, and betalains and polyphenolic compounds in red beet waste (Lazăr et al. 2021).

Table 4. Studies of carrot, cabbage, and red beet by-products in pharmaceutical products and cosmetics

By-products as raw materials	Product or application examples: main components or compounds utilized	Reference
Carrot peel	Pharmaceuticals or cosmetics: carotenoids	de Andrade Lima et al. 2018
	Pharmaceutical applications: carotenoids	El-Sawi et al. 2022
Carrot pomace	Pharmaceuticals, functional ingredient for intestinal health	Roohinejad et al. 2014
	Pharmaceutical against cancer and microbial infections: phenolics, flavonoids	Purohit and Gogate 2015
	Dietary fiber	Ma et al. 2016
Carrot discards	Pigments for pharmaceutical industry: sugars, acids	Ramos-Andrés et al. 2021
Cabbage outer leaves	Additives for cosmetics (red cabbage): polyphenols, glucosinolates	Gonzales et al. 2015
	Pharmaceutics, food, and feed: butyric, valeric, and caproic acids	Selder et al. 2021
	Antibacterial and antifungal lipophilic mixture for the clinical, veterinary, and agricultural fields	Arrais et al. 2022
Red beet peel	Pharmaceuticals and nutraceuticals: betalains, polyphenols	Lazăr et al. 2021
Red beet pomace	Pharmaceutical and cosmetic industry: phytochemical profile, antiradical, antimicrobial and cytotoxic activities	Vulic et al. 2013
	Encapsulated extract for pharmaceuticals, food additives: phenolics, flavonoids, betalain	Tumbas Šaponjac et al. 2016
	Medicinal and food applications: betacyanin, beta-xanthin, phenolics, antioxidants	Kuswaha et al. 2018
Red beet leaves and stems	Nutraceuticals, functional food, or pharmaceuticals: phenolics	Battistella Lasta et al. 2019
	Supplementation, adjuvant in obesity	Micheletti Lorizola et al. 2021

Biosorbents

Activated carbon was the most common active component from carrot or cabbage by-products in biosorbent studies (Table 5), and it was examined mostly for wastewater purification and, in a few studies, for purifying soil and vapor. Lignocellulose-based biosorbents were also common for wastewater purification. Single studies concerning carrot by-products in biosorbents for wastewater were based on pectin (Hastuti et al. 2018) and peroxidase enzyme (Joel et al. 2020), while one study focused on the bioremediation of polluted soil

(Hamoudi-Belarbi et al. 2018). Regardless of the active component or structure examined, the by-products examined in most of the studies showed potential as biosorbent materials, with cabbage being the only exception in Hossain et al. (2014) and Xue et al. (2019).

Table 5. Studies of carrot, cabbage, and red beet by-products in biosorbents

By-products as raw materials	Product or application examples (potential raw material, active component)	Reference
Carrot discards	Biochar/active carbon for removing phosphorus from wastewater	de Carvalho Eufrásio Pinto et al. 2019
	Biochar/active carbon for dye removal from wastewater	Hira et al. 2020, Moradi et al. 2021
Carrot residue or discards	Lignocellulose as biosorbent/adsorbent for heavy metal removal	Nasernejad et al. 2005, Bhatti et al. 2010
Carrot peel and pulp	Composite PET (Polyethylene terephthalate) adsorbent containing tomato and carrot for removing cobalt from wastewater	Changmai et al. 2018
Carrot peel	Biostimulation medium for bioremediation of crude petroleum	Hamoudi-Belarbi et al. 2018
	Pectin as biosorbent for heavy metals from wastewater	Hastuti et al. 2018
Carrot pulp	Biochar for heavy metal fractionation in contaminated soil	Gholami and Rahimi 2021
Carrot leaves and stems	Lignocellulose as biosorbent for removing dyes from wastewater	Kuswaha et al. 2014
Cabbage outer leaves	Lignocellulose as biosorbent/adsorbent for removing heavy metals or dyes from wastewater	Hossain et al. 2014, Wekoye et al. 2020
	Peroxidase enzyme for biodegradation of phenol and synthetic dyes from wastewater	Joel et al. 2020
	Activated carbon as adsorbent for mercury vapour removal	Vakili et al. 2021a, b
	Biochar for purifying off-gas or removing VOCs	Zhang et al. 2022
Red beet fiber	Lignocellulose as biofilter/adsorbent for removing dyes and heavy metals from wastewater, hard water softener	Rima et al. 2014

Composites, films, and packages

Using carrot and red beet discards or waste in composites and packages has been examined in some studies (Table 6), all of which showed that carrot or red beet by-products have at least preliminary potential in the production of films and packaging materials. The most typical end products examined or mentioned were food packages (Tran et al. 2017, Otoni et al. 2018, Perotto et al. 2018, Sogut and Cakmak 2020, Amoroso et al. 2022, Rodríguez-Félix et al. 2022), while the end product was not clearly specified in the remaining studies (Table 6).

Table 6. Valorization studies of carrot and red beet by-products in composites, films, and packages

By-products as raw materials	Product or application examples	Reference
Carrot pomace	Cellulose nanofibers for, e.g., reinforcement in polymer composites	Berglund et al. 2016
	Cellulose and nanocellulose film	Szymanska-Chargot et al. 2019
Carrot peel	Cellulose nanofibers for biocomposites for packaging	Otoni et al. 2018
Carrot discards and pulp	Cellulose nanofibers for bioplastics	Perotto et al. 2018
Carrot discards	Cellulose nanofiber film for packaging	Amoroso et al. 2022
	Dietary fibers, syrup to produce lactic acid for biodegradable plastic	Salvañal et al. 2021
Carrot pulp	Fibers and microcrystalline cellulose, packaging films	Sogut and Cakmak 2020
Red beet discards, peel	Biocomposite for packaging for pharmaceutical, medical, and food applications	Tran et al. 2017
Red beet pomace	Active food packaging	Rodríguez-Félix et al. 2022

Energy products

Energy products made from vegetable by-products include biogas, ethanol, and biodiesel. Cabbage (Solowski et al. 2019, Wei et al. 2021, Czubaszek et al. 2022), red beet (Surendran and Shanmugam 2021), and mixtures of vegetable wastes containing carrot (Yang and Cosolini 2019) and carrot and cabbage (Ravi et al. 2018) showed potential for biogas production. In some studies, co-fermentation of a mixture of cabbage waste and other raw materials produced more biogas than the wastes separately (Bozym et al. 2015, Wu et al. 2016, Mu et al. 2017), but opposite results have also been obtained (Arifan et al. 2021). Ethanol was produced from carrot discards (Aimaretti and Yibalo 2012, Aimaretti et al. 2012) or pomace (Yu et al. 2013, Khoshkho et al. 2022) through fermentation, while the residue could be used as animal feed. Biodiesel was prepared from carrot pomace by transesterification (Karatay et al. 2019, 2020), carrot being the growth medium or carbon source for microbial lipid production.

Soil amendments and pesticides

Improving soil properties or fertilization are the most common applications of vegetable by-products for agricultural purposes, while only a few studies have focused on pesticide use. Processed vegetable residues are used in agriculture mainly in the form of biochar, as residues from anaerobic digestion in energy production, or as compost. Cabbage was the most examined vegetable for this application sector (Table 7).

In agriculture, biochar can be used in various ways. As a soil ameliorant, it is used to optimize soil structure and composition, and to increase the availability of nutrients and the water retention capacity of the soil. Biochar buried in the soil, either as such or as a compost additive, serves as a long-term carbon storage, increases soil health by limiting the availability of pesticides and heavy metals, and contributes to the improvement of plant growth and crop production by affecting soil microbiology and enzyme activity (Enaime and Lübken 2021). Biochar as a soil amendment was discussed by Bakshi et al. (2021). Biochar can also be used as animal feed (Enaime and Lübken 2021). Most of the studies on the removal of contaminants from water or more widely from the environment (Table 8) focus on heavy metals. However, phosphorus, in particular, is relevant for agriculture. Carrot or cabbage by-products have shown potential as raw materials of biochar (de Carvalho Eufrásio Pinto et al. 2019, Pradhan et al. 2020a, b).

Anaerobic digestion produces biogas and digestate as a co-product, which can be used as fertilizer. Composting as a treatment of cabbage for soil amendment applications showed promising results in most studies (Table 7). Studies focusing on other treatments or active components for soil improvement were few: polymer gel from cabbage waste (Zhang et al. 2021) and lime mud from red beet juice production (Ławińska et al. 2020) showed potential, while carrot acid extracts did not (Suzuki et al. 2001) (Table 7). Results from studies examining the utilization of cabbage waste for biopesticides varied from positive (Zhang et al. 2022) to unpromising (Roshan-Bakhsh et al. 2019, Supyani et al. 2021).

Table 7. Studies of by-product use in soil amendments and pesticides

By-products as raw materials	Product or application examples	Reference
Cabbage outer leaves	Co-composting with phosphate rock	Walker et al. 2012
	Co-composting with poultry manure	Saleem et al. 2017, 2018
	Composting: carrier of fungi, control of plant pathogens in soil	Wolna-Maruwka et al. 2019
	Composting with fungus	Ntsohi et al. 2021
	Vermicomposting: soil amendment	Mago et al. 2022
Red cabbage outer leaves	Biochar for use in P recovery and as fertilizer	de Carvalho Eufrásio Pinto et al. 2019
Cabbage outer leaves	Extracts by various solvents, natural nematicide	Roshan-Bakhsh et al. 2019
	Biochar for soil amendment, balancing soil properties	Pradhan et al. 2020a&b
	Fermentation extract with cow urine to produce biopesticide	Supyani et al. 2021
	Applications of superabsorbent polymer gel, e.g. soil amendment	Zhang et al. 2021
	Extract before pyrolysis for use as pesticides or insecticides, biochar, activated carbon, pyrolytic liquid, pyrolytic gas	Zhang et al. 2022
Carrot pomace	Acid extract of carrot pomace as a plant growth substance	Suzuki et al. 2001
Red beet pomace	Carbonation lime mud, raw material for producing mineral–organic fertilizers	Ławińska et al. 2020

Enzymes, dyes, and biosurfactants

Some studies showed good potential for using carrot or cabbage waste in enzyme production or for using red color extract from cabbage or red beet as a textile dye (Table 8). Some results were contradictory: the study by George and Jayachandran (2008) did not favor carrot peels in biosurfactant production, while positive results were obtained by Andrade et al. (2014) for red beet waste from juice production for the same purpose. Red beet skin was the most promising for producing biocatalysts to produce pure compounds for several fields of application (Vandenberghé et al. 2013).

Table 8. Studies of carrot, cabbage, and red beet by-products in enzyme, dye, and biosurfactant production

By-products as raw materials	Product or application examples	Reference
Vegetable by-products, e.g., carrot peel	Growing media for extremophile biomass production, enzymes and polymers	Di Donato et al. 2011
Carrot peel	Biosurfactants	George and Jayachandran 2008
Carrot pulp	Fatty acids precursor, fibrous booster to maintain in wastewater treatment	Hastuti et al. 2018
Cabbage outer leaves	Enzymes for dry-cleaning, detergents, meat processing, cheese making, silver recovery, digestive and medical treatments, waste management	Madhumithah et al. 2011
	Enzymes for textile and paper industries	Das et al. 2012
	Enzyme production with slaughterhouse effluent and cabbage waste mixture	Ramakodi et al. 2020
Carrot, red cabbage, red beet by-products	Biocatalyst for pharmaceutically important molecules, agrochemicals, flavours, and asymmetric chiral ligands	Vandenberghé et al. 2013
Red cabbage outer leaves	Colourant for textile industry	Priyadarshini et al. 2021
Red beet peel	Betalains for colour industry	Fernando et al. 2021, Popescu et al. 2021

Electrical uses

Electrical products are a marginally examined product sector for the vegetable by-products. However, Ahmed et al. (2018) and Hoang et al. (2019) found carrot waste to be suitable for energy storage products (Table 9). Of the materials examined, Mannarmannan and Biswas (2021) found carrot peel extract to be the best for preparing copper oxide nanoparticles with antibacterial properties.

Table 9. Studies of carrot, cabbage, and red beet by-products in electrical use

By-products as raw materials	Active component or structure and product or application examples	Reference
Rotten carrots and other vegetable by-products	Porous activated carbon-based electrode for energy storage applications	Ahmed et al. 2018
Carrot discards	Precursor for Ni/P-doped carbon composite electrocatalyst for e.g. batteries, fuel cells, and electrocatalytic water splitting	Hoang et al. 2019
Carrot peel	Synthesis of Cu ₂ O nanoparticles with antibacterial properties from carrot peel extract	Mannarmannan and Biswas 2021
Red beet discards	Single-crystalline gold (Au) nanoplates from extract for use, e.g., in catalysis, photonics, biosensing, electronics, or nanomedicines	Deokar and Ingale 2018
Cabbage outer leaves	Carbon aerogel for supercapacitors and adsorbent for oil/water separation	Cai et al. 2017

Discussion

Valorization options

This review summarizes the literature on valorization options for by-products of carrot, red beet, and cabbage that are important vegetables in northern climates. The review showed that there is no single answer about the best use for the by-products of the vegetables under focus. The Waste Framework Directive sets the basic concepts and definitions related to waste management. Preventing waste is the preferred option, and preparing by-products for re-use, recycling, recovery, and sending waste to landfill should be the last options (EU 2008). Vegetable wastes are rich sources of vitamins and minerals for producing high-value compounds or metabolites (Yadav et al. 2023). Studies on various valorization options provide valuable insights into the potential uses of vegetable by-products for different-sized production companies. Some potential for using the by-products of carrot, red beet, and cabbage in food applications was observed. There were only a few studies about these by-products in medicine and cosmetics, and food and pharmaceutical uses were often lumped together. These by-products or discards are used as such in animal feed, but quality upgrading for different animal species appeared positive in several studies. According to Ajila et al. (2012), the production of animal feed from agro-residues could be one of the most suitable technologies for finding an effective way to attain income by the agro-community and for the better management of environmental pollution. On the other hand, Esparza et al. (2020) mentioned bioactive compounds, enzymes, exopolysaccharides, bioplastics, and biofuels as the most promising options for valorization. In addition to food applications, Rapa et al. (2024) mentioned biogas and organic fertilizers, animal feed and innovative food packaging, and construction and water purification materials as potential applications.

Particularly, cabbage was promising for biogas production via anaerobic digestion. By-products, particularly of carrot and cabbage, appeared suitable for producing biochar that can be used as a fertilizer or to remove contaminants, such as heavy metals or phosphorus, from the environment. Carrot or cabbage by-products in the form of activated carbon were common in wastewater purification. Both activated carbon, lignocellulose, and pectin from the by-products were mostly potential as biosorbents. Composting of cabbage for soil amendment was mostly promising, while studies on other treatments for soil improvement and biopesticides were few and contradictory. Carrot and red beet by-products had potential in the production of films and packaging materials. Carrot or cabbage wastes showed potential for the production of the enzymes, and red beet also for textile dyes. However, in this case, contradictory results were found too. Some technical valorization options, e.g., electrical products, were examined only in a few studies, in which the by-products were nevertheless found functionally as potential product sectors.

In practice, the most profitable valorization options are those related to centralizing the production and capitalizing on economies of scale (Cristobal et al. 2018). For example, the production of flour from vegetable by-products is affordable only if high value-added ingredients and products are developed (Ratti 2001). Economic profitability was not the focus of many of the studies reviewed. Technical feasibility, including pre-treatment, extraction and production methods, is highly important for both production technical and economic reasons. Sustainable recovery technologies eliminate harmful effects compared to conventional technologies (Czubaszek 2022). Green extraction methods in the separation of various bioactive compounds from vegetable by-products correspond to the aims of the circularity of resources and sustainability (Rapa et al. 2024). Many studies exploit technologies such as ultrasounds, microwaves, high pressure, and supercritical fluid processing, for by-product processing. However, these incur huge investment and maintenance costs, along with requiring specialized know-how and production plants (Galanakis 2013). The costs are high even when using technologies characterized by a high industrial maturity level, e.g., air-drying (Ferreira et al. 2015). For example, biofuel applications (approximately 186–372 €/tonne biomass) create more value compared to generating electricity (approximately 56–139 €/tonne biomass) and cattle feed (65–186 €/tonne biomass), but the turnover that could be generated by high-value biochemicals is much higher and estimated to be around 930 €/tonne biomass (Tuck et al. 2012). An accurate cost-benefit analysis should be performed to evaluate the environmental and economic sustainability of the proposed vegetable waste valorization strategies (Coelho et al. 2020).

In this review, valorization options were examined, including all by-product sources. However, industrial by-products and wastes are typically more homogeneous and constant in composition than those generated during the consumption phase, which makes them a more attractive option for valorization (Fava et al. 2015, Rao et al. 2021). In addition, the type of by-product affects its properties and thus its use potential, although the part of the vegetable examined as a by-product was not specified in all studies, or vegetable waste was examined as a whole. For example, red beet has a high antioxidant capacity (Porto Dalla Costa et al. 2017, Šeremet et al. 2020, Salamatullah

et al. 2021, Zin and Bánvölgyi 2021, El-Beltagi et al. 2022), but the composition is not equal in all parts of the vegetable (Kujala et al. 2000). In Zin and Bánvölgyi (2022), the contents of betalains, phenolics, and flavonoids were also the highest in red beet peel, followed by the flesh and stalk. Based on this review, industrial-scale studies are needed to confirm the utilization of valuable components of vegetable by-products.

Challenges in valorization of by-products

According to Socas-Rodríguez et al. (2021), the valorization of food by-products is challenging for several reasons: the different stabilities of the added-value components during processing, the technological difficulty in large-scale production, the low energy efficiency and high costs of conventional extraction processes, and the use of non-food-grade solvents during conventional extraction processes. A central challenge is the perishable nature and heterogeneity of vegetable by-products and logistical arrangements (Esparza et al. 2020). The presence of moisture is connected to the stability and microbiological safety of food products (San Martín et al. 2016). For this reason, many processes applied for the valorization of food waste involve a drying step in which the water content is preferably reduced to less than 12% (Gomes et al. 2020, Iriondo-DeHond et al. 2018, Ribeiro et al. 2020, San Martín et al. 2016). Drying consumes a lot of energy, and in the northern countries, it is expensive when solar energy is limited.

Finding reference values that allow determining the suitability of the obtained by-products for certain uses has been difficult, as no legislation has specifically been established in this regard. The transfer of contaminants from the raw material to the valorized product must be evaluated, along with the generation of new hazardous substances or their enrichment during food by-product processing (Socas-Rodríguez et al. 2021). The lack of regulation creates risks for the valorization of many potentially useful by-products from the agro-food industry. This problem brings about the necessity for determining specific regulations for food by-product valorization and safety (Socas-Rodríguez et al. 2021). By-products from organic farming lack residual pesticides or potentially toxic chemicals (Barbulova et al. 2015), which is a benefit for food applications.

Environmental sustainability

The strategy of the Circular Economy Action Plan is to use waste material and energy as inputs for other industrial processes or as regenerative resources for nature (EU 2020). However, when developing the processing chains of vegetables, waste prevention should also be emphasized. The US Environmental Protection Agency's waste hierarchy ranks waste prevention as the most preferable option, and energy recovery as a less preferable strategy of waste management compared to others (EPA 2023).

Vegetable by-products are of lower consequence to the environment compared to some other waste streams, but their leachates and CH₄ emissions still present a risk (Misi and Foster 2002). According to the Food and Agriculture Organization of the United Nations (FAO), food loss and waste is the second-highest cause of greenhouse gas emissions. Statistically, 1.3 billion tonnes of wasted food caused approximately 4.4 gigatonnes of greenhouse gas emissions (FAO 2015). Vegetable by-products obtained from the food bioprocessing industry remain a major surplus, leading to environmental pollution (FAO 2013). The global carbon footprint of food loss and waste, excluding emissions from land use change, is estimated at 3.3 gigatonnes of CO₂ equivalent, while surface and groundwater resource use is attributable to circa 250 km³ of food lost or wasted, and almost 1.4 billion hectares are used to produce food that is later lost or wasted (FAO 2013). Approximately 40 823 kg of carrot rejects and waste destined for landfills released 6 988 kg of CH₄, equivalent to 174 710 kg of CO₂ (Kaur et al. 2022a). Improper disposal of vegetable waste and combustion leads to environmental pollution with the release of dioxins. Landfilling with vegetable wastes also causes water and air pollution and leads to CH₄ gas emissions (ElMekawy et al. 2015).

Implementing non-thermal technologies, adding sustainable solvents and safer materials, and possessing GRAS (Generally Recognized As Safe) status are strongly recommended. Conducting integral investigations that include recovery protocols and preservation assays is necessary to ensure industrial exploitation and sustainability of the final product (Galanakis et al. 2015). Innovative valorization strategies have been shown to maximize the economic performance of the considered system (Plazzotta et al. 2020). A holistic approach is needed for utilizing vegetable by-products, including environmental aspects, technical feasibility, processing techniques, and economic profitability. Holistic examination of the sustainability of by-products, including different processing techniques, is also needed, particularly via life cycle assessment.

Conclusions

This literature review showed that plenty of valorization ideas in a wide variety of applications, including technical applications, have been examined for carrot, cabbage, and red beet by-products. Thus, there is not a single best use for the by-products of the vegetables under focus. Food applications were the most examined product sector. More studies for the food sector were found concerning carrots and red beets when compared to cabbage. Most studies indicated at least some potential for these by-products in food use. Fewer studies focused on pharmaceuticals and cosmetics. Vegetable by-products are often used untreated for animal feed, but valorization studies for several animal groups were also found. Technical applications as valorization targets included biosorbents (mostly based on activated carbon), composites and films (mostly food packages), energy products (mostly biogas but also bioethanol and biodiesel), soil amendments and pesticides, enzymes, dyes and biosurfactants, and electrical applications.

From an economic point of view, balancing between the amount, quality, logistics, and processing technologies of the by-products is a key issue. According to the literature, high-value biochemicals may be among the most profitable valorization options. To handle the significant amounts of by-product biomass created, tailored nutraceutical or feed additive products, bioplastics, or biofuels might also be realistic options on a large scale. However, since most of the studies reviewed were of laboratory scale, research on an industrial scale is needed.

Food loss and waste are a significant cause of greenhouse gas emissions. Concerning environmental sustainability, waste prevention is the primary option. Vegetable waste contains valuable biomass and nutrients, and the valorization of vegetable by-products can result in relative environmental benefits. To decrease emissions from the treatment of vegetable by-products, non-thermal technologies and sustainable solvents should be considered.

The transformation of vegetable by-products into new products with maximum added value aligns with the precepts of the circular economy and the desire for natural ingredients. Many studies on the use of vegetable residues of interest in northern climates were reviewed from the viewpoints of valorization, but it is important to increase the scope of such studies, interrelating many other aspects that have a direct impact on the amount, type, and management of the generated residues. More studies about the economic and environmental aspects of utilizing vegetable by-products are needed. Many product ideas and research results exist, and these could be used, for example, in creating a company's by-product utilization strategy.

References

- Abdo, E.M, Shaltout, O.E-S., Ali, S. & Mansour, H.M.M. 2022. A functional orange juice fortified with beetroot by-products attenuates hyperlipidemia and obesity induced by a high-fat diet. *Antioxidants* 11: 457. <https://doi.org/10.3390/antiox11030457>
- Aguirre Calvo, T.R., Santagapita, P.R. & Perullini, M. 2019. Functional and structural effects of hydrocolloids on Ca(II)-alginate beads containing bioactive compounds extracted from beetroot. *Food Science and Technology* 111: 520-526. <https://doi.org/10.1016/j.lwt.2019.05.047>
- Ahmed, S., Ahmed, A. & Rafat, M. 2018. Supercapacitor performance of activated carbon derived from rotten carrot in aqueous, organic and ionic liquid based electrolytes. *Journal of Saudi Chemical Society* 22: 9931002. <https://doi.org/10.1016/j.jscs.2018.03.002>
- Aimaretti, N. & Ybalo, C. 2012. Valorization of carrot and yeast discards for the obtention of ethanol. *Biomass & Bioenergy* 42: 18–23. <https://doi.org/10.1016/j.biombioe.2012.03.022>
- Aimaretti, N.R., Ybalo, C.V., Rojas, M.L., Plou, F.J. & Yori, J.C. 2012. Production of bioethanol from carrot discards. *Bioresource Technology* 123: 727–732. <https://doi.org/10.1016/j.biortech.2012.08.035>
- Ajila, C.M., Brar, S.K., Verma, M., Tyagi, R.D., Godbout, S. & Valéro, J.R. 2012. Bio-processing of agro-byproducts to animal feed. *Critical Reviews in Biotechnology* 32: 382–400. <https://doi.org/10.3109/07388551.2012.659172>
- Alam, Md S., Khaira, H., Pathania, S., Kumar, S. & Singh, B. 2015. Extrusion process optimization for soy-carrot pomace powder incorporated wheat-based snacks. *Journal of Agricultural Engineering* 53: 1–13.
- Amany, E.M., El-Abasy, Youssef, M.M., Abou-Gharbia, H.A. & Hamida, M.M. 2010. Utilization of carrot pomace in formulating functional biscuits and cakes. *Alexandria Journal of Food Science & Technology* 7: 25–32. <https://doi.org/10.21608/ajfs.2010.29081>
- Amin, S., Jung, S., Kang, I. & Duval, A. 2021. Valorization of baby carrot processing waste. *Journal of Culinary Science & Technology* 21: 1–17. <https://doi.org/10.1080/15428052.2021.1879338>
- Amoroso, L., De France, K.J., Milz, C.I., Siqueira, G. Zimmermann, T. & Nyström, G. 2022. Sustainable cellulose nanofiber films from carrot pomace as sprayable coatings for food packaging applications. *ACS Sustainable Chemistry & Engineering* 10: 342–352. <https://doi.org/10.1021/acssuschemeng.1c06345>
- Anal, A.K. 2018. Food processing by-products and their utilization. John Wiley & Sons Ltd., Hoboken, NJ. 570 p. <https://doi.org/10.1002/9781118432921>

- Andrade, M., Rabelo de Oliveira Torres, L., Sanches Silva, A., Barbosa, C.H., Vilarinho, F., Ramos, F., Rodrigues Bernaldo de Quirós, A., Khwaldia, K. & Sendon, R. 2020. Industrial multi-fruits juices by-products: total antioxidant capacity and phenolics profile by LC-MS/MS to ascertain their reuse potential. *European Food Research and Technology* 246: 2271–2282. <https://doi.org/10.1007/s00217-020-03571-3>
- de Andrade Lima, M., Charalampopoulos, D. & Chatzifragkou, A. 2018. Optimisation and modelling of supercritical CO₂ extraction process of carotenoids from carrot peels. *Journal of Supercritical Fluids* 133: 94–102. <https://doi.org/10.1016/j.supflu.2017.09.028>
- Araújo-Rodrigues, H., Santos, D., Campos, D.A., Ratinho, M., Rodrigues, I.M. & Pintado, M.E. 2021. Development of frozen pulps and powders from carrot and tomato by-products: Impact of processing and storage time on bioactive and biological properties. *Horticulturae* 7: 2–22. <https://doi.org/10.3390/horticulturae7070185>
- Arifan, F., Abdullah, A. & Sumardiono, S. 2021. Effect of organic waste addition into animal manure on biogas production using anaerobic digestion method. *International Journal of Renewable Energy Development* 10: 623–633. <https://doi.org/10.14710/ijred.2021.36107>
- Aristi, H., Mudji, E.H., Soepriananondo, K., Haridjani, N., Aprilia, Z., Gholib, G., Duboscq, J., Sutriana, A., Sahara Z.R. & Engelhardt, A. 2020. Levels of protein and fat produced by black soldier fly (*Hemestia illucens*) larvae in the bioconversion of organic waste. *E3S web of conferences* 151: 1041. <https://doi.org/10.1051/e3sconf/202015101041>
- Arrais, A., Testori, F., Calligari, R., Gianotti, V., Roncoli, M., Caramaschi, A., Todeschini, V., Massa, N. & Bona, E. 2022. Extracts from cabbage leaves: Preliminary results towards a “universal” highly-performant antibacterial and antifungal natural mixture. *Biology* 11: 1080. <https://doi.org/10.3390/biology11071080>
- Asadi, S.Z. & Khan, M.A. 2021. The effect of beetroot (*Beta vulgaris* L.) leaves powder on nutritional, textural, sensorial and antioxidant properties of cookies. *Journal of Culinary Science & Technology* 19: 424–438. <https://doi.org/10.1080/15428052.2020.1787285>
- Babalola, O.O. 2018 Comparative evaluation of lettuce wastes, pawpaw leaves, whole lettuce and cabbage wastes as a sole feed ingredient for growing *Archachatina marginata* snails. *Agricultural Resources* 7: 83–88. <https://doi.org/10.1007/s40003-018-0289-7>
- Bakshi, M.P.S., Wadhwa, M. & Makkar, P.S. 2016. Waste to worth: vegetable wastes as animal feed. *CAB Review* 11: 1–26. <https://doi.org/10.1079/PAVSNNR201611012>
- Bakshi, S., Banik, C., Laird, D.A., Smith, R.G. & Brown, R. 2021. Enhancing biochar as scaffolding for slow release of nitrogen fertilizer. *ACS Sustainable Chemistry & Engineering* 9: 1–17. <https://doi.org/10.1021/acssuschemeng.1c02267>
- Baljeet, S.Y., Ritika, B.Y. & Reena, K. 2014. Effect of incorporation of carrot pomace powder and germinated chickpea flour on the quality characteristics of biscuits. *International Food Resource Journal* 21: 217–222.
- Barbulova, A., Colucci, G. & Apone, F. 2015. New trends in cosmetics: By-products of plant origin and their potential use as cosmetic active ingredients. *Cosmetics* 2: 82–92. <https://doi.org/10.3390/cosmetics2020082>
- Battistella Lasta, H.F., Lentz, L., Mezzomo, N. & Ferreira, S.R.S. 2019. Supercritical CO₂ to recover extracts enriched in antioxidant compounds from beetroot aerial parts. *Biocatalysis and Agricultural Biotechnology* 19: 101169. <https://doi.org/10.1016/j.cbab.2019.101169>
- Ben-Othman, S., Joudu, I. & Bhat, R. 2020. Bioactives from agri-food wastes: Present insights and future challenges. *Molecules* 25: 510. <https://doi.org/10.3390/molecules25030510>
- Berglund, L., Noël, M., Aitomäki, Y., Öman, T. & Oksman, K. 2016. Production potential of cellulose nanofibers from industrial residues: Efficiency and nanofiber characteristics. *Industrial Crops and Products* 92: 84–92. <https://doi.org/10.1016/j.indcrop.2016.08.003>
- Bhatti, H.N., Nasir, A.W. & Hanif, M.A. 2010. Efficacy of *Daucus carota* L. waste biomass for the removal of chromium from aqueous solutions. *Desalination* 253: 78–87. <https://doi.org/10.1016/j.desal.2009.11.029>
- Bozym, M., Florczak, I., Zdanowska, P., Wojdalski, J. & Klimkiewicz, M. 2015. An analysis of metal concentrations in food wastes for biogas production. *Renewable Energy* 77: 467–472. <https://doi.org/10.1016/j.renene.2014.11.010>
- Brito, T.B.N., Pereira, A.P.A., Pastore, G.M., Moreira, R.F.A., Ferreira, M.S.L & Fai, A.E.C 2020. Chemical composition and physico-chemical characterization for cabbage and pineapple by-products flour valorization. *Food Science and Technology* 124: 109028. <https://doi.org/10.1016/j.lwt.2020.109028>
- van Broekhoven, S., Oonincx, D.G.A.B., van Huis, A. & van Loon, J.J.A. 2015. Growth performance and feed conversion efficiency of three edible mealworm species (Coleoptera: Tenebrionidae) on diets composed of organic by-products. *Journal of Insect Physiology* 3: 1–10. <https://doi.org/10.1016/j.jinsphys.2014.12.005>
- Cai, T., Wang, H., Jin, C., Sun, Q. & Nie, Y. 2017. Fabrication of nitrogen-doped porous electrically conductive carbon aerogel from waste cabbage for supercapacitors and oil/water separation. *Journal of Materials Science: Materials in Electronics* 29: 4334–4344. <https://doi.org/10.1007/s10854-017-8381-5>
- Caldeira, C., De Laurentiis, V., Corrado, S., van Holsteijn, F. & Sal, S. 2019. Quantification of food waste per product group along the food supply chain in the European Union: a massflow analysis. *Resources, Conservation & Recycling* 149: 479–488. <https://doi.org/10.1016/j.resconrec.2019.06.011>
- de Carvalho Eufrásio Pinto, M., da Silva, D., Gomes, A., Luiza, A., dos Santos, M., Morais, R., de Couto, A., Alexandre, R., Ferreira de Novais, R., Constantino, L., Regina, V., Jairo, T. & Garcia, P.F. 2019. Biochar from carrot residues chemically modified with magnesium for removing phosphorus from aqueous solution. *Journal of Cleaner Production* 222: 36–46. <https://doi.org/10.1016/j.jclepro.2019.03.012>
- Castrica, M., Tedesco, D.E.A., Panseri, S., Ferrazzi, G., Ventura, V., Frisio, D.G. & Balzaretto, C.M. 2018. Pet food as the most concrete strategy for using food waste as feedstuff within the European context: A feasibility study. *Sustainability* 10: 2035. <https://doi.org/10.3390/su10062035>
- Changmai, M., Banerjee, P., Nahar, K. & Purkait, M.K. 2018. A novel adsorbent from carrot, tomato and polyethylene terephthalate waste as a potential adsorbent for Co(II) from aqueous solution: Kinetic and equilibrium studies. *Journal of Environmental Chemical Engineering* 6: 246–257. <https://doi.org/10.1016/j.jece.2017.12.009>

- Chantaro, P., Devahastin, S. & Chiewchan, N. 2008. Production of antioxidant high dietary fiber powder from carrot peels. *LWT* 41: 1987–1994. <https://doi.org/10.1016/j.lwt.2007.11.013>
- Chauhan, S. & Rajput, H. 2018. Production of gluten free and high fibre cookies using beet root waste powder and wheat flour husk. *The Pharma Innovation Journal* 7: 556–558.
- Christiaens, S., Uwibambe, D., Uyttebroek, M., Van Droogenbroeck, B., Van Loey, A.M. & Hendrickx, M.E. 2015. Pectin characterisation in vegetable waste streams: A starting point for waste valorisation in the food industry. *Food Science and Technology* 61: 275–282. <https://doi.org/10.1016/j.lwt.2014.12.054>
- Coelho, M.C., Pereira, R.N., Rodrigues, A.S., Teixeira, J.A. & Pintado, M.E. 2020. The use of emergent technologies to extract added value compounds from grape by-products. *Trend in Food Science & Technology* 106: 182–197. <https://doi.org/10.1016/j.tifs.2020.09.028>
- Coman, V., Teleky, B.-E., Mitrea, L., Martau, G.A., Szabo, K., Calinoiu, L.-F. & Vodnar, D.C. 2020. Bioactive potential of fruit and vegetable wastes. *Advances in Food and Nutrition Research* 91: 157–225. <https://doi.org/10.1016/bs.afnr.2019.07.001>
- Cristobal, J., Caldeira, C., Corrado, S. & Sala, S. 2018. Techno-economic and profitability analysis of food waste biorefineries at European level. *Bioresource Technology* 259: 244–252. <https://doi.org/10.1016/j.biortech.2018.03.016>
- Czubaszek, R., Wysocka-Czubaszek, A. & Tyborowski, R. 2022. Methane Production Potential from Apple Pomace, Cabbage Leaves, Pumpkin Residue and Walnut Husks. *Applied Science* 12: 6128. <https://doi.org/10.3390/app12126128>
- Das, A., Ghosh, U., Mohapatra, P.K.D., Pati, B.R. & Mondal, K.C. 2012. Study on thermodynamics and adsorption kinetics of purified endoglucanase (CMCase) from *Penicillium notatum* NCIM NO-923 produced under mixed solid-state fermentation of waste cabbage and bagasse. *Brazilian Journal of Microbiology* 43: 1103–1111. <https://doi.org/10.1590/S1517-83822012000300037>
- Deokar, G.K. & Ingale, A.G. 2018. Unveiling an unexpected potential of beetroot waste in green synthesis of single crystalline gold nanoplates: A mechanistic study. *Arabian Journal of Chemistry* 11: 950–958. <https://doi.org/10.1016/j.arabjc.2018.03.016>
- Di Donato, P., Fiorentino, G., Anzelmo, G., Tommonaro, G., Nicolaus, B. & Poli, A. 2011. Re-use of vegetable wastes as cheap substrates for extremophile biomass production. *Waste and Biomass Valorization* 2: 103–111. <https://doi.org/10.1007/s12649-011-9062-x>
- Du, G., Shi, J., Zhang, J., Ma, Z., Liu, X., Yuan, C., Zhang, B. & Zhang, Z. & Harrison, M.D. 2021. Exogenous probiotics improve fermentation quality, Microflora phenotypes, and trophic modes of fermented vegetable waste for animal feed. *Microorganisms* 9: 644. <https://doi.org/10.3390/microorganisms9030644>
- Duval, A. 2020. Valorization of carrot processing waste. A. thesis. The faculty of California polytechnic state university. San Luis Obispo. [https://digitalcommons.calpoly.edu/cgi/viewcontent.cgi?article=3694&context=theses\(2020\)](https://digitalcommons.calpoly.edu/cgi/viewcontent.cgi?article=3694&context=theses(2020)). Accessed 3 June 2023.
- El-Beltagi, H.S., El-Mogy, M.M., Parmar, A., Mansour, A.T., Shalaby, T.A. & Ali, M.R. 2022. Phytochemical characterization and utilization of dried red beetroot (*Beta vulgaris*) peel extract in maintaining the quality of Nile Tilapia fish fillet. *Antioxidants* 11: 906. <https://doi.org/10.3390/antiox11050906>
- ElMekawy, A., Srikanth, S., Bajracharya, S., Hegab, H.M., Nigam, P.S., Singh, A. & Pant, D. 2015. Food and agricultural wastes as substrates for bioelectrochemical system (BES): The synchronized recovery of suitable energy and waste treatment. *Food Research International* 73: 213–225. <https://doi.org/10.1016/j.foodres.2014.11.045>
- El-Sawi, S.A., Ibrahim, M.E., Sleem, A.A., Farghaly, A.A., Awad, G.E.A. & Merghany, R.M. 2022. Development of alternative medicinal sources from golden berry, bananas and carrot wastes as antioxidant, cytotoxic and antimicrobial agents. *Acta Ecologica Sinica* 42: 224–232. <https://doi.org/10.1016/j.chnaes.2021.04.006>
- EPA 2023. US Environmental Protection Agency waste hierarchy EPA 2023 <https://www.epa.gov/smm/sustainable-materials-management-non-hazardous-materials-and-waste-management-hierarchy>. Accessed 3 June 2023.
- Enaime, G. & Lübken, M. 2021. Agricultural Waste-Based Biochar for Agronomic Applications. *Applied Sciences* 11: 8914. <https://doi.org/10.3390/app11198914>
- Esparza, I., Jimenez-Moreno, M., Bimbela, F., Ancin-Azpilicueta, C., Luis M. & Gandía, L.M. 2020. Fruit and vegetable waste management: Conventional and emerging approaches. *Journal of Environmental Management* 265: 110510. <https://doi.org/10.1016/j.jenvman.2020.110510>
- EU 2020. European Union, Outcome Report FOOD 2030 Pathways Workshop. Future Research & Innovation Needs in view of the transition to sustainable, healthy, safe and inclusive food systems. <https://fit4food2030.eu/wp-content/uploads/2020/04/20200424-Food-2030-Pathways-Workshop-Report.pdf>. Accessed 3 June 2023.
- EU 2008. The Waste Framework Directive 98/EC. https://environment.ec.europa.eu/topics/waste-and-recycling/waste-framework-directive_en. Accessed 20 July 2025.
- de Evan, T., Vintimilla, A., Marcos, C.N., Ranilla, M.J. & Carro, M.D. 2019. Evaluation of Brassica vegetables as potential feed for ruminants. *Animals* 9: 588. <https://doi.org/10.3390/ani9090588>
- FAO 2011. Global food losses and food waste. Study conducted for the international congress Save food! Düsseldorf, Germany.
- AO 2013. Food Wastage Footprint Impacts on Natural Resources. <http://www.fao.org/3/i3347e/i3347e.pdf> Accessed 3 June 2023
- FAO 2015. Food Loss and Food Waste. <http://www.fao.org/food-lossand-foodwaste/en/>. Accessed 3 June 2023
- FAO 2019. Key factors on food loss and waste you should know. <http://www.fao.org/savefood/resources/keyfindings/en/> Accessed 3 June 2023
- Fava, F., Totaro, G., Diels, L., Reis, M., Duarte, J., Carioca, B.O., Poggi-Varaldo, H.M. & Ferreira, S.B. 2015. Biowaste biorefinery in Europe: opportunities and research & development needs. *New Biotechnology* 32: 100–108. <https://doi.org/10.1016/j.nbt.2013.11.003>
- Fernando, G.S.N., Wood, K., Papaioannou, E.H., Marshall, L.J., Sergeeva, N.N. & Boesch, C. 2021. Application of an ultrasound-assisted extraction method to recover betalains and polyphenols from red beetroot waste. *ACS Sustainable Chemistry & Engineering* 9: 8736–8747. <https://doi.org/10.1021/acssuschemeng.1c01203>

- Ferreira, M.S.L., Santos, M.C.P., Moro, T.M.A., Basto, G.J., Andrade, R.M.S. & Gonçalves, E.C.B.A. 2015. Formulation and characterization of functional foods based on fruit and vegetable residue flour. *Journal of Food Science and Technology* 52: 822–830. <https://doi.org/10.1007/s13197-013-1061-4>
- Galanakis, C.M. 2012. Recovery of high added-value components from food wastes: Conventional, emerging technologies and commercialized applications. *Trends of Food Science Technology* 26: 68–87. <https://doi.org/10.1016/j.tifs.2012.03.003>
- Galanakis, C.M. 2013. Emerging technologies for the production of nutraceuticals from agricultural by-products: A viewpoint of opportunities and challenges. *Food and Bioproducts Processing* 91: 575–579. <https://doi.org/10.1016/j.fbp.2013.01.004>
- Galanakis, C.M., Martinez-Saez, N., del Castillo, M.D., Barba, F.J. & Mitropoulou, V.S. 2015. Patented and commercialized applications. In: *Food Waste Recovery*. Elsevier Inc. Chapter 15. p. 337–360. <https://doi.org/10.1016/B978-0-12-816449-5.00015-8>
- Galanakis, C. 2020. Food waste valorization opportunities for different food industries. In: Galanakis, C.M. (ed.). *The Interaction of Food Industry and Environment*. Elsevier, London. p. 341–422. <https://doi.org/10.1016/B978-0-12-816449-5.00011-4>
- Garces, C.L. & Heinen, J.M. 1993. Evaluation of beef liver, fruits, and vegetables as dietary supplements for indoor nursery culture of freshwater prawns *Macrobrachium rosenbergii*. *Journal of the World Aquaculture Society* 24: 58–65. <https://doi.org/10.1111/j.1749-7345.1993.tb00150.x>
- George, S. & Jayachandran, K. 2008. Analysis of Rhamnolipid Biosurfactants Produced Through Submerged Fermentation Using Orange Fruit Peelings as Sole Carbon Source. *Applied Biochemistry & Biotechnology* 158: 694–705. <https://doi.org/10.1007/s12010-008-8337-6>
- Gholami, L. & Rahimi, G. 2021. Chemical fractionation of copper and zinc after addition of carrot pulp biochar and thiourea-modified biochar to a contaminated soil. *Environmental Technology* 42: 3523–3532. <https://doi.org/10.1080/09593330.2020.1733101>
- Giller, K., Bossut, L., Eggenschwiler, L. & Terranova, M. 2021. In vitro ruminal fermentation, methane production and nutrient degradability as affected by fruit and vegetable pomaces in differing concentrations. *Journal of Animal Physiology and Animal Nutrition* 106: 957–967. <https://doi.org/10.1111/jpn.13656>
- Gomes, L.A., Santos, A.F., Pinhiera, C.T., Gois, J.C. & Quina, M.J. 2020. Screening of waste materials as adjuvants for drying sewage sludge based on environmental, technical and economic criteria. *Journal of Cleaner Production* 259: 120927. <https://doi.org/10.1007/978-3-030-23165-1>
- Gonzales, G.B., Raes, K., Vanhoutte, H., Coelus, S., Smagghe, G. & Van Camp, J. 2015. Liquid chromatography-mass spectrometry coupled with multivariate analysis for the characterization and discrimination of extractable and nonextractable polyphenols and glucosinolates from red cabbage and Brussels sprout waste streams. *Journal of Chromatography A* 1402: 60–70. <https://doi.org/10.1016/j.chroma.2015.05.009>
- Green, B.N., Johnson, C.D. & Adams, A. 2006. Writing narrative literature reviews for peer-reviewed journals: secrets of the trade. *Journal of Chiropractic Medicine* 5: 101–117. [https://doi.org/10.1016/S0899-3467\(07\)60142-6](https://doi.org/10.1016/S0899-3467(07)60142-6)
- Gül, H., Yanik, A. & Acun, S. 2013 Effects of white cabbage powder on cookie quality. *Journal of Food, Agriculture & Environment* 11: 69–72.
- Hammershøj, M., Kidmose, U. & Steinfeldt, S. 2010. Deposition of carotenoids in egg yolk by short-term supplement of coloured carrot (*Daucus carota*) varieties as forage material for egg-laying hens. *J. Science of Food and Agriculture* 90: 1163–71. <https://doi.org/10.1002/jsfa.3937>
- Hamoudi-Belarbi, L., Hamoudi, S., Belkacemi, K., Nouri, L., Bendifallah, L. & Khodja, M. 2018. Bioremediation of polluted soil sites with crude oil hydrocarbons using carrot peel waste. *Environments (Basel)* 5: 124. <https://doi.org/10.3390/environments5110124>
- Hang, B.P.T., Lam, V. & Preston, T.R. 2011. Effect of different sources of carbohydrate as supplements to basal diets of water spinach and cabbage leaves on the growth rate in rabbits. *Livestock Research of Rural Development* 23: 227.
- Hastuti, B., Totiana, F. & Winiasih, R. 2018. The role of pectin in Pb binding by carrot peel biosorbents: Isoterm adsorption study. IOP conference series. *Materials Science and Engineering* 349: 12011. <https://doi.org/10.1088/1757-899X/349/1/012011>
- Hidalgo, A., Brandolini, A., Čanadanović-Brunet, J., Četković, G. & Tumbas Šaponjac, V. 2018. Microencapsulates and extracts from red beetroot pomace modify antioxidant capacity, heat damage and colour of pseudocereals-enriched einkorn water biscuits. *Food Chemistry* 268: 40–48. <https://doi.org/10.1016/j.foodchem.2018.06.062>
- Hira, S.A., Yusuf, M., Annas, D., Hui, H.S. & Park, K.H. 2020. Biomass-Derived Activated Carbon as a Catalyst for the Effective Degradation of Rhodamine B dye. *Processes* 8: 926. <https://doi.org/10.3390/pr8080926>
- Hoang, V.C., Gomes, V.G. & Dinh, K.N. 2019. Ni- and P-doped carbon from waste biomass: A sustainable multifunctional electrode for oxygen reduction, oxygen evolution and hydrogen evolution reactions. *Electrochimica Acta* 314: 49–60. <https://doi.org/10.1016/j.electacta.2019.05.053>
- Hossain, M.A., Ngo, H.H., Guo, W.S., Nghiem, L.D., Hai, F.I., Vigneswaran, S. & Nguyen, T.V. 2014. Competitive adsorption of metals on cabbage waste from multi-metal solutions. *Bioresources Technology* 160: 79–88. <https://doi.org/10.1016/j.biortech.2013.12.107>
- Hussein, A.M.S., Kamil, M.M., Hegazy, N.A., Mahmoud, K.F. & Ibrahim, M.A. 2015. Utilization of some fruits and vegetables by-products to produce high dietary fiber jam. *Food Science and Quality Management* 37: 39–45.
- Iriondo-DeHond, M., Miguel, E. & del Castilla, M.D. 2018. Food by-products as sustainable ingredients for innovative and healthy dairy foods. *Nutrients* 10: 1358. <https://doi.org/10.3390/nu10101358>
- Janiszewska-Turak, E., Kotakowska, W., Pobiega, K. & Gramza-Michałowska, A. 2021. Influence of drying type of selected fermented vegetables pomace on the natural colorants and concentration of lactic acid bacteria. *Applied Science* 11: 7864. <https://doi.org/10.3390/app11177864>
- Joel, E.B., Mafulul, S.G., Adamu, H.E., Goje, L.J., Tijjani, H., Igunnu, A. & Malomo, S.O. 2020. Peroxidase from waste cabbage (*Brassica oleracea capitata* L.) exhibits the potential to biodegrade phenol and synthetic dyes from wastewater. *Scientific African* 10: e00608. <https://doi.org/10.1016/j.sciaf.2020.e00608>

- Jongaroontaprangsee, S., Tritromg, W. & Chokanaporn, W. 2007. Effects of drying temperature and particle size on hydration properties of dietary fiber powder from lime and cabbage by-products. *International Journal of Food Properties* 10: 887–897. <https://doi.org/10.1080/10942910601183619>
- Jovanović, M., Zlatanovic, S., Micic, D., Bacic, D., Mitic-Culafic, D., Duris, N. & Gorjanovic, S. 2021. Functionality and palatability of yogurt produced using beetroot pomace flour granulated with lactic acid bacteria. *Foods* 10: 1696. <https://doi.org/10.3390/foods10081696>
- Karatay, S.E., Demiray, E. & Dönmez, G. 2020. Usage potential of apple and carrot pomaces as raw materials for newly isolated yeast lipid-based biodiesel production. *Biomass Conversion and Biorefinery* 12: 4773–4783. <https://doi.org/10.1007/s13399-020-01174-2>
- Karatay, S.E., Demiray, E. & Dönmez, G. 2019. Efficient approaches to convert *Coniochaeta hoffmannii* lipids into biodiesel by *in-situ* transesterification. *Bioresources Technology* 285: 121321. <https://doi.org/10.1016/j.biortech.2019.121321>
- Kaur, N., Kaur, A., Sridhar, K., Sharma, M., Singh, T.P. & Kumar, S. 2021. Development and quality characteristics of functional Kulfi fortified with microencapsulated betalains. *International Journal of Food Science & Technology* 56: 5362–5370. <https://doi.org/10.1111/ijfs.15139>
- Kaur, G.J., Kumar, D., Orsat, V. & Singh, A. 2022a. Assessment of carrot rejects and wastes for food product development and as a biofuel. *Biomass Conversion and Biorefinery* 12: 757–768. <https://doi.org/10.1007/s13399-020-01096-z>
- Kaur, G.J., Orsat, V. & Singh, A. 2022b. Sustainable usage of carrot discards in food processing. *International Journal of Sustainable Development & World Ecology* 29: 18–26. <https://doi.org/10.1080/13504509.2021.1896589>
- Khan, M.K.I., Asif, M., Razzaq, Z.U., Nazir, A. & Maan, A.A. 2022. Sustainable food industrial waste management through single cell protein production and characterization of protein enriched bread. *Food Bioscience* 46: 101406. <https://doi.org/10.1016/j.fbio.2021.101406>
- Khoshkho, S.M., Mahdavian, M., Karimi, F., Karimi-Maleh, H. & Razaghi, P. 2022. Production of bioethanol from carrot pulp in the presence of *Saccharomyces cerevisiae* and beet molasses inoculum. A biomass based investigation. *Chemosphere* 286: 131688. <https://doi.org/10.1016/j.chemosphere.2021.131688>
- Kim, D.-O., Padilla-Zakour, O.L. & Griffiths, P.D. 2004. Flavonoids and antioxidant capacity of various cabbage genotypes at juvenile stage. *Journal of Food Science* 69: 9. <https://doi.org/10.1111/j.1365-2621.2004.tb09916.x>
- Kohajdová, Z., Karovičová, J., Kuchtová, V. & Lauková, M. 2018. Utilisation of beetroot powder for bakery applications. *Chemistry Paper* 72: 1507–1515. <https://doi.org/10.1007/s11696-018-0392-z>
- Kop, A., Durmaz, Y. & Hekimoglu, M. 2010. Effect of natural pigment sources on colouration of cichlid (*Cichlasoma severum* sp. Heckel, 1840). *Journal of Animal and Veterinary Advances* 9: 566–569. <https://doi.org/10.3923/javaa.2010.566.569>
- Kujala, T.S., Loponen, J.M., Klika, K.D. & Pihlaja, K. 2000. Phenolics and betacyanins in red beetroot (*Beta vulgaris*) root: Distribution and effect of cold storage on the content of total phenolics and three individual compounds. *Journal of Agriculture and Food Chemistry* 48: 5338–5342. <https://doi.org/10.1021/jf000523q>
- Kujala, T., Loponen, J. & Pihlaja, K. 2001. Betalains and phenolics in red beetroot (*Beta vulgaris*) peel extracts: extraction and characterisation. *Zeitschrift für Naturforschung C* 56: 343–348. <https://doi.org/10.1515/znc-2001-5-604>
- Kumar, V., Kushwaha, R., Goyal, A., Tanwar, R. & Kaur, J. 2018. Process optimization for the preparation of antioxidant rich ginger candy using beetroot pomace extract. *Food Chemistry* 245: 168–177. <https://doi.org/10.1016/j.foodchem.2017.10.089>
- Kushwaha, A.K., Gupta, N. & Chattopadhyaya, M.C. 2014. Removal of cationic methylene blue and malachite green dyes from aqueous solution by waste materials of *Daucus carota*. *Journal of Saudi Chemical Society* 18: 200–207. <https://doi.org/10.1016/j.jscs.2011.06.011>
- Kushwaha, A.K., Kumar, V., Vyas, G. & Kaur, J. 2018. Optimization of different variable for eco-friendly extraction of betalains and phytochemicals from beetroot pomace. *Waste and biomass Valorization* 9: 1485–1494. <https://doi.org/10.1007/s12649-017-9953-6>
- Kusznierewicz, B., Bartoszek, A., Wolskaa, L., Drzewieckic, J., Gorinsteind, S. & Namieśnik, J. 2008. Partial characterization of white cabbages (*Brassica oleracea* var. *capitata* f. *alba*) from different regions by glucosinolates, bioactive compounds, total antioxidant activities and proteins. *LWT- Food Science and Technology* 41: 1–9. <https://doi.org/10.1016/j.lwt.2007.02.007>
- Laflamme, L.F. 1992. Carrot/grass silage as cattle feed. *Canadian Journal of Animal Science* 72: 441–443. <https://doi.org/10.4141/cjas92-056>
- Lai, W.T., Khong, N.M.H., Lim, S.S., Hee, Y.Y., Sim, B.I. & Lau, K.Y. 2017. A review: modified agricultural by-products for the development and fortification of food products and nutraceuticals. *Trends in Food Science & Technology* 59: 148–160. <https://doi.org/10.1016/j.tifs.2016.11.014>
- Ławińska, K., Szufa, S., Obraniak, A., Olejnik, T., Siuda, R., Kwiatek, J. & Ogrodowczyk, D. 2020. Disc granulation process of carbonation lime mud as a method of post-production waste management. *Energies* 13: 3419. <https://doi.org/10.3390/en13133419>
- Lazar (Mistrieanu), S., Constantin, O.E., Stanciuc, N., Aprodu, I., Croitoru, C. & Rapeanu, G. 2021. Optimization of betalain pigments extraction using beetroot by-products as a valuable source. *Inventions* 6: 50. <https://doi.org/10.3390/inventions6030050>
- Leite, C.W., Boroski, M., Boeing, J.S., Aguiar, A.C., França, P.B., Souza, Nilson, E. & Visentainer, J.V. 2011. Chemical characterization of leaves of organically grown carrot (*Daucus carota* L.) in various stages of development for use as food. *Ciência e Tecnologia de Alimentos* 31: 735–738. <https://doi.org/10.1590/S0101-20612011000300028>
- Liang, J.-L., Yeow, C.C., Teo, K.C., Gnanaraj, C. & Chang, Y.P. 2019. Valorizing cabbage (*Brassica oleracea* L. var. *capitata*) and capsicum (*Capsicum annuum* L.) wastes: *in vitro* health-promoting activities. *Journal of Food Science & Technology* 56: 4696–4704. <https://doi.org/10.1007/s13197-019-03912-5>
- Liang, J.L. 2016. Functional properties of wastes from cabbage (*Brassica oleracea* L. *capitata*) and capsicum (*Capsicum annuum* L. var. *annuum*). Bachelor of science biomedical science. Universiti Tunku Abdul Rahman p. 99.
- Lili, W., Sunarto, Santoso, B.J., Zidni, I. 2018. Carrot starch addition effect on artificial feed towards color enhancement of sword-tail fish (*Xiphophors Helleri*). *Global Science Journal* 6: 31–41.

- Liu, C., Masri, J., Perez, V., Maya, C. & Zhao, J. 2020. Growth performance and nutrient composition of mealworms (*Tenebrio molitor*) fed on fresh plant materials-supplemented diets. *Foods* 9: 151. <https://doi.org/10.3390/foods9020151>
- Ma, S., Ren, B., Diao, Z., Chen, Y., Qiao, Q. & Liu, X. 2016. Physicochemical properties and intestinal protective effect of ultra-micro ground insoluble dietary fibre from carrot pomace. *Food Function* 7: 392–399. <https://doi.org/10.1039/C6FO00665E>
- Madhumithah, C.G., Raghavan, K., Sridhar, S., Changam, S.S., Soma, G. & Kotturathu, M.C. 2011. Utilization of vegetable wastes for production of protease by solid state fermentation using *Aspergillus niger*. *World Journal of Agricultural Sciences* 7: 550–555.
- Mago, M., Gupta, R., Yadav, A. & Kumar G.V. 2022. Sustainable treatment and nutrient recovery from leafy waste through vermicomposting. *Bioresource Technology* 347: 126390. <https://doi.org/10.1016/j.biortech.2021.126390>
- Mahgoub, O., Kadim, I.T., Eltahir, Y., Al-Lawatia, S. & Al-Ismaili, A.M. 2018. Nutritional value of vegetable wastes as livestock feed. *Sultan Qaboos University Journal of Science [SQUJS]* 23: 78–84. <https://doi.org/10.24200/squjs.vol23iss2pp78-84>
- Mainali, B.P., Moadeli, T., Ponton, F., Taylor, P.W. & Neven, L. 2019. Comparison of gel larval diet with traditional lucerne chaff and carrot solid diets for rearing of Queensland fruit fly (Diptera: Tephritidae). *Journal of Economic Entomology* 112: 2278–2286. <https://doi.org/10.1093/jee/toz140>
- Majerska, J., Michalskab, A. & Figiel, A. 2019. A review of new directions in managing fruit and vegetable processing byproducts. *Trends in Food Science & Technology* 88: 207–219. <https://doi.org/10.1016/j.tifs.2019.03.021>
- Mannarmannan, M. & Biswas, K. 2021. Phytochemical-assisted Synthesis of cuprous oxide nanoparticles and their antimicrobial studies. *Chemistry Select (Weinheim)* 6: 3534–3539. <https://doi.org/10.1002/slct.202004471>
- Marcillo-Parra, V., Tupuna-Yerovi, D.S., Gonzalez, Z. & Ruales, J. 2021. Encapsulation of bioactive compounds from fruit and vegetable by-product for food application - A review. *Trends in Food Science & Technology* 116: 11–23. <https://doi.org/10.1016/j.tifs.2021.07.009>
- Micheletti Lorizola, I., Miyamoto, J.E., Figueireda Vieira, A.L., Rocchetti Sumere, B., Neves Bezerra, R.M., Torsoni, M.A., Torsoni, A.S., Rostagno, M.A., Milanski, M. & Dário Capitani, C. 2021. Beet (*Beta vulgaris* L.) stalk and leaf supplementation changes the glucose homeostasis and inflammatory markers in the liver of mice exposed to a high-fat diet. *Food Chemistry- Molecular Science* 2: 100018. <https://doi.org/10.1016/j.fochms.2021.100018>
- Mirzaei-Aghsaghali, A. & Maheri-Sis, N. 2008. Nutritive value of some agro-industrial by-products for ruminants - a review. *World Journal of Zoology* 3: 40–46.
- Misi, S.N. & Foster, C.F. 2002. Semi-continuous anaerobic co-digestion of agro-waste. *Environmental Technology* 23: 445–451. <https://doi.org/10.1080/09593332508618405>
- Moradi, R., Kashefialasl, M., Marandi, R., Salahi, E. & Moradidehaqi, S. 2021. Application of box-behnken design and response surface methodology of acid red 18 adsorption onto PAC (synthesized carrot waste) coated with Fe₃O₄ nanoparticles from aquatic solution: kinetic and isotherm studies. *Archive of Hygiene Science* 10: 30–48. <https://doi.org/10.52547/ArchHygSci.10.1.30>
- Morales-Ramos, J.A., Rojas, M.G., Dossey, A.T. & Berhow, M. 2020. Self-selection of food ingredients and agricultural by-products by the house cricket, *Acheta domesticus* (Orthoptera: Gryllidae): a holistic approach to develop optimized diets. *PLoS ONE* 15: e0227400. <https://doi.org/10.1371/journal.pone.0227400>
- Mu, H., Zhao, C., Zhao, Y., Li, Y., Hua, D., Zhang, X. & Xu, H. 2017. Enhanced methane production by semi-continuous mesophilic co-digestion of potato waste and cabbage waste: Performance and microbial characteristics analysis. *Bioresource Technology* 236: 68–76. <https://doi.org/10.1016/j.biortech.2017.03.138>
- Mukodiningsih, S., Achmadi, J., Wahyono, F., Taswanda, T.R., Sugiono & Ohh, S.J. 2019a. The effect of lactic acid bacteria-containing calf starter on Holstein calf performance. *IOP conference series. Earth and Environmental Science* 250: 12110. <https://doi.org/10.1088/1755-1315/250/1/012110>
- Mukodiningsih, S., Pujaningsih, R.I., Resniati, R. & Astuti, A. 2019b. The effect of storage time on the total lactic acid bacteria and presence of gram positive and negative bacteria in calf starter pellet added with fermented cabbage waste. *ISSTAP Buletin Peternakan* 43: 27–30. <https://doi.org/10.21059/buletinpeternakan.v43i1.38478>
- Mustafa, A.F. & Baurhoo, B. 2018. Evaluation of dried vegetable residues for poultry: III Effects of feeding cabbage leaf residues on laying performance, egg quality, and apparent total tract digestibility. *Journal of Applied Poultry Research* 27: 45–151. <https://doi.org/10.3382/japr/pfx031>
- Nalecz-Tarwacka, T., Karaszewska, A. & Zdziarski, K. 2003. The influence of carrot addition to cow's ration on the level of vitamins and fatty acids in cow milk. *Polish Journal of Food Nutrition Sciences* 12: 53–6.
- Nasernejad, B., Zadeh, T.E., Pour, B.B., Bygi, M.E. & Zamani, A. 2005. Comparison for biosorption modeling of heavy metals (Cr (III), Cu (II), Zn (II)) adsorption from wastewater by carrot residues. *Process Biochemistry* 40: 1319–1322. <https://doi.org/10.1016/j.procbio.2004.06.010>
- Naviglio, D., Scarano, P., Ciaravolo, M. & Gallo, M. 2019. Rapid solid-liquid dynamic extraction (RSLDE): A powerful and greener alternative to the latest solid-liquid extraction techniques. *Foods* 8: 245. <https://doi.org/10.3390/foods8070245>
- Ngu, N.T. & Ledin, I. 2005. Effects of feeding wastes from Brassica species on growth of goats and pesticide/insecticide residues in goat meat. *Asian-Australasian Journal of Animal Sciences* 18: 197–202. <https://doi.org/10.5713/ajas.2005.197>
- Nguen, H.T., Vo Thanh, T., Vo, L., Bui Phan, T.H. & Preston, T.R. 2009. Effects on growth of rabbits of supplementing a basal diet of water spinach (*Ipomoea aquatica*) with vegetable wastes and paddy rice. *Livestock Research for Rural Development* 21: 174.
- Nilnakara, S., Chiewchan, N. & Devahastin, S. 2009. Production of antioxidant dietary fibre powder from cabbage outer leaves. *Food and Bioproducts Processing* 87: 301–307. <https://doi.org/10.1016/j.fbp.2008.12.004>
- Nkosi, B.D. & Ratsaka, M.M. 2010. Effect of dietary inclusion of discarded beetroot and potato hash silage on growth performance and digestibility in South African dorper lambs. *Journal of Animal and Veterinary Advances (JAVA)* 9: 853–856. <https://doi.org/10.3923/javaa.2010.853.856>

- Ntsohi, N., Fanadzo, M., Le Roes-Hill, M. & Nchu, F. 2021. Effects of *Clonostachys rosea* f. *catenula* inoculum on the composting of cabbage wastes and the endophytic activities of the composted material on tomatoes and red spider mite infestation. *Microorganisms* 9: 1184. <https://doi.org/10.3390/microorganisms9061184>
- Nutter, J., Fernandez, M.V., Jagus, M.R. & Agüero, M.V. 2021. Development of an aqueous ultrasound-assisted extraction process of bioactive compounds from beet leaves: A proposal for reducing losses and increasing biomass utilization. *Journal of Science of Food and Agriculture* 101: 1989–1997. <https://doi.org/10.1002/jsfa.10815>
- Nynäs, A.-L., Newson, W.R. & Johansson, E. 2021. Protein fractionation of green leaves as an underutilized food source - Protein yield and the effect of process parameters. *Foods* 10: 2533. <https://doi.org/10.3390/foods10112533>
- Omolara, B.O. & Olaleye, A.A. 2010. Performance, carcass analysis and sensory evaluation of cooked meat of snaillets of African Giant Land snail (*Archachatina marginata*) fed pawpaw leaves, whole lettuce, lettuce waste and cabbage waste as sole feed ingredient. *African Journal of Agricultural Research* 5: 2386–2391.
- de Oliveira, A.N., Stevanato, F.B. & Visentainer, J.V. 2015. Food supplementation for workers: flour enriched with omega-3. *Acta Scientiarum-Technology* 37: 133–139. <https://doi.org/10.4025/actascitechnol.v37i1.20108>
- Olsson, M. 2023. Livsmedelsförluster vid produktion av morötter. Jordbruksverket, Sweden. Rapport 4. (in Swedish). <https://webbutik.jordbruksverket.se/sv/artiklar/ra234.html>
- Ooninx, D.G.A.B. & de Boer, I.J.M. 2012. Environmental impact of the production of mealworms as a protein source for humans - A Life cycle assessment. *PLoS One* 7: e51145. <https://doi.org/10.1371/journal.pone.0051145>
- Ooninx, D.G.A.B., van Broekhoven, S., van Huis, A. & van Loon, J.J.A. 2015. Feed conversion, survival and development, and composition of four insect species on diets composed of food by-products. *PLoS One* 10: e0144601. <https://doi.org/10.1371/journal.pone.0144601>
- Otoni, C.G., Lodi, B.D., Lorevice, M.V., Leitão, R.C., Ferreira, M.D., de Moura, M.R. & Mattoso, L.H.C. 2018. Optimized and scaled-up production of cellulose-reinforced biodegradable composite films made up of carrot processing waste. *Industrial Crops and Products* 21: 66–72. <https://doi.org/10.1016/j.indcrop.2018.05.003>
- Patras, A. 2019. Stability and colour evaluation of red cabbage waste hydroethanolic extract in presence of different food additives or ingredients. *Food Chemistry* 275: 539–548. <https://doi.org/10.1016/j.foodchem.2018.09.100>
- Van Peer, M., Frooninx, L., Coudron, C., Berrens, S., Alvarez, C., Derutter, D., Verheyen, G. & Van Miert, S. 2021. Valorisation Potential of Using Organic Side Streams as Feed for *Tenebrio molitor*, *Acheta domesticus* and *Locusta migratoria*. *Insects* 12: 796. <https://doi.org/10.3390/insects12090796>
- Perotto, G., Ceseracciu, L., Simonutti, R., Paul, U.P., Guzman-Puyol, S., Tran, T-N., Bayer, I.S. & Athanassiou, A. 2018. Bioplastics from vegetable waste via an eco-friendly water-based process. *International Journal of Green Chemistry Resources (GC)* 20: 894–902. <https://doi.org/10.1039/C7GC03368K>
- Petrović, M., Veljović, S., Tomić, N., Zlatanović, S., Tosti, T., Vukosavljević, P. & Gorjanović, S. 2021. Formulation of novel liqueurs from juice industry waste: consumer acceptance, phenolic profile and preliminary monitoring of antioxidant activity and colour changes during storage. *Food Technology and Biotechnology* 59: 282–294. <https://doi.org/10.17113/ftb.59.03.21.6759>
- Pinotti, L. & Ottoboni, M. 2021. Substrate as insect feed for bio-mass production *Journal of Insects as Food and Feed* 7: 585–596. <https://doi.org/10.3920/JIFF2020.01110>
- Pirie, A. & Wood, C. 1946. Effects of vitamin A deficiency in the rabbit. *Biochemical Journal* 40: 557–60. <https://doi.org/10.1042/bj0400557>
- Plazzotta, S., Manzocco, L. & Nicolini, M.C. 2020. Fruit and vegetable waste management and the challenge of fresh-cut salad. *Trends in Food Science & Technology* 63: 51–59. <https://doi.org/10.1016/j.tifs.2017.02.013>
- Popescu, V., Blaga, A.C., Pruneanu, M., Cristian, I.N., Pişlaru, M., Popescu, A., Rotaru, V., Creţescu, I. & Caşcaval, D. 2021. Green chemistry in the extraction of natural dyes from colored food waste, for dyeing protein textile materials. *Polymers* 13: 3867. <https://doi.org/10.3390/polym13223867>
- Porto Dalla Costa, A., Stahl Hermes, V. & de Oliveira Rios, A. 2017. Minimally processed beetroot waste as an alternative source to obtain functional ingredients. *Journal of Food Science & Technology* 54: 2050–2058. <https://doi.org/10.1007/s13197-017-2642-4>
- Pradhan, S., Abdelaal, A.H., Mroue, K., Al Ansari, T., Mackey, H.R. & McKay, G. 2020a. Biochar from vegetable wastes: agro environmental characterization. *Biochar* 2: 439–453. <https://doi.org/10.1007/s42773-020-00069-9>
- Pradhan, S., Shahbaz, M., Abdelaal, A., Al-Ansari, T., Mackey, H.R. & McKay, G. 2020b. Optimization of process and properties of biochar from cabbage waste by response surface methodology. *Biomass Conversion and Biorefinery* 12: 5479–5491. <https://doi.org/10.1007/s13399-020-01101-5>
- Priyadarshini, M., Subash, A., Shobana, A. & Sujatha, A. 2021. Potential use of Brassica oleracea L. (purple cabbage) waste as a valuable source of natural colorant for the textile industry. *Journal of Advanced Scientific Research* 12: 183–189. <https://doi.org/10.55218/JASR.202112123>
- Purohit, A.J. & Gogate, P.R. 2015. Ultrasound-assisted extraction of β -carotene from waste carrot residue: effect of operating parameters and type of ultrasonic irradiation. *Separation Science and Technology* 50: 1507–1517. <https://doi.org/10.1080/01496395.2014.978472>
- Ralla, T., Salminen, H., Wolfangel, T., Edelmann, M., Dawid, C., Hofmann, T. & Weiss, J. 2019. Value addition of red beet (*Beta vulgaris* L.) by-products: Emulsion formation and stability. *International Journal of Food Science & Technology* 54: 619–625. <https://doi.org/10.1111/ijfs.13886>
- Ramakodi, M.P., Santhosh, N., Pragadeesh, T., Mohan, S. V. & Basha, S. 2020. Production of protease enzyme from slaughter-house effluent: An approach to generate value-added products from waste. *Bioresource Technology Reports* 12: 100552. <https://doi.org/10.1016/j.biteb.2020.100552>

- Ramos-Andrés, M., Aguilera-Torre, B. & García-Serna, J. 2021. Biorefinery of discarded carrot juice to produce carotenoids and fermentation products. *Journal of Cleaner Production* 323: 129139. <https://doi.org/10.1016/j.jclepro.2021.129139>
- Rao, M., Bast, A. & de Boer, A. 2021. Valorized food processing by-products in the EU: Finding the balance between safety, nutrition, and sustainability. *Sustainability* 13: 4428. <https://doi.org/10.3390/su13084428>
- Râpă, M., Darie-Niță, R.N. & Coman, G. 2024. Valorization of fruit and vegetable waste into sustainable and value-added materials. *Waste* 2: 258–278. <https://doi.org/10.3390/waste2030015>
- Rashad, M.M. & Abdou, H.M. 2001. Fungal degradation of cabbage leaf liquid waste. *Advances Food Science* 23: 124–130.
- Ratti, C. 2001. Hot air and freeze-drying of high-value foods: A review. *Journal of Food Engineering* 49: 311–319. [https://doi.org/10.1016/S0260-8774\(00\)00228-4](https://doi.org/10.1016/S0260-8774(00)00228-4)
- Ravi, P.P., Lindner, J., Oechsner, H. & Lemmer, A. 2018. Effects of target pH-value on organic acids and methane production in two-stage anaerobic digestion of vegetable waste. *Bioresource Technology* 247: 96–102. <https://doi.org/10.1016/j.biortech.2017.09.068>
- Razzaq, Z.U., Maan, A.A., Nazir, A., Hafeez, M.A. & Khan, M.K.I. 2022. Characterizing the single cell protein enriched noodles for nutritional and organoleptic attributes. *Journal of Food Measurement and Characterization* 16: 1725–1732. <https://doi.org/10.1007/s11694-022-01300-w>
- Ren, H., Wang, L., Sun, Y., Zhao, Q., Sun, Y., Li, J. & Zhang, B. 2021. Enhancing the co-ensiling performance of corn stover and cabbage waste via the addition of cellulase. *Bioresources* 16: 6342–6362. <https://doi.org/10.15376/biores.16.3.6342-6362>
- de Rezende, A.V., Silveira Rabelo, C.H., da Silva, M.R.M., Härter, C.J. & Veiga, R.M. 2015. Wasted cabbage (*Brassica oleracea*) silages treated with different levels of ground corn and silage inoculant. *Revista Brasileira de Zootecnia* 44: 296–302. <https://doi.org/10.1590/S1806-92902015000800005>
- Ribeiro, T.B., Voss, G.B., Coelho, M.C. & Pintado, M.E. 2022. Food waste and by-product valorization as an integrated approach with zero waste: Future challenges. In: *Future foods*. Chapter 33. Elsevier Inc. p. 569–596. <https://doi.org/10.1016/B978-0-323-91001-9.00017-7>
- Rima, J., Rahme, K. & Assaker, K. 2014. The use of modified beetroot fibers by sodium dodecyl sulfate (SDS) cleaning water contaminated by organic and inorganic compounds. *Journal of Food Resources* 3: 19. <https://doi.org/10.5539/jfr.v3n5p19>
- Rodríguez-Félix, F., Corte-Tarazón, J.A, Rochín-Wong, S., Fernández-Quiroz, J.D., Garzón-García, A.M., Santos-Sauceda, I., Plascencia-Martínez, D.F., Chan-Chan, L.H., Vásquez-López, C., Barreras-Urbina, C.G., Olguin-Moreno, A. & Tapia-Hernández, J.A. 2022. Physicochemical, structural, mechanical and antioxidant properties of zein films incorporated with no-ultrafiltered and ultrafiltered betalains extract from the beetroot (*Beta vulgaris*) bagasse with potential application as active food packaging. *Journal of Food Engineering* 334: 111153. <https://doi.org/10.1016/j.jfoodeng.2022.111153>
- Roohinejad, S., Oey, I., Everett, D.W., Niven, B.E. 2014. Evaluating the effectiveness of β -carotene extraction from pulsed electric field-treated carrot pomace using oil-in-water microemulsion. *Food and Bioprocess Technology* 7: 3336–3348. <https://doi.org/10.1007/s11947-014-1334-6>
- Roshan-Bakhsh, A., Pourjam, E., Ayyari, M. & Pedram, M. 2019. Biocontrol properties of some agricultural waste extracts on three nematode species in in vitro and in vivo conditions. *Nematology: International Journal of Fundamental and Applied Nematological Research* 21: 837–846. <https://doi.org/10.1163/15685411-00003258>
- Rovai, D., Ortgies, M., Amin, S., Kuwahara, S., Schwartz, G., Lesniauskas, R., Garza, J. & Lammert, A. 2022. Utilization of carrot pomace to grow mealworm larvae (*Tenebrio molitor*). *Sustainability* 13: 9341. <https://doi.org/10.3390/su13169341>
- Rust, S. & Buskirk, D. 2008. Feeding carrots or sugar beets to cattle. *Cattle Call*, Michigan State University 13: 1–6.
- Sagar, N.A., Pareek, S., Sharma, S., Yahia, E.M. & Lobo, M.G. 2018. Fruit and vegetable waste: bioactive compounds, their extraction, and possible utilization. *Comprehensive Reviews in Food Science and Food Safety* 17: 512–531. <https://doi.org/10.1111/1541-4337.12330>
- Salamatullah, A.M., Hayat, K., Alkaltham, M.S., Ahmed, M.A., Arzoo, S., Husain, F.M., Al-Dossari, A.M., Shamlan, G. & Al-Harbi, L.N. 2021. Bioactive and antimicrobial properties of oven-dried beetroot (pulp and peel) using different solvents. *Processes* 9: 588. <https://doi.org/10.3390/pr9040588>
- Saleem, A., Irshad, M., Eneji, A.E., Hassan, A., Mahmood, Q. & Irshad, U. 2017. Fractionation of phosphorus in soils amended with poultry manure co-composted with sugarcane and cabbage wastes. *Journal of Biosciences* 33: 1230–1241. <https://doi.org/10.14393/BJ-v33n5a2017-37179>
- Saleem, A., Fareed, I., Irshad, M., Mahmood, Q., Eneji, A. E. & Shahzad, M. 2018. Transformations of phosphorus and other plant nutrients in poultry litter composted with sugarcane and cabbage wastes. *Compost Science and Utilization* 26: 114–127. <https://doi.org/10.1080/1065657X.2017.1396268>
- Salvañal, L., Clementz, A., Guerra, L., Yori, J.C. & Romanini, D. 2021. L-lactic acid production using the syrup obtained in biorefinery of carrot discards. *Food and Bioproducts Processing* 127: 465–471. <https://doi.org/10.1016/j.fbp.2021.04.002>
- Šamec, D., Piljac-Zegarac, J., Bogović, M., Habjanič, K. & Grúz, J. 2011. Antioxidant potency of white (*Brassica oleracea* L. var. capitata) and Chinese (*Brassica rapa* L. var. *pekinensis* (Lour.)) cabbage: The influence of development stage, cultivar choice and seed selection. *Scientia Horticulturae* 128: 75–83. <https://doi.org/10.1016/j.scienta.2011.01.009>
- San Martín, D., Ramos, S. & Zufáet, J. 2016. Valorisation of food waste to produce new raw materials for animal feed. *Food Chemistry* 198: 68–74. <https://doi.org/10.1016/j.foodchem.2015.11.035>
- Sedlar, T., Čakarević, J., Tomić, J. & Popovic, L. 2021. Vegetable by-products as new sources of functional proteins. *Plant Foods for Human Nutrition* 76: 31–36. <https://doi.org/10.1007/s11130-020-00870-8>
- Selder, L., Turunen, R., Vuoristo, K.S., Uusitalo, J. & Zeng, A.-P. 2021. Volatile and medium chain fatty acid production from cabbage residues using mixed culture fermentations and in-situ product removal by electrodialysis. *Bioresource Technology Reports* 36: 100828. <https://doi.org/10.1016/j.biteb.2021.100828>

- Sepulveda, L. Contreras, E., Cerro, D. & Quintulen, L. 2021. Technical feasibility of natural antioxidant recovery from the mixture of the inedible fractions of vegetables produced in wholesale market. *CYTA-Journal of Food* 19: 418–428. <https://doi.org/10.1080/19476337.2021.1915878>
- Šeregelj, V., Škrobot, D., Kojić, J., Pezo, L., Šovljanski, O., Tumbas, Š.V., Vulić, J., Hidalgo, A., Brandolini, A., Čanadanović-Brunet, J. & Četković, G. 2022. Quality and sensory profile of durum wheat pasta enriched with carrot waste encapsulates. *Foods* 11: 1130. <https://doi.org/10.3390/foods11081130>
- Šeremet, D., Durgo, K., Jokić, S., Huđek, A., Vojvodić C.A., Mandura, A., Jurasović, J. & Komes, D. 2020. Valorization of banana and red beetroot peels: determination of basic macrocomponent composition, application of novel extraction methodology and assessment of biological activity *in vitro*. *Sustainability* 12: 4539. <https://doi.org/10.3390/su12114539>
- Setya Utama, C., Suthama, N., Sulistiyanto, B. & Setiani, B.E. 2013. Utility of rice bran mixed with fermentation extract of vegetable waste unconditioned as probiotics from vegetable market. *International Journal of Science and Engineering* 4: 97–102. <https://doi.org/10.12777/ijse.4.2.97-102>
- Sharma, K.D., Karki, S., Thakur, N.S. & Attri, S. 2012. Chemical composition, functional properties and processing of carrot - A review. *Journal of Food Science and Technology* 49: 22–32. <https://doi.org/10.1007/s13197-011-0310-7>
- Sidker, A.C., Chowdhury, S.D., Rashid, M.H., Sarker, A.K. & Das, S.C. 1998. Use of dried carrot meal (DCM) in lauing hen diet for egg yolk Pigmentation. *AJAS* 11: 239–244. <https://doi.org/10.5713/ajas.1998.239>
- Singh, B., Panesar, P.S. & Nanda, V. 2006. Utilization of carrot pomace for the preparation of a value added product. *World Journal of Dairy & Food Sciences* 1: 22–27.
- Singh, B.K., Sharma, S.R., Kalia, P. & Singh, B. 2010. Character association and path analysis of morphological and economic traits in cabbage (*Brassica oleracea* var. *capitata*). *The Indian Journal of Agricultural Sciences* 80: 116–118.
- Singh, R.S., Chauhan, K., Singh, J., Pandey, A. & Larroche, C. 2018. Solid-state fermentation of carrot pomace for the production of inulinase by *Penicillium oxalicum* BGPUP-4. *Food Technology and Biotechnology* 56: 31–39.
- Socas-Rodriguez, B., Alvarez-Rivera, G., Valdes, A., Ibanez, E. & Cifuentes, A. 2021. Food by-products and food wastes: are they safe enough for their valorization? *Trends in Food Science & Technology* 114: 133–147. <https://doi.org/10.1016/j.tifs.2021.05.002>
- Sogut, E. & Cakmak, H. 2020. Utilization of carrot (*Daucus carota* L.) fiber as a filler for chitosan based films. *Food Hydrocolloids* 106: 105861. <https://doi.org/10.1016/j.foodhyd.2020.105861>
- Solowski, G., Konkol, I., Hrycak, B. & Czynkowski, D. 2019. Hydrogen and methane production under conditions of anaerobic digestion of key-lime and cabbage waster. *Agritech* 39: 243–250. <https://doi.org/10.22146/agritech.35848>
- Steenfeldt, S., Kjaer, J.B. & Engberg, R.M. 2007. Effect of feeding silages or carrots as supplements to laying hens on production performance, nutrient digestibility, gut structure, gut microflora and feather pecking behaviour. *British Poultry Science* 48: 454–68. <https://doi.org/10.1080/00071660701473857>
- Stockhammer, S., Stolze, K., Rohr-Udilova, N., Chizzola, R., Zitterl-Eglseer, K. & Franz, C. 2009. Antioxidant activity of phyto-genous industrial waste and derived extracts for the production of feed and food additives. *International Journal of Food Science & Technology* 44: 702–710. <https://doi.org/10.1111/j.1365-2621.2008.01829.x>
- Supyani, Hadiwiyono, Poromarto, S.H, Widono, S., Septariani, D.N. & Putri, C.P. 2021. Effectiveness of fermentation extract of cabbage leaves from harvesting waste to control purple blotch of garlic. *IOP conference series. Earth and Environmental Science* 828: 12020. <https://doi.org/10.1088/1755-1315/828/1/012020>
- Surendran, P.A.C. & Shanmugam, P. 2021. Correlation between empirical formulae based stoichiometric and experimental methane potential and calorific energy values for vegetable solid wastes. *Energy Reports* 7: 19–31. <https://doi.org/10.1016/j.egyr.2020.10.071>
- Suzuki, T., Usui, I., Tomita Y.K., Kono, S., Tsubura, H., Miki, Y. & Hasegawa, K. 2001. Effects of acid extracts of tomato (*Lycopersicon esculentum* Mill.) and carrot (*Daucus carota* L.) wastes from the food industry on the growth of some crops and weeds. *Weed Biology and Management* 1: 226–230. <https://doi.org/10.1046/j.1445-6664.2001.00036.x>
- Syahrizal, Ediwarman, Safratilofo & Ridwan, M. 2022. Analysis of the use of media resulting from bioconversion of organic waste in the production of maggots BSF (black soldier fly). *Jurnal Akuakultur Indonesia* 21: 1–10. <https://doi.org/10.19027/jai.21.1.1-10>
- Szymańska-Chargot, M., Chylińska, M., Pieczywek, P.M. & Zdunek, A. 2019. Tailored nanocellulose structure depending on the origin. Example of apple parenchyma and carrot root celluloses. *Carbohydrate Polymers* 210: 186–195. <https://doi.org/10.1016/j.carbpol.2019.01.070>
- Tran, T.N., Athanassiou, A., Basit, A. & Bayer, I.S. 2017. Starch-based bio-elastomers functionalized with red beetroot natural antioxidant. *Food Chemistry* 216: 324–333. <https://doi.org/10.1016/j.foodchem.2016.08.055>
- Tsutsui, Y., Shinihara, T. & Hachinohe, Y. 1967. Some physiological aspects in body weight in rabbits. *Journal of Faculty of Agriculture, Hokkaido University* 55: 111–132.
- Tuck, C.O., Perez, E., Horvaeth, I.T., Sheldon, R.A. & Polakoff, M. 2012. Valorization of biomass: deriving more value from waste. *Science* 10: 695–699. <https://doi.org/10.1126/science.1218930>
- Tumbas Šaponjac, V., Čanadanović-Brunet, J., Četković, G., Jakišić, M., Djilas, S., Vulić, J. & Stajčić, S. 2016. Encapsulation of beetroot pomace extract: RSM optimization, storage and gastrointestinal stability. *Molecules* 21: 584–584. <https://doi.org/10.3390/molecules21050584>
- UH (University of Helsinki) 2025. Helka Guide. https://libraryguides.helsinki.fi/helka_en
- Vakili, F., Rashidi, A., Taghavi, L. & Mansouri, N. 2021a. Conversion of biomass to N, S co-doped porous graphene as an adsorbent for mercury vapor removal: optimization and DFT study. *Journal of Environmental Health Science and Engineering* 19: 1569–1582. <https://doi.org/10.1007/s40201-021-00712-y>
- Vakili, F., Rashidi, A., Taghavi, L. & Mansouri, N. 2021b. Single-step synthesis of N, S co-doped waste-derived nanoporous carbon sorbent for mercury vapor removal. *Environmental Science and Pollution Reseach* 28: 17265–17274. <https://doi.org/10.1007/s11356-020-12075-2>

- Vandenbergh, A., Markó, I.E., Lucaccioni, F. & Lutts, S. 2013. Enantioselective hydrolysis of racemic 1-phenylethyl acetate by an enzymatic system from fresh vegetables. *Industrial Crops and Products* 42: 380–385. <https://doi.org/10.1016/j.indcrop.2012.06.003>
- Varelas, V. 2019. Food wastes as a potential new source for edible insect mass production for food and feed: A review. *Fermentation* 5: 81. <https://doi.org/10.3390/fermentation5030081>
- Vulić, J.J., Cebović, T.N., Canadanović, V.M., Cetković, G.S., Djilas, S.M., Canadanović-Brunet, J.M., Velicanski, A.S., Cvetkovci, D.D. & Tumbas, V.T. 2013. Antiradical, antimicrobial and cytotoxic activities of commercial beetroot pomace. *Food and Function* 4: 713–721. <https://doi.org/10.1039/c3fo30315b>
- Vulić, J.J., Cebović, T.N., Canadanović-Brunet, J.M., Cetković, G.S., Čanadanović, V.M., Djilas, S.M. & Tumbas Šaponjac, V.T. 2014. *In vivo* and *in vitro* antioxidant effects of beetroot pomace extracts. *Journal of Functional Foods* 6: 168–175. <https://doi.org/10.1016/j.jff.2013.10.003>
- Wadhwa, M., Kaushal, S. & Bakshi, M.P.S. 2006. Nutritive evaluation of vegetable wastes as complete feed for goat bucks. *Small Ruminant Research* 64: 279–284. <https://doi.org/10.1016/j.smallrumres.2005.05.017>
- Wagde, M.S., Sharma, S.K. & Sharma, B.K. 2018. Effect of natural β -carotene sources-carrot (*Daucus carota*) and spinach (*Spinacia oleracea*) on the growth of an ornamental fish - sword tail (*Xiphophorus hellerii*). *Journal of Entomology and Zoology Studies* 6: 2112–2115.
- Walker, R.L., Edwards, A.C., Maskell, P., Watson, C.A., Rees, R.M., Knox, O.G.G. & Stockdale, E.A. 2012. The effect of co-composted cabbage and ground phosphate rock on the early growth and P uptake of oilseed rape and perennial ryegrass. *Journal of Plant Nutrition and Soil Science* 175: 595–603. <https://doi.org/10.1002/jpln.201100124>
- Wassef, E.A., Chatzifotis, S., Sakr, E.M., Norhan E. & Saleh, N.E. 2010. Effect of two natural carotenoid sources in diets for gilt-head seabream, *Sparus aurata*, on growth and skin coloration. *Journal of Applied Aquaculture* 22: 216–229. <https://doi.org/10.1080/10454438.2010.497741>
- Weerakkody, W.S. & Cumaratunga, P.R.T. 2016. Effects of dried, powdered carrot (*Daucus carota*) incorporated diet on the skin and flesh colouration of *Catla catla*. *Sri Lanka Journal of Aquatic Sciences* 21: 95–103. <https://doi.org/10.4038/sljias.v21i2.7505>
- Wei, D., Liu, J., Deng, C., Wang, C., Zhao, X., Kai, W., Fang, Y., Yang, B. & Wudi, Z. 2021. Chlorophyll changes during anaerobic digestion of cabbage waste. *Energy Sources, Part A: Recovery, Utilization, and Environmental Effects* 47: 1–14. <https://doi.org/10.1080/15567036.2021.2009063>
- Wekoye, J.N., Wanyonyi, W.C., Wangila, P.T. & Tonui, M.K. 2020. Kinetic and equilibrium studies of Congo red dye adsorption on cabbage waste powder. *Environmental Chemistry and Ecotoxicology* 2: 24–31. <https://doi.org/10.1016/j.enceco.2020.01.004>
- Wennberg, M., Enqvist, G. & Nyman, E. 2003. Effect of boiling on dietary fiber components in fresh and stored white cabbage (*Brassica oleracea* var. *capitata*). *Journal of Food Science* 68: 1615–1621. <https://doi.org/10.1111/j.1365-2621.2003.tb12301.x>
- Wolna-Maruwka, A., Dach, J., Cáceres, R., Czekala, W., Niewiadomska, A., Janczak, D. & Budka, A. 2019. An effective method of utilizing vegetable waste in the form of carriers for *Trichoderma* strains with phytosanitary properties. *Science of the Total Environment* 671: 795–804. <https://doi.org/10.1016/j.scitotenv.2019.03.120>
- Wu, W., Chen, Y., Faisal, S., Khan, A., Chen, Z., Ling, Z., Liu, P. & Li, X. 2016. Improving methane production in cow dung and corn straw co-fermentation systems via enhanced degradation of cellulose by cabbage addition. *Scientific Reports* 6: 33628–33628. <https://doi.org/10.1038/srep33628>
- Xue, S., Zhang, X., Ngo, H.H., Guo, W., Wen, H., Li, C., Zhang, Y. & Ma, C. 2019. Food waste based biochars for ammonia nitrogen removal from aqueous solutions. *Bioresource Technology* 292: 121927–121927. <https://doi.org/10.1016/j.biortech.2019.121927>
- Yadav, S., Arora, S.K. & Vats, S. 2023. Vitamins and minerals: A review on processing losses and strategies to control it. *Modern Concepts & Developments in Agronomy MCDA* 12: 000783. <https://doi.org/10.31031/MCDA.2023.12.000783>
- Yang, L. & Cosolini, S.I. 2019. A case study on converting organic farm waste vegetables to biogas using a cartridge design anaerobic digester. *Applied Biochemistry and Biotechnology* 189: 638–646. <https://doi.org/10.1007/s12010-019-03034-w>
- Yu, C.-Y., Jiang, B.-H. & Duan, K.-J. 2013. Production of bioethanol from carrot pomace using the thermotolerant yeast *Kluyveromyces marxianus*. *Energies* 6: 1794–1801. <https://doi.org/10.3390/en6031794>
- Zanoni, F., Primiterra, M., Angeli, N. & Zoccatelli, G. 2020. Microencapsulation by spray-drying of polyphenols extracted from red chicory and red cabbage: Effects on stability and color properties. *Food Chemistry* 307: 125535–125535. <https://doi.org/10.1016/j.foodchem.2019.125535>
- Zhang, Y., Cheng, X., Wang, Z., Tahir, M.H., Wang, Z., Wang, X. & Wang, C. 2022. Full recycling of high-value resources from cabbage waste by multi-stage utilization. *Science of the Total Environment* 804: 149951–149951. <https://doi.org/10.1016/j.scitotenv.2021.149951>
- Zhang, W., Liu, Q., Guo, L., Wang, P., Liu, S., Cjen, J. & Lei, Z. 2021. White cabbage (*Brassica oleracea* L.) waste, as biowaste for the preparation of novel superabsorbent polymer gel. *Journal of Environmental Chemistry* 9: 106689. <https://doi.org/10.1016/j.jece.2021.106689>
- Zin, M.M. & Bánvölgyi, S. 2021. Portfolio of beetroot (*Beta vulgaris* L.) peel extracts concentrated by nanofiltration membrane. *International Journal of Food Studies* 10: 334–345. <https://doi.org/10.7455/ijfs/10.2.2021.a5>
- Zin, M.M. & Bánvölgyi, S. 2022. Thermal emerging technology approach for the extraction of bioactive compounds from *Cylindra* beetroot (peel, flesh, and stalk) with green solvent. *Journal of Food Processing and Preservation*. <https://doi.org/10.1111/jfpp.16436>