



Antimicrobial Efficacy of Four Different Extracts of *Plantago major*: An *In vitro* Study

G. Akkuş^a and M. M. HIZ-ÇİÇEKLIYURT^{b*}

^a Department of Biology, Faculty of Arts & Science, Çanakkale Onsekiz Mart University, Turkey.
^b Department of Medical Biology, Faculty of Medicine, Çanakkale Onsekiz Mart University, Turkey.

Authors' contributions

This work was carried out in collaboration between both authors. Author GA is a student responsible for all antimicrobial assays and performed the statistical analysis and wrote the first draft of the manuscript. Author MMÇ wrote the protocol, managed the analyses of the study and the literature searches. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85117>

Original Research Article

Received 27 October 2021
Accepted 29 December 2021
Published 30 December 2021

ABSTRACT

Aims: *Plantago major* is frequently used in traditional treatment for upper respiratory tract diseases such as pneumonia, cough, pharyngitis, and skin, eye, and urinary tract infections. This study aims to evaluate the antimicrobial activity and minimum effective dose of hexane, methanol, ethanol, and water extract of *Plantago major* leaves.

Study design: *In vitro* experimental study.

Methodology: *P. major* leaves was crushed into a fine powder and dissolved in different solvents (hexane, methanol, ethanol, or water) using a Soxhlet device, then the extracts were purified by evaporation of the solvent. All extracts were analyzed for antibacterial and antifungal properties using broth dilution method depending on MIC value determined according to the solvent-microorganism-time trio in DDM.

Results: The *in vivo* test showed that all methods to extract of *Plantago major* leaves have activity against all test microorganisms. Both hexane and water extract showed the same activity on *Staphylococcus aureus*, *Bacillus subtilis*, and *Pseudomonas aeruginosa*, *C. albicans* and *Candida tropicalis* at 4 mg/ml. The lowest activity of *Plantago major*'s (PM's) hexane and water extract was on *Escherichia coli* and *Pseudomonas vulgaris* bacteria as 8 mg/ml. Methanol and ethanol extracts of PM showed higher activity than hexane and water extract. PM ethanol extract showed antimicrobial activity against *Staphylococcus aureus*, *Enterococcus faecalis*, *Pseudomonas aeruginosa* at 2ml/mL, and *Bacillus subtilis* *Escherichia coli* and *Proteus vulgaris* at 4 ml/mL.

Conclusion: The results nominate the *Plantago major* extract has potential antimicrobials and antifungals. However, the development of antimicrobial agents requires purifying the active bio components by high throughput techniques to achieve effective activity as positive controls.

Keywords: *Plantago major*; antimicrobial activity; MIC.

ABBREVIATIONS

Here is the Definitions section. This is an optional section.

T_B : Boiling point
 PM : *Plantago Major*

1. INTRODUCTION

Plantago major (*P. major*) is a widely used medicinal plant in folk medicine [1]. The plant contains bioactive components such as flavonoids, polysaccharides, terpenoids, lipids, iridoid glycosides, and caffeic acid derivatives [2,3]. Due to its rich components, the plant was used to treat various medical conditions such as coughs, infection, fever, bleeding, and inflammation [3,4]. Currently, animal studies focus on using the *Plantago major* (*PM*) on different medicinal conditions. Parhizgar and colleagues have demonstrated the protective effect of *PM* extract in the presence of kidney damage. Parhizgar has shown glomerular filtration rate (GFR), urine osmolarity, and urinary excretion rate of potassium were increased via the treatment of *PM* extract of kidney tissue-damaged rats [5]. In another study, hydroalcoholic extract of *Plantago major* also had a protective role in doxorubicin-induced nephropathy. These two animal studies have been demonstrated the protective role of *PM* extract on the renal system. Boskabadi et al. [6] have been shown potent relaxant effects of *PM* extract on Tracheal Smooth Muscles of rats. The randomized, double-blind placebo-controlled clinical trial done by Jazayeri and colleagues has achieved elevated liver enzymes and a better prognosis for Nonalcoholic Fatty Liver Disease by using 2 gr of *PM* supplementation twice daily for 12 weeks [7]. Depending on the therapeutic properties shown in both animals and clinical studies, *PM* extract protects the kidneys and liver against toxicity.

In another way of view, *PM* extract is not harmful to animals and humans by its protective role, thus potent bioactive compound for drug development. *PM* extracts were widely used for

antibiotics, antioxidants, analgesics, and wound healer purposes from ancient times. The antibacterial activity of *PM* extract was evaluated by Sharma et al. They found no activity for periodontal pathogens by the Kirby-Bauer disc diffusion technique [8]. However, none of the studies were performed to evaluate *PM* extract's antimicrobial activity by MIC method against standard test microorganisms. Thus, this study investigated the bioactivity of a *Plantago major* against pathogens: *Staphylococcus aureus*, *Enterococcus faecalis*, *Bacillus subtilis*, *Pseudomonas aeruginosa*, *Proteus vulgaris* as antimicrobial, and *Candida albicans* and *Candida tropicalis* as antifungal.

2. MATERIALS AND METHODS

2.1 Chemicals

The medium used for this study was Nutrient Broth for bacterial, and RPMI-1640 medium with L-glutamine for fungi were purchased from Sigma-Aldrich (Hamburg, Germany). The microbial lines were purchased from American Type Culture Collection (ATCC®, Manassas, VA, USA). The antibiotics and antifungal pharmaceuticals were used as a positive control as follows: Ampicillin (Mustafa Nevzat Pharmaceuticals, Turkey); Gentamicin (Bilim Pharmaceuticals, Turkey); Fluconazole (Pfizer Pharmaceuticals, Roerig Division (New York, N.Y.)). All other analytical grade chemicals used without further purification were purchased from Sigma-Aldrich (Hamburg, Germany).

2.2 Preparation of Plant Extract

The plants leaves were washed with distilled water, dried in the shade with continuous airflow, and then grounded into powder as 1-3 mm pieces. The powder of *PM* was randomized aliquoted into 20 mg using filter paper (Whatmann No.1) and then was macerated with 200 ml solution for 8 hours in Soxhlet (Wisd, Wise Therm) for each solution. The extraction step was done with four different solvents as follows: hexane (C_6H_{14} , $T_B:69^\circ C$), methanol (CH_3OH , $T_B:64,7^\circ C$), ethanol (C_2H_5OH , $T_B:$

78.4°C and water (H₂O, T_B:100°C). After this step, all extracts were evaporated until dried at 50 °C to obtain a solvent-free extract, which will be stored in the refrigerator (0-4 °C) until experiments. Each extract was redissolved in Dimethyl sulfoxide (DMSO) to yield a final concentration of 32 mg/mL. All of them were sterilized by a membrane filter (0.2 µm) before use [9].

2.3 Antimicrobial and Antifungal Activity Assay

Antimicrobial activity of extracts was applied to Gram-positive bacteria cultures: *Staphylococcus aureus* (ATCC 6538), *Enterococcus faecalis* (ATCC 292112), *Bacillus subtilis* (ATCC 6633), whereas gram-negative bacterial cultures *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (27853), *Proteus vulgaris* (ATCC 13315). Antifungal activity has been studied against *Candida albicans* (ATCC 60193) and *Candida tropicalis* (ATCC 13803).

Antimicrobial activity analyses of four different extracts of *Plantago major* were performed using Broth microdilution methods as recommended by Clinical & Laboratory Standards Institute [10-11]. In brief, the microorganism inoculum was prepared by 18 hours of fresh incubated microbial cultures adjusted with turbidity, which means a final concentration-1.5 × 10⁸ CFU/mL. The inoculums were pipetted in each well. The plant extracts were dissolved in dimethyl sulfoxide (DMSO), and after two times concentrated extracts at 32-0,156 mg/ml

concentration were pipetted in each well of microtiter plates. In brief, the wells were filled with 100 µL culture suspension as nutrient broth (Sigma-Aldrich, Germany) for antimicrobial activity and RPMI-1640 medium with L-Glutamin (Sigma-Aldrich, Germany) for antifungal activity as a growth medium. As per protocol, all extracts have been applied to wells 32 mg/mL and diluted 1/2 respectively. From 1-12 wells, it was from 32 to 0.156 mg/mL. The standard antibiotics and fungicide were used as Ampicillin, Gentamicin, and Fluconazole 2 mg/ml, respectively as control. The aerobic incubation condition was applied at 37 °C 18-24 h for bacteria and 48 h for fungi. The minimum inhibitory concentration (MIC) was calculated by visible inhibition of the microbial growth at the lowest concentration.

3. RESULTS AND DISCUSSION

Table 1 and Table 2 are summarized the antibacterial and antifungal effects of different extracts of *Plantago Major*. The antimicrobial activity of hexane and aqueous extracts of *PM* against all test microorganisms except *Enterococcus faecalis* was the same.

The results revealed that the efficiency of both ethanol and methanol extracts of *PM* was similar to all test microorganisms. Methanol and ethanol showed the highest activity (2 mg/mL) in *S. aureus*, *E. faecialis* and *P. aeruginosa* bacteria and *C. albicans* yeast however had low activity on *B.subtilis*, *E. coli* *P. vulgaris* bacteria, and *C. tropicalis* yeast.

Table 1. Antimicrobial activity of different extracts of *Plantago major*

Test Materials	MIC (mg/ml)					
	Gram (+)			Gram (-)		
	<i>Staphylococcus aureus</i> ATCC 6538	<i>Enterococcus faecalis</i> ATCC 292112	<i>Bacillus subtilis</i> ATCC 6633	<i>Escherichia coli</i> ATCC 25922	<i>Pseudomonas aeruginosa</i> ATCC 27853	<i>Proteus vulgaris</i> ATCC 13315
<i>PM</i> -nHexOH	4	2	4	8	4	8
<i>PM</i> -MeOH	2	2	4	4	2	4
<i>PM</i> -EtOH	2	2	4	4	2	4
<i>PM</i> -H ₂ O	4	4	4	8	4	8
Gentamicin	10 ⁻³	10 ⁻³	8.10 ⁻³	16.10 ⁻³	8.10 ⁻³	16.10 ⁻³
Ampicilin	1610 ⁻³	16.10 ⁻³	16.10 ⁻³	32.10 ⁻³	32.10 ⁻³	16.10 ⁻³
DMSO	-	-	-	-	-	-

Table 2. Antifungal activity of different extracts of *Plantago major*

Test Materials	MIC (mg/ml)	
	<i>Candida albicans</i> ATCC 60193	<i>Candida tropicalis</i> ATCC 13803
<i>PM</i> -nHexOH	4	4
<i>PM</i> -MeOH	2	4
<i>PM</i> -EtOH	2	4
<i>PM</i> -H ₂ O	4	4
Fluconazole	625. 10 ⁻³	25. 10 ⁻¹
DMSO	-	-

In addition, the results (Tables 1-2) show that methanol and ethanol extracts showed higher activity than hexane and water extract. It is known that hexane has polar solubility, methanol and ethanol have semi-polar (methanol is more polar than ethanol), and water has apolar solubility. These differences explain that ethanol and methanol would be better solvents for *PM*, and bioactive compounds were more easily resolved in ethanol and methanol. The four types of *PM* extracts exhibited inhibitory effects against test microorganisms; however, these antimicrobial activities were below the MIC value of positive controls (ampicillin gentamicin and fluconazole). thus, we can conclude that *PM* extracts show antimicrobial activity but are not as effective as standard therapeutics.

This study provides further documentation of the applicability of the *Plantago major* extract against microorganisms. Previous studies have shown that *PM* has no antimicrobial activity against primary plaque colonizers or periodontal pathogens [8]. However, we clearly showed that *PM* ethanol extract showed antimicrobial activity against *Staphylococcus aureus*, *Enterococcus faecalis*, *Pseudomonas aeruginosa* at 2ml/mL, and *Bacillus subtilis* *Escherichia coli* and *Proteus vulgaris* at 4 ml/mL. In addition to ethanol extract, hexane, methanol, and water extract showed antimicrobial activity. Together with this, all extracts showed antifungal activity against *Candida albicans* and *Candida tropicalis*. In a previous study, the Kirby-Bauer disc diffusion technique was used to investigate the antimicrobial activity, which is not recommended as a reference method.

In contrast, we clearly showed antimicrobial activity by the MIC method, which is a gold standard used in clinics to calculate the antimicrobial activity at routine practice. In addition, Sharma and colleagues prepared their text extract with maceration; thus, the preparation procedure should affect the concentration of bioactive compounds in total. Similar to our study, Ferrazzano and colleagues [12] demonstrated a significant antimicrobial effect of *Plantago lanceolata* and evaluated the plant extract as a natural anti-cariogenic agent. Although Ferrazzano et al. studied different species, both plants have the same genus and have similar phytochemicals. The broth dilution MICs for *Plantago major* extracts tested against *C. albicans* and *C. tropicalis* showed significant inhibition, which means *PM* extracted by different solvents might be used with its antifungal activity. In summary, we have provided documentation of *PM* extracts with antimicrobial and antifungal activity; therefore, *PM* extracts should be a good candidate for drug development.

4. CONCLUSION

This study demonstrated the antimicrobial activity of *Plantago major* against all test microorganisms. In addition, n-EtOH and n-MeOH extract of *PM* showed better antimicrobial and antifungal activity when compared to hexane and aqueous extracts. The difference in antimicrobial efficiency at different extracts clearly showed that the extraction method is important to obtain bioactive molecules. Despite significant improvements in synthetic molecules in the pharmaceutical industry, the growing antibiotic resistance problem still requires finding

alternative antimicrobial agents. In this frame, the discovery of plant-based antibiotics is also essential to reduce the side effects of synthetic antimicrobials. Thus, we recommended testing different extract types and using gold standards as antimicrobial activity determination. The results of our study show that *PM* extracts are effective as an antimicrobial agent. However, the active component of the plant extracts should be enhanced, or active components should be purified by high-throughput techniques to develop bio-based pharmaceuticals.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Wang H, Zhao C, Huang Y, Wang F, Li Y, Chung HY. Chemical Constituents and Bioactivities of Plantaginis Herba. *Hong Kong Med J*. 2015;22:29–35.
2. Samuelsen AB. The traditional uses, chemical constituents and biological activities of *Plantago major* L. A review. *J Ethnopharmacol*. 2000;71:1–21.
3. Najafian Y, Hamed SS, Farshchi MK, Feyzabadi Z. *Plantago major* in Traditional Persian Medicine and modern phytotherapy: A narrative review. *Electron Physician*. 2018;10(2):6390-9.
4. Ghanadian M, Soltani R, Homayouni A, Khorvash F, Jouabadi SM, Abdollahzadeh M. The Effect of *Plantago major* Hydroalcoholic Extract on the Healing of Diabetic Foot and Pressure Ulcers. A Randomized Open-Label Controlled Clinical Trial. *Int J Low Extrem Wounds* 2022; (In press).
5. Parhizgar S, Hosseinian S, Soukhtanloo M, Bideskan AE, Hadjzadeh MA, Shahraki S et al. *Plantago major* protects against cisplatin-induced renal dysfunction and tissue damage in rats. *Saudi J Kidney Dis Transpl*. 2018;29(5):1057-64.
6. Boskabadi J, Saadat S, Boskabady MH. The Relaxant Effect of *Plantago Major* on Rat Tracheal Smooth Muscles and Its Possible Mechanisms. *Iran J Allergy Asthma Immunol*. 2020;19(4):386-96.
7. Jazayeri SF, Ghods R, Hashem Dabaghian F, Shojaii A, Moravej SAA, Khadem E et al. A Randomized Double-Blind Clinical Trial. *Evid Based Complement. Alternat Med*. 2021;6693887.
8. Sharma H, Yunus GY, Mohapatra AK, Kulshrestha R, Agrawal R, Kalra M. Antimicrobial efficacy of three medicinal plants *Glycyrrhiza glabra*, *Ficus religiosa*, and *Plantago major* on inhibiting primary plaque colonizers and periodontal pathogens: An *in vitro* study. *Indian J Dent Res*. 2016;27(2):200-4.
9. Kaur K, Arora S, Kumar S, Nagpal A.. Antimutagenic Activities of Acetone and Methanol Fractions of *Terminalia arjuna*. *Food and Chemical Toxicology*. 2002;40: 1475-82.
10. CLSI, Method for Antifungal Disk Diffusion Susceptibility Testing of Yeasts, Approved Guideline. CLSI document M44-A. CLSI, 940 West Valley Road, Suite 1400, Wayne, Pennsylvania 19087-1898, USA; 2004.
11. CLSI, Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically; Approved Standard-Ninth Edition. CLSI Document M07-A9. Wayne, Pennsylvania; 2012b.
12. Ferrazzano GF, Cantile T, Roberto L, Ingenito A, Catania MR, Roscetto E et al. Determination of the *in vitro* and *in vivo* antimicrobial activity on salivary Streptococci and Lactobacilli and chemical characterisation of the phenolic content of a *Plantago lanceolata* infusion. *Biomed Res Int*. 2015;286817.

© 2021 Akkuş and HIZ-ÇİÇEKLYURT; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85117>