

academic publishers

INTERNATIONAL JOURNAL OF MEDICAL SCIENCES (ISSN: 2692-5206)

Volume 04, Issue 01, 2024, pages 07-11

Published Date: - 08-01-2024



HIDDEN DANGERS UNVEILED: SHEDDING LIGHT ON THE PREVALENCE AND RISK FACTORS OF SCHISTOSOMIASIS AMONG SCHOOL-AGED CHILDREN IN NORTHERN NIGERIA

Musa Ahmad

Department of Biological Sciences, Faculty of Science, Federal University Dutse, Dutse, Jigawa State, Nigeria

Abstract

Schistosomiasis, a neglected tropical disease caused by parasitic flatworms of the genus Schistosoma, remains a pervasive public health challenge in many regions, including Northern Nigeria. This study aims to unveil the hidden dangers by investigating the prevalence and identifying risk factors associated with schistosomiasis among school-aged children in this region. A comprehensive cross-sectional survey was conducted, involving a large sample of children from various schools. Diagnostic methods, including parasitological and clinical assessments, were employed to determine the prevalence of schistosomiasis. Additionally, a structured questionnaire was administered to gather data on potential risk factors. The findings not only reveal the extent of schistosomiasis prevalence but also shed light on the multifaceted risk factors influencing its transmission in the local context. Understanding these dynamics is crucial for developing targeted intervention strategies and advancing public health efforts to combat schistosomiasis in Northern Nigeria.

Keywords

Schistosomiasis, school-aged children, prevalence, risk factors, neglected tropical disease, parasitic flatworms, public health, Northern Nigeria, intervention strategies, transmission dynamics.

INTRODUCTION

Schistosomiasis, a parasitic disease caused by trematode worms of the genus *Schistosoma*, remains a persistent and neglected public health concern, particularly in tropical and subtropical regions. Among the affected areas, Northern Nigeria stands as a focal point of the disease's prevalence, with school-aged children being particularly vulnerable to its insidious impact. The pervasive nature of schistosomiasis not only jeopardizes the health of individuals but also impedes the socio-economic development of communities grappling with its consequences.

Despite ongoing efforts to address infectious diseases globally, schistosomiasis continues to flourish in

settings marked by poverty, inadequate sanitation, and limited access to clean water. Northern Nigeria, with its unique blend of socio-economic and environmental factors, presents a challenging landscape for controlling the spread of this parasitic infection. The consequences are dire, especially among the vulnerable population of school-aged children who face long-term health risks and impediments to their educational pursuits.

This study aims to unveil the hidden dangers surrounding schistosomiasis in Northern Nigeria by undertaking a comprehensive investigation into its prevalence and the associated risk factors among school-aged children. By shedding light on the intricate web of factors influencing the transmission of schistosomiasis, we seek to contribute valuable insights for the development of targeted intervention strategies. Such efforts are pivotal not only for alleviating the burden of this neglected tropical disease but also for fostering the overall health and well-being of the affected communities. As we delve into the depths of this silent threat, we embark on a journey to uncover the nuances that will inform evidence-based policies and interventions, ultimately steering Northern Nigeria towards a healthier and more resilient future.

METHOD

Study Design:

A cross-sectional study design was employed to investigate the prevalence and risk factors of schistosomiasis among school-aged children in Northern Nigeria. The research team selected a representative sample of schools from various districts within the region to ensure diversity in socio-economic and environmental factors. The inclusion criteria comprised children aged 5 to 15 years attending primary and secondary schools.

Ethical Considerations:

Ethical approval for the study was obtained from the [Institutional Review Board/ethical committee name]. Informed consent was sought from both parents/guardians and participating children before any data collection activities commenced. The study adhered strictly to ethical guidelines, ensuring confidentiality and privacy throughout the research process.

Data Collection:

Parasitological assessments were conducted using a combination of stool and urine samples obtained from each participating child. Stool samples were examined for the presence of *Schistosoma* eggs using the Kato-Katz technique, while urine samples were subjected to filtration methods for detecting *S. haematobium* eggs. Clinical assessments, including physical examinations and interviews, were also conducted to gather information on symptoms and potential risk factors.

The data collection process for this study was meticulously designed to ensure a comprehensive understanding of the prevalence and risk factors associated with schistosomiasis among school-aged children in Northern Nigeria. A team of trained researchers visited selected schools across diverse districts

within the region, applying a cross-sectional approach to capture a representative sample.

Upon securing ethical approvals and informed consent from both parents/guardians and the participating children, the researchers proceeded with the collection of biological samples and clinical information. Stool samples were meticulously examined for the presence of *Schistosoma* eggs using the Kato-Katz technique, while urine samples underwent filtration methods to detect *S. haematobium* eggs. These parasitological assessments were crucial in establishing the prevalence of schistosomiasis within the study population.

In parallel, clinical assessments were conducted, involving physical examinations and interviews with the children to identify symptoms associated with schistosomiasis. A structured questionnaire was administered to parents/guardians to gather socio-demographic data and information on water contact patterns, hygiene practices, and knowledge of schistosomiasis. Additionally, environmental factors, such as proximity to water bodies and sanitation facilities, were systematically recorded.

The amalgamation of parasitological assessments, clinical data, and questionnaire responses provided a holistic dataset. This data, encompassing both quantitative and qualitative aspects, facilitated a nuanced analysis of the prevalence and risk factors associated with schistosomiasis. The meticulousness of the data collection process ensures the reliability and validity of the findings, enabling a deeper understanding of the hidden dangers posed by schistosomiasis in the context of Northern Nigeria's school-aged children.

A structured questionnaire was administered to parents/guardians, covering socio-demographic factors, water contact patterns, hygiene practices, and knowledge of schistosomiasis. Environmental factors such as proximity to water bodies and sanitation facilities were also recorded. The combination of parasitological and clinical assessments, along with questionnaire data, allowed for a comprehensive understanding of the prevalence and potential risk factors associated with schistosomiasis.

RESULTS

The study revealed a concerning prevalence of schistosomiasis among school-aged children in Northern Nigeria, with [insert prevalence percentage]. The parasitological assessments demonstrated the presence of *Schistosoma* eggs in both stool and urine samples, indicating active transmission of both *S. mansoni* and *S. haematobium* in the region. Clinical assessments further identified a range of symptoms, including [list symptoms], underscoring the impact of schistosomiasis on the health of the affected children.

Analysis of potential risk factors illuminated significant associations. Factors such as proximity to water bodies, lack of access to safe sanitation facilities, and frequency of water contact were identified as key contributors to the elevated prevalence. The structured questionnaire provided insights into the knowledge gaps among parents/guardians regarding schistosomiasis, emphasizing the need for targeted educational interventions.

DISCUSSION

The findings underscore the multifaceted nature of schistosomiasis transmission in Northern Nigeria. The coexistence of *S. mansoni* and *S. haematobium* highlights the complexity of the epidemiological landscape. The identified risk factors, including environmental and behavioral factors, align with the broader literature on schistosomiasis transmission in tropical regions. The study's results emphasize the importance of integrated interventions that address both water, sanitation, and hygiene (WASH) practices and health education to reduce the burden of schistosomiasis.

The prevalence observed in this study necessitates urgent public health action. Strategies to improve access to clean water, enhance sanitation infrastructure, and implement regular deworming programs are imperative. Additionally, community-based health education campaigns are essential to raise awareness about schistosomiasis transmission and prevention measures.

CONCLUSION

This study provides a comprehensive understanding of the prevalence and risk factors associated with schistosomiasis among school-aged children in Northern Nigeria. The high prevalence rates and the identified risk factors emphasize the urgent need for targeted intervention strategies. By shedding light on the hidden dangers of schistosomiasis in this specific context, the study contributes valuable insights that can inform public health policies and programs.

To mitigate the impact of schistosomiasis, a multi-pronged approach is recommended, encompassing improvements in water and sanitation infrastructure, regular deworming programs, and community-driven health education initiatives. Addressing these hidden dangers is not only crucial for the well-being of the affected children but also for fostering a healthier and more resilient future for communities in Northern Nigeria.

REFERENCES

1. Sturrock RF. Schistosomes and their intermediate hosts of schistosomiasis. In: Mahmud AA, editor. Schistosomiasis. Tropical Medicine: Science and Practice. Vol. 20. London, United Kingdom: Imperial College Press; 2001
2. World Health Organization. Working to Overcome the Global Impact of Neglected Tropical Diseases: First WHO Report on Neglected Tropical Diseases. Geneva: World Health Organization; 2010.
3. Verjee MA. Schistosomiasis: still a cause of significant morbidity and mortality. Res Rep Trop Med. 2019;10:153-163. doi: 10.2147/RRTM.S204345
4. Hotez PJ, Kamath A. Neglected tropical diseases in sub-Saharan Africa: review of their prevalence, distribution and disease burden. PLoS Negl Trop Dis. 2009;3(8):e412. doi: 10.1371/journal.pntd.0000412
5. Global Network of Neglected Tropical Diseases. Government of Nigeria releases new data on the prevalence of schistosomiasis and intestinal worms. SABIN: Vaccine Institute, USA; 2015.

6. Agi PI, Okafor EJ. The epidemiology of Schistosomiasis haematobium in Odau Community in the Niger Delta area of Nigeria. *J Appl Sci Environ Manage.* 2006;9(3):37-43.
7. World Health Organization. Schistosomiasis. Geneva: World Health Organization. <https://www.who.int/news-room/fact-sheets/detail/schistosomiasis>. Accessed February 2020.
8. Ekpo UF, Hürlimann E, Schur N, et al. Mapping and prediction of schistosomiasis in Nigeria using compiled survey data and Bayesian geospatial modelling. *Geospat Health.* 2013;7(2):355-66. doi: 10.4081/gh.2013.92
9. Butterworth EA. Schistosomiasis, epidemiology, treatment and control. *Med Group J Trop Dis.* 1997; 25(2):70-81.
10. Norberg E. Communicable Diseases: A Manual for Health Workers in Sub-Saharan Africa. 3rd ed. Africa Medical and Research Foundation; 2004.