



Surgical Management of Early Stage Invasive Breast Cancer - A Practice Guideline

Hakeem Sulthan. Z¹, Dr. Dhananjayan Govindarajan², Dr. N. K. S. Shaik Mujibur Rahman^{3*}

¹Senior Resident, Department of General Surgery, Shri Sathya Sai Medical College & Research Institute, Ammapettai - 603108, Tamilnadu.

²Senior Resident, Department of General Surgery, Shri Sathya Sai Medical College & Research Institute, Ammapettai - 603108, Tamilnadu.

³Associate Professor, Department of General Surgery, Shri Sathya Sai Medical College & Research Institute, Ammapettai - 603108, Tamilnadu.

Corresponding Author: Dr. N. K. S. Shaik Mujibur Rahman, Associate Professor, Department of General Surgery, Shri Sathya Sai Medical College & Research Institute, Ammapettai - 603108, Tamilnadu.

Received Date: 15/04/2025

Revised Date: 10/05/2025

Accepted Date: 08/06/2025

KEYWORDS

Ayurveda, Plantar Psoriasis, Vaipadika, Shodhana, Shamana, Case Report, Kshudra Kushta

ABSTRACT:

Background: Objectives: To assess the available evidence on sentinel lymph-node biopsy, and to examine the long term follow-up data from large randomized phase III trials comparing breast-conserving therapy with mastectomy in order to make recommendations on the surgical management of early invasive breast cancer (stages I and II), including the optimum management of the axillary nodes: for the breast - modified radical mastectomy or breast-conserving therapy; for the axilla - complete axillary node dissection, axillary dissection of levels I and II lymph nodes, sentinel lymph-node biopsy or no axillary node surgery.

Outcomes: Overall survival, disease-free survival, local recurrence, distant recurrence and quality of life.

Evidence: MEDLINE, EMBASE, the Cochrane Library databases and relevant conference proceedings were searched to identify randomized trials and meta-analyses. Two members of the Practice Guidelines Initiative, Breast Cancer Disease Site Group (BCDSG) selected and reviewed studies that met the inclusion criteria. The systematic literature review was combined with a consensus process for interpretation of the evidence to develop evidence-based recommendations. This practice guideline has been reviewed and approved by the BCDSG, comprising surgeons, medical oncologists, radiation oncologists, pathologists, a medical sociologist, a nurse representative and a community representative.

Benefits, harms and costs: Breast-conserving therapy (lumpectomy with levels I and II axillary node dissection, plus radiotherapy) provides comparable overall and disease-free survival to modified radical mastectomy. Levels I and II axillary dissection accurately stages the axilla and minimizes the morbidity of axillary recurrence but is associated with lymphedema in approximately 20% of patients and arm pain in approximately 33%. Currently, there is insufficient data regarding locoregional recurrence and long term morbidity associated with sentinel-node biopsy to advocate it as the standard of care. Breast conserving therapy may offer an advantage over mastectomy in terms of body image, psychological and social adjustment but appears equivalent with regard to marital adjustment, global adjustment and fear of recurrence.

Recommendations: Women who are eligible for breast-conserving surgery should be offered the choice of either breast-conserving therapy with axillary dissection or modified radical mastectomy.



Removal and pathological examination of levels I and II axillary lymph nodes should be the standard practice in most cases of stages I and II breast carcinoma. There is promising but limited evidence to support recommendations regarding sentinel lymph-node biopsy alone. Patients should be encouraged to participate in clinical trials investigating this procedure.

Validation: A draft version of this practice guideline and a 21-item feedback questionnaire was circulated to 201 practitioners in Ontario. Of the 131 practitioners who returned the questionnaire, 98 (75%) completed the survey and indicated that the report was relevant to their clinical practice. Eighty (82%) of these practitioners agreed that the draft document should be approved as a practice guideline.

INTRODUCTION

Breast cancer is the most commonly diagnosed cancer among Indian women, excluding non melanoma skin cancer. In 2004, this disease will have been diagnosed in an estimated 21 200 women in India.¹ The majority of these women will have operable breast cancer.

Surgery continues to play a prominent role in the management of early invasive breast cancer. Over the last 2 decades, breast-conserving therapy has become established as a viable alternative to modified radical mastectomy. More recently, sentinel lymph-node biopsy has shown promise as a conservative approach to staging the axilla. At present, axillary dissection of levels I and II lymph nodes is standard in India. This procedure accurately stages the axilla and minimizes the risk of axillary recurrence. However, in some women, axillary dissection is associated with infection, cutaneous numbness and dysesthesia, and significantly lymphedema.² A less invasive method of axillary evaluation is very appealing given the potential morbidity associated with axillary dissection. However, caution is warranted before adopting a more conservative staging procedure. Axillary lymph node status remains the single most important prognostic factor for breast cancer.³ Adjuvant systemic and regional therapy decisions are based, in large part, on whether, and to what extent, axillary lymph nodes are involved.

A practice guideline for the surgical management of early stage invasive breast cancer was originally completed in 1996 and subsequently published.⁴ The rationale for revisiting the practice in this evidence based practice guideline is 2-fold: (1) to assess the available evidence on sentinel-node biopsy, and (2) to examine the long-term follow-up data from the large randomized phase III

trials comparing breast conserving therapy to mastectomy.

METHODS

Literature search strategy

We searched MEDLINE (1966–April 2004), EMBASE (1980–April 2004) and the Cochrane Library (issue 1, 2004). The strategy used disease specific terms (breast neoplasms/ or breast cancer [tw] or mammary neoplasms/) in combination with treatment-specific terms (mastectomy/ or mastectomy [tw, sh] or mastectomy or segmental/ or lumpectomy [tw] or breast conserv [tw] or conserv [tw] or sentinel [tw] or axilla [tw]) and terms specifically for study design and publication type (meta-analysis [pt, sh, tw] or randomized controlled trial [sh, pt, tw] or randomized controlled trials/ or random [tw]). The search was not limited by language of publication.

The Physician Data Query (PDQ) clinical trials database on the Internet (www.cancer.gov/search/clinical_trials/) and the proceedings of the annual meetings (1999–2003) of the American Society of Clinical Oncology and the American Society for Therapeutic Radiology and Oncology were searched for reports of new trials. Relevant studies were retrieved and reviewed by 2 Breast Cancer Disease Site Group (BCDSG) members. References lists of included studies were searched for additional trials.

Inclusion criteria

Published abstracts and full reports were eligible for inclusion if they were randomized controlled trials (RCTs) or meta-analyses comparing (a) breast-conserving therapy to mastectomy or (b) the surgical management of the axilla by either complete axillary node clearance, dissection of levels I and II lymph nodes,



sentinel node biopsy or no axillary surgery. Randomized trials or meta-analyses investigating the efficacy and safety of sentinel-node biopsy were also eligible. Outcomes of interest included overall or disease-free survival, local recurrence, distant recurrence and quality of life.

Synthesizing the evidence

A quantitative synthesis of the data from randomized trials comparing breast-conserving surgery and mastectomy was not undertaken for this practice guideline, as 3 meta-analyses have been conducted and published on this topic.^{5–7} Similarly, it was decided not to pool the results of randomized trials comparing mastectomy or lumpectomy plus radiation with or without axillary dissection because a published meta-analysis was available⁸ that synthesized the data from all of the randomized trials identified on this topic.

Guideline development

This practice guideline report was developed by the Practice Guidelines Initiative of Cancer Care Ontario's Program in Evidence-Based Care, using the methodology of the Practice Guidelines Development Cycle.⁹ Evidence was selected and reviewed by members of the Practice Guidelines Initiative's BCDSG and methodologists.

The guideline is a convenient and up-to-date source of the best available evidence on the surgical management of early stage breast cancer, developed through systematic reviews, evidence synthesis and input from practitioners in Ontario. The body of evidence in this report primarily comprises RCT data; therefore, recommendations by the BCDSG are offered. The report is intended to enable evidence-based practice. The Practice Guidelines Initiative is editorially independent of Cancer Care Ontario and the Ontario Ministry of Health and Long-Term Care. External review by Ontario practitioners was obtained through a mailed survey consisting of items that address the quality of the draft practice guideline report and its recommendation and whether the recommendations should serve as a practice guideline. Final approval of the original guideline report

has been obtained from the Practice Guidelines Coordinating Committee.

The Practice Guidelines Initiative has a formal standardized process to ensure the currency of each guideline report. This consists of the periodic review and evaluation of the scientific literature and, where appropriate, integration of the literature with the original guideline information. This document replaces the practice guideline report on the surgical management of early stage breast cancer originally completed in 1996.⁴ The recommendation concerning breast conserving therapy versus mastectomy is similar to that made in 1996, but new recommendations on axillary lymph-node dissection and sentinel node biopsy have been added.

RESULTS

Literature search results

We identified 10 RCTs and 2 un-published studies (see Table 110–19) and 3 meta-analyses^{5–7} that compared the effect of breast-conserving therapy to mastectomy on overall survival or recurrence. Surgical management of the axilla was evaluated in 6 RCTs^{17,18,20–28} and 1 meta-analysis⁸ on axillary dissection. One RCT²⁹ and 1 meta-analysis³⁰ of case series on sentinel-node biopsy were found. The literature search also identified 14 papers that reported quality-of-life data from randomized trials of breast-conserving therapy versus mastectomy^{31–44} and 1 meta-analysis⁴⁵ on the topic.

Surgical management: breast conserving therapy versus mastectomy

Key results from the 12 randomized trials that compare breast-conserving therapy with mastectomy for women with early stage breast cancer are summarized in Table 1 (unpublished data).^{10–19} Six trials are considered the standard in the field.^{10–15} Of the remaining 6 trials, the Argentinean¹⁶ and the 2 Guy's Hospital series^{17,18} had significant methodologic irregularities, and results from 3 trials reported in the meta-analysis by the Early Breast Cancer Trialists' Collaborative Group (EBCTCG)⁵ were never published (2 trials) or only published in abstract form¹⁹ (1 trial).

**Table 1: Randomized trials comparing breast-conserving therapy with mastectomy**

Table 1 Randomized trials comparing breast-conserving therapy with mastectomy						
Study	Comparison*	No. of patients	Time point, yr	Overall survival, %	Disease-free survival, %	Local recurrence, %†
Fisher et al ¹⁰	Lumpectomy	634		53	64	39‡
	Lumpectomy and radiation	628	20	50	62	14
	Total mastectomy	589		51	63	—
Veronesi et al ¹¹	Quadrantectomy and radiation	352	20	41	NR	8‡
	Modified radical mastectomy	349		42	NR	2
van Dongen et al ¹²	Breast-conserving surgery	448	10	65	47	20‡
	Modified radical mastectomy	420		66	53	12
Arriagada et al ¹³	Tumorectomy and radiation	88	22	60	NR	9
	Modified radical mastectomy	91		49	NR	14
Poggi et al ¹⁴	Lumpectomy and radiation	121	18	54	63	NR
	Modified radical mastectomy	116		58	67	NR
Blicherf-Toff et al ¹⁵	Breast-conserving surgery	430	6	79	70	2
	Total mastectomy	429		82	66	—
Gori et al ¹⁶	Tumorectomy	58	5	95	NR	8
	Madden's procedure	63		93	NR	2
Hayward ¹⁷	Wide excision and radiation	122	6	NR	NR	30‡
	Total mastectomy and radiation	130				8
Atkins et al ¹⁸	Wide excision and radiation	184	10	NR	NR	40‡
	Total mastectomy and radiation	192				18
D'Aluto et al ¹⁹	Breast-conserving surgery	170	NR	88	NR	NR
	Mastectomy	170		85		
CRC UK, 1995§	Breast-conserving surgery	71	NR	80	NR	NR
	Mastectomy	74		82		
BMFT 01 Germany, 1995§	Breast-conserving surgery	41	NR	90	NR	NR
	Mastectomy	31		95		

*Axillary dissection was carried out in all patients except for the breast conservation arms in the 2 Guy's Hospital trials.^{17,18}
†There was substantial variability between studies in how local or locoregional recurrence was defined.
‡Indicates a significant difference at $p < 0.05$.
§The results of these studies are published in abstract form only.
BMFT = Bundesministerium für Forschung und Technologie; CRC UK = Cancer Research Campaign United Kingdom.

With the exception of the 2 Guy's Hospital trials,^{17,18} no significant differences were reported in overall survival, disease-free survival or distant disease-free survival in any of the studies comparing breast-conserving surgery and mastectomy.

All 3 published meta-analyses of RCTs comparing mastectomy and breast-conserving surgery reported no significant difference in survival between the 2 groups.^{5–7} The EBCTCG study included data from 4891 women who participated in 9 RCTs of mastectomy versus breast-conserving surgery plus radiotherapy.⁵ They reported a non-significant odds reduction for death of -2% (standard error, m7), which represented a 2% increase in the odds of death in the mastectomy group compared with the breast-conserving therapy group ($p = 0.7$).

In 1997, Morris and colleagues⁶ published a meta-analysis using a combination of individual patient data and published results from 6 randomized trials. They reported pooled odds ratios for death of 0.90 (95%

confidence interval [CI] 0.74–1.09) at 5 years after randomization and 0.91 (95% CI 0.78–1.05) at 10 years. In 1998, abstract data from another meta-analysis by Morris and associates⁷ reported long-term data from 3 RCTs. After up to 20 years' follow up, no significant differences were detected between pooled overall survival (55% v. 48%, log rank $p = 0.95$) or distant recurrence (38% v. 36%, log rank $p = 0.61$) in the mastectomy and the breast-conserving therapy arms respectively.

Axillary node dissection

Results from 6 randomized trials of axillary node dissection versus no axillary node dissection^{17,18,20–28} were summarized in the meta-analysis by Orr.⁸ This meta-analysis was based on 4 decades of data from 2936 women who participated in 6 randomized trials comparing mastectomy or lumpectomy plus radiation with or without axillary dissection. Trials were eligible for inclusion if they included patient populations with



stage I or a combination of stages I and II disease. In 2 trials, the mean tumour size was not reported; 3 trials reported average tumour sizes greater than 3 cm, with positive nodes in 39%–54% of women. The authors of the meta-analysis reported that it was unlikely that any of the women had mammographically detected tumours and that adjuvant treatment with chemotherapy or tamoxifen would rarely have been used at the time these trials were conducted. The 6 trials reported an absolute survival benefit with axillary dissection ranging from 4% to 16%, which corresponds to a 7%–46% relative reduction in the risk of death. Orr reported a significant pooled survival benefit of 5.4% (95% CI 2.7%–8.0%, $p < 0.01$) favouring axillary dissection. However, the results must be viewed with caution since this meta-analysis was based only on published data rather than on individual patient data. Also, procedures other than levels I and II axillary node dissection were used in some of the studies, and in others axillary radiation was given to those who did not undergo axillary surgery. Although this meta-analysis suggests a significant survival benefit with axillary dissection, evolving approaches in surgical management, radiotherapy, adjuvant systemic therapy and screening practices may limit the magnitude of the survival benefit on women treated with current breast cancer therapy.

Longer term results for 220,28 of the 6 trials were identified.^{46,47} In the study from the Institut Curie, overall survival had decreased from 97% and 93% ($p = 0.014$) at 4.5 years to 76% and 74% ($p = \text{NS}$) at 15 years in the axillary dissection group versus the axillary radiotherapy group.⁴⁷ The lack of difference in disease-free survival, metastases and local recurrence was maintained. Recurrences in axillary nodes were still less frequent in the axillary dissection group (1% v. 3%, $p = 0.04$). In the National Surgical Adjuvant Breast and Bowel Project (NSABP) B-04 study, 25-year follow-up failed to show a significant difference in overall survival between the groups with and without axillary dissection (25% v. 26% respectively).⁴⁶ Rates of distant disease-free survival were 46% and 43% in the groups with and without axillary dissection respectively (HR = 1.10, 95% CI 0.89–1.35). These results tend to diminish the conclusions from the meta-analysis.

Sentinel lymph-node biopsy versus axillary dissection

To date, only 1 RCT has published data comparing sentinel-node biopsy with axillary node dissection.²⁹ In this trial from Milan, 516 women who had undergone quadrantectomy and sentinel-node biopsy were randomized to axillary node dissection or axillary node dissection only if the sentinel node contained metastatic cancer (SNB group). At a median follow-up of 46 months, there were 15 breast cancer events (ipsilateral or contralateral breast cancer or regional nodal or distant metastases) in the axillary node dissection group compared with 10 in the SNB group ($p = 0.26$). Two women died of breast cancer in the axillary node dissection group compared with 1 in the SNB group ($p = 0.26$).

One large ongoing randomized trial is comparing sentinel-node biopsy and axillary dissection. Similar to the Milan trial, the NSABP B-32 trial will evaluate sentinel-node biopsy with or without axillary node dissection.⁴⁸ Furthermore, a number of ongoing trials are studying women with positive sentinel nodes. The American College of Surgeons Oncology Group (ACOSOG) Z0010 trial is a prognostic study of sentinel-node and bone marrow micro metastases in women with early breast cancer.⁴⁹ The ACOSOG Z0011 trial will evaluate the effectiveness of axillary node dissection in women who have a positive sentinel node.⁵⁰ The International Breast Cancer Study Group (IBCSG) 23-01 trial will also compare the effectiveness of axillary node dissection in women with sentinel-node-positive disease.⁵¹ The European Organization for the Research and Treatment of Cancer (EORTC) 10981 trial will compare axillary node dissection with axillary radiotherapy in sentinel node-positive women.⁵² Of note, only the ACOSOG Z0011, the IBCSG 23-01, and the EORTC 10981 trials were still recruiting patients as of May 2004.

In 1999, Miltenburg and colleagues³⁰ published a meta-analysis of 11 case series in the literature between 1993 and 1998. Data were reported for 912 women with breast cancer who had sentinel-node biopsy followed by axillary node dissection. Overall, sentinel lymph nodes were successfully identified in 84% of women and concordance with pathological results from axillary dissection was confirmed in 98% of women. There was a 5% false-negative rate associated with sentinel node



biopsy. The highest identification rates were reached using either radiocolloid or dye and radio colloid combined. In fact, between January 1991 and December 2000, over 50 studies (involving more than 9000 women) have been reported.⁵³ The studies were all case series, some prospective and some retrospective. In all of these studies, patients first had a sentinel-node biopsy, followed by an axillary dissection. The false-negative rate ranged from 0% to 22%. It is important to note that falsely staging the presence of nodal metastases may affect the treatment a patient receives after surgery and possibly the chances of breast cancer recurrence.

Quality of life for women who choose breast-conserving therapy over mastectomy

Fourteen papers on quality of life, using data from randomized trials of breast-conserving surgery versus mastectomy, have been published.^{31–44} Poulsen and colleagues³¹ reported on 184 women who participated in the Danish Breast Cancer Cooperative Group trial. Over an average follow up of 31 months, no significant differences were found between the 2 types of surgery on measures of physical state, emotional state, social activity, work activity, body image, marital and sexual life or level of anxiety.

Curran and associates³² analyzed data from 278 women who participated in the European Organization for the Treatment of Cancer trial. Two years postoperatively, women in the breast-conserving therapy group had better body image ($p = 0.001$) and were more satisfied with treatment ($p = 0.001$) than those in the mastectomy group; there was no significant difference between the 2 groups with respect to fear of cancer recurrence ($p = 0.236$).

Although survival results are not yet forthcoming, quality-of-life data from the EORTC 10850 trial have been published.⁴⁴ In 136 women aged 70 years and older, women who underwent breast-conserving and tamoxifen therapy did not differ from those receiving mastectomy in terms of treatment preference, fatigue, emotional function, fear of recurrence, social support, physical functioning and leisure-time activities. Women who underwent breast conserving therapy did, however, report fewer arm problems ($p = 0.04$) and a trend toward improved body image ($p = 0.06$). Data from the remaining 11 randomized trials^{33–43} form the basis of the meta-analysis by Moyer.⁴⁵ Results favouring breast-conserving therapy were reported in 10 trials for

psychological adjustment (mean weighted effect size [MWES] = 0.060, standard deviation [SD] 0.66, $p < 0.001$) and in 3 trials for social adjustment (MWES = 0.334, SD 0.140, $p < 0.05$). No significant differences were detected in 7 trials measuring marital and sexual adjustment ($p > 0.05$) or body self image ($p > 0.05$), or in 6 trials measuring cancer-related fears and concerns ($p > 0.05$). The pooled effect size for global adjustment from 3 studies favoured mastectomy but was not statistically significant (MWES = -0.20, SD 0.108, $p > 0.05$).

Preoperative chemotherapy

Preoperative chemotherapy has been compared to the same chemotherapy given postoperatively in a number of randomized trials.^{54–56} None have shown any difference in disease-free or overall survival, but all have shown an increased likelihood of breast-conserving surgery in the preoperative chemotherapy arms. A study conducted at the Royal Marsden Hospital, London, using mitoxantrone and methotrexate chemotherapy, detected an 87% breast-conserving surgery rate in the preoperative chemotherapy arm compared with 72% in the postoperative arm ($p < 0.005$).⁵⁴ In the NSABP B-18 trial using doxorubicin and cyclophosphamide chemotherapy, 68% versus 60% of patients had breast-conserving surgery in favour of the preoperative treatment (p value not reported).⁵⁵ The EORTC study used fluorouracil, epirubicin and cyclophosphamide preoperatively versus postoperatively. In the preoperative group, 23% of the women who were considered initially to require mastectomy had breast-conserving surgery after preoperative chemotherapy.⁵⁶ The European Cooperative Trial in operable breast cancer used doxorubicin and paclitaxel for 4 cycles followed by 4 cycles of cyclophosphamide, methotrexate and fluorouracil, either pre- or postoperatively.⁵⁷ The primary tumour had to be more than 2 cm in dimension. Breast conserving surgery rates were 71% in the preoperative group versus 35% in the postoperative group ($p < 0.0001$).

Practitioner feedback results

In October 2001, 201 practitioners (42 medical oncologists, 41 radiation oncologists and 118 surgeons) were surveyed. Of the 131 practitioners who returned the questionnaire, 98 (75%) completed the survey and indicated that the report was relevant to their clinical practice; 89 (91%) of them indicated that they would use



the draft recommendations in their practice, and 80 (82%) agreed that the draft guideline should be approved as a practice guideline. Twenty-six respondents provided written comments. There was mixed feedback on the role of sentinel-node biopsy outside clinical trials. Some practitioners urged the adoption of sentinel-node biopsy by adequately trained surgeons. Others wanted clear evidence of survival equivalence before adopting sentinel-node biopsy as standard practice. Some practitioners questioned the need for axillary node dissection in elderly women with receptor-positive cancers who would be receiving tamoxifen regardless of the results of the dissection.

In response to the practitioner feedback survey, minor changes were made to the guideline report but not to the recommendations. The rationale for full dissection when the sentinel lymph node is positive for metastatic disease was added to the guideline report. The issue of sentinel node biopsy alone, outside a clinical trial, was discussed by the committee, as well as the reference by a number of respondents to the Canadian practice guideline on sentinel lymph-node biopsy by the Steering Committee on Clinical Practice Guidelines for the Care and Treatment of Breast Cancer.² The Canadian guideline recommends that axillary node dissection remain the standard of care and that, if a patient requests or is offered sentinel-node biopsy alone, she needs to be made aware of the risks and benefits and what is and what is not yet known about the procedure. The Canadian guideline confirms the lack of data from randomized trials comparing outcomes from sentinel-node biopsy to those with axillary node dissection; therefore, participation in randomized trials is encouraged. Since there is little evidence from randomized trials, the Canadian guideline is based on a consensus of the Steering Committee. In the opinion of the BCDSG although this consensus statement was reasonable, sentinel-node biopsy alone could not be recommended in the absence of high-quality evidence at the time of writing. No change was made to the guideline.

With regard to the omission of axillary node dissection in elderly hormone-receptor-positive women receiving tamoxifen, it is recognized that there may be some cases in which the omission of axillary node dissection could be justified. However, there are no randomized data confirming that such women do as well without as with axillary node dissection, and so omission of axillary node

dissection cannot be recommended as standard care. The International Breast Cancer Study Group is currently conducting a randomized trial comparing complete axillary dissection to tamoxifen in elderly women, and the group's study addresses this question.⁵⁸

DISCUSSION

For eligible candidates, surgical treatment options for early stage invasive breast carcinoma include breast-conserving surgery plus radiation or mastectomy. Six fully published RCTs^{10–15} have demonstrated comparable results between standard breast-conserving therapy and mastectomy in terms of overall survival and disease-free survival.

Although evidence relating quality of life to the extent of breast surgery is conflicting, women should be fully informed of the treatment implications involved with both mastectomy and breast-conserving surgery (i.e., the potential need for additional surgery and for adjuvant radiation therapy following breast-conserving surgery).

Evaluation of axillary lymph-node disease is an integral part of adjuvant treatment planning for most patients with stages I and II breast cancer. Although the surgical treatment of the axilla in cases of early stage breast cancer may or may not contribute significantly to a reduction in mortality in today's patient populations, it reduces the morbidity of axillary recurrence.

Axillary lymph-node dissection is the current standard of surgical care. It carries significant risk of lymphedema and long-term postoperative dysesthesias. With no set criteria used to define lymphedema and a variety of assessment techniques in use, there is wide variation in reported rates of lymphedema following axillary dissection. Rates ranging from 2% to 70% have been reported.⁵⁹ In a recent study, arm morbidity was assessed in 110 women after partial mastectomy with axillary dissection and in most cases, irradiation.⁶⁰ Lymphedema (defined as a >10% increase in arm volume) developed in 19% of the women, and 49% had reduced arm mobility (defined as a 15° impairment of shoulder mobility). After 5 years, 31% of women continued to report some arm pain after breast conserving therapy.

However promising, investigations for axillary staging such as sentinel node biopsy have not yet demonstrated acceptable specificity and sensitivity to be used routinely



outside the context of a clinical trial. Although sentinel-node biopsy alone is currently not a standard practice, a position paper by McCreedy and colleagues⁶¹ recommends that surgeons consider acquiring the necessary equipment, training and infrastructure to perform this technique. Surgeons should also collaborate with their colleagues in pathology and nuclear medicine to develop techniques and standards for proper handling and pathological assessment of these nodes.

Breast Cancer Disease Site Group consensus

With no observed differences in overall survival or distant recurrence, the opinion of the BCDSG was that, for eligible candidates, the choice between breast-conserving therapy and modified radical mastectomy should be based on patient preference.

To make an informed decision, patients should be fully aware of the risks and benefits of each procedure. Breast-conserving therapy typically involves tumour excision with uninvolved margins, axillary dissection and adjuvant breast irradiation. There is also a potential need for further surgery, possibly a mastectomy, in cases of local recurrence. A modified radical mastectomy involves the removal of the entire breast, including the nipple and areola complex, and the fascia over the pectoralis muscles, while sparing the underlying muscles and innervation. Breast reconstruction is an option for women who choose mastectomy.

The DSG agreed that at present there is insufficient evidence to make recommendations regarding sentinel node biopsy alone. The group acknowledged that some clinicians in Ontario are beginning to train for the procedure and are building expert teams in anticipation of the potential demand should sentinel-node biopsy alone become standard practice. The BCDSG agreed that patients should be encouraged to participate in clinical trials investigating this procedure.

Given that quality-of-life measures are difficult to capture objectively, the BCDSG believed that the evidence surrounding quality of life after surgery was conflicting. Whereas some evidence suggests that women who undergo breast-conserving therapy may have higher body self image than those who undergo mastectomy, other measures of psychosocial well-being were inconclusive.

Preoperative chemotherapy in patients with operable breast cancer does not appear to improve disease free or overall survival, but it does increase the likelihood of breast conserving surgery. The BCDSG therefore felt that preoperative chemotherapy could be considered for those who desire breast-conserving surgery but are not believed initially to be a good candidate because of the size of the tumour in relation to the size of the breast.

Practice guideline

This practice guideline applies to women with early stage (stages I and II) invasive breast cancer who are eligible for either breast-conserving therapy or mastectomy.

Recommendations

- Women who are eligible for breast-conserving therapy should be offered the choice of either breast-conserving therapy with axillary dissection or modified radical mastectomy.
- Removal and pathological examination of levels I and II axillary lymph nodes should be the standard practice in most cases of stages I and II breast carcinoma.
- Because, there is promising but limited evidence, not yet sufficient, to support a recommendation for sentinel lymph-node biopsy alone, patients should be encouraged to participate in clinical trials investigating this procedure.

Qualifying statements

- With no difference in survival or distant recurrence, the choice between breast-conserving therapy with axillary dissection and modified radical mastectomy should depend on patient preference when appropriate.
- Each patient should be fully informed of the risks and benefits of each procedure.
- Patients should be aware that breast-conserving therapy involves tumour excision with uninvolved margins, axillary dissection and adjuvant breast irradiation.
- Patients who choose breast conserving therapy should be aware that there is the potential need for further surgery, possibly a mastectomy, in cases of local recurrence.
- Evidence surrounding quality of life after surgery is conflicting, but there is some



evidence suggesting that women who receive breast-conserving therapy may have higher body self image than those who undergo mastectomy.

- In some instances, preoperative chemotherapy can shrink a larger primary tumour and allow for breast-conserving surgery followed by breast irradiation rather than mastectomy.

REFERENCES

1. National Cancer Institute of Canada. Canadian Cancer Statistics. Toronto: Canadian Cancer Society; 2004.
2. The Steering Committee on Clinical Practice Guidelines for the Care and Treatment of Breast Cancer. Canadian Association of Radiation Oncologists. Axillary dissection. *CMAJ* 1998;158(Suppl 3):S22-6.
3. Carter CL, Allen C, Henson DE. Relation of tumor size, lymph node status, and survival in 24,740 breast cancer cases. *Cancer* 1989;63:181-7.
4. Mirsky D, O'Brien SE, McCready DR, Newman TE, Whelan TJ, Levine MN. Surgical management of early stage invasive breast cancer (stage I and II). Provincial Breast Disease Site Group. *Cancer Prev Control* 1997;1:10-7.
5. Early Breast Cancer Trialists' Collaborative Group. Effects of radiotherapy and surgery in early breast cancer. An overview of the randomized trials. Early Breast Cancer Trialists' Collaborative Group. *N Engl J Med* 1995;333:1444-55.
6. Morris AD, Morris RD, Wilson JF, White J, Steinberg S, Okunieff P, et al. Breast-conserving therapy vs mastectomy in early-stage breast cancer: a meta-analysis of 10-year survival. *Cancer J Sci Am* 1997;3:6-12.
7. Morris AD, Arriagada R, Ancukiewicz M. When is survival a cure in early stage breast cancer? Twenty year follow-up from 3 randomized trials [abstract]. *Int J Radiat Oncol Biol Phys* 1998;42(Suppl 1):182.
8. Orr RK. The impact of prophylactic axillary node dissection on breast cancer survival— a Bayesian meta-analysis. *Ann Surg Oncol* 1999;6:109-16.
9. Browman GP, Levine MN, Mohide EA, Hayward RS, Pritchard KI, Gafni A, et al. The practice guidelines development cycle: a conceptual tool for practice guidelines development and implementation. *J Clin Oncol* 1995;13:502-12.
10. Fisher B, Anderson S, Bryant J, Margolese RG, Deutsch M, Fisher ER, et al. Twenty-year follow-up of a randomized trial comparing total mastectomy, lumpectomy, and lumpectomy plus irradiation for the treatment of invasive breast cancer. *N Engl J Med* 2002;347:1233-41.
11. Veronesi U, Cascinelli N, Mariani L, Greco M, Saccozzi R, Luini A, et al. Twenty-year follow-up of a randomized study comparing breast-conserving surgery with radical mastectomy for early breast cancer. *N Engl J Med* 2002;347:1227-32.
12. van Dongen JA, Voogd AC, Fentiman IS, Le Grand C, Sylvester RJ, Tong D, et al. Long-term results of a randomized trial comparing breast-conserving therapy with mastectomy: European Organization for Research and Treatment of Cancer 10801 trial. *J Natl Cancer Inst* 2000;92:1143-50.
13. Arriagada R, Le MG, Guinebretiere JM, Dunant A, Rochard F, Tursz T. Late local recurrences in a randomised trial comparing conservative treatment with total mastectomy in early breast cancer patients. *Ann Oncol* 2003;14:1617-22.
14. Poggi MM, Danforth DN, Sciuto LC, Smith SL, Steinberg SM, Liewehr DJ, et al. Eighteen-year results in the treatment of early breast carcinoma with mastectomy versus breast conservation therapy: the National Cancer Institute Randomized Trial. *Cancer* 2003;98:697-702.
15. Blichert-Toft M, Rose C, Andersen JA, Overgaard M, Axelsson CK, Andersen KW, et al. Danish randomized trial comparing breast conservation therapy with



- mastectomy:six years of life-table analysis. Danish Breast Cancer Cooperative Group. *J Natl Cancer Inst Monogr* 1992;(11):19-25.
16. Gori J, Castano R, Engel H, Toziano M, Fischer C, Maletti G. Conservative treatment vs. mastectomy without radiotherapy in aged women with breast cancer — a prospective and randomized trial. *Zentralbl Gynakol* 2000;122:311-7.
 17. Hayward JL. The Guy's trial of treatment of "early" breast cancer. *World J Surg* 1977;1:314-6.
 18. Atkins H, Hayward JL, Klugman DJ, Wayte AB. Treatment of early breast cancer: a report after ten years of a clinical trial. *BMJ* 1972;2:423-9.
 19. D'Aiuto G, Parisi V, Grasso M, *et al.* editors. *Conservative surgery in early breast cancer*. 5th European Conference on Clinical Oncology, London, UK: 1989. p. 0939.
 20. Fisher B, Redmond C, Fisher ER, Bauer M, Wolmark N, Wickerham DL, *et al.* Ten-year results of a randomized clinical trial comparing radical mastectomy and total mastectomy with or without radiation. *N Engl J Med* 1985;312:674-81.
 21. Johansen H, Kaae S, Schiødt T. Simple mastectomy with postoperative irradiation versus extended radical mastectomy in breast cancer. A twenty-five-year follow-up of a randomized trial. *Acta Oncol* 1990;29:709-15.
 22. Kaae S, Johansen H. Does simple mastectomy followed by irradiation offer survival comparable to radical procedures? *Int J Radiat Oncol Biol Phys* 1977;2(11-12):1163-6.
 23. Bruce J. Operable cancer of the breast. A controlled clinical trial. *Cancer* 1971;28:1443-52.
 24. Hamilton T, Langlands AO, Prescott RJ. The treatment of operable cancer of the breast: a clinical trial in the South-East region of Scotland. *Br J Surg* 1974;61:758-61.
 25. Langlands AO, Prescott RJ, Hamilton T. A clinical trial in the management of operable cancer of the breast. *Br J Surg* 1980;67:170-4.
 26. Hayward J, Caleffi M. The significance of local control in the primary treatment of breast cancer. Lucy Wortham James clinical research award. *Arch Surg* 1987;122:1244-7.
 27. Hayward JL. The Guy's Hospital trials on breast conservation. In: Harris JR, Hellman S, Silen W, editors. *Conservative management of breast cancer: new surgical and radiotherapeutic techniques*. Philadelphia: JB Lippincott; 1983. p. 77-90.
 28. Cabanes PA, Salmon RJ, Vilcoq JR, Durand JC, Fourquet A, Gautier C, *et al.* Value of axillary dissection in addition to lumpectomy and radiotherapy in early breast cancer. The Breast Carcinoma Collaborative Group of the Institut Curie. *Lancet* 1992;339:1245-8.
 29. Veronesi U, Paganelli G, Viale G, Luini A, Zurrida S, Galimberti V, *et al.* A randomized comparison of sentinel-node biopsy with routine axillary dissection in breast cancer. *N Engl J Med* 2003;349:546-53.
 30. Miltenburg DM, Miller C, Karamlou TB, Brunicardi FC. Meta-analysis of sentinel lymph node biopsy in breast cancer. *J Surg Res* 1999;84:138-42.
 31. Poulsen B, Graversen HP, Beckmann J, Blichert-Toft M. A comparative study of post-operative psychosocial function in women with primary operable breast cancer randomized to breast conservation therapy or mastectomy. *Eur J Surg Oncol* 1997;23:327-34.
 32. Curran D, van Dongen JP, Aaronson NK, Kiebert G, Fentiman IS, Mignolet F, *et al.* Quality of life of early-stage breast cancer patients treated with radical mastectomy or breast-conserving procedures: results of EORTC Trial 10801. The European Organization for Research and Treatment of Cancer (EORTC), Breast Cancer Cooperative Group (BCCG). *Eur J Cancer* 1998;34:307-14.



33. Schain W, Edwards BK, Gorrell CR, deMoss EV, Lippman ME, Gerber LH, *et al.* Psychosocial and physical outcomes of primary breast cancer therapy: mastectomy versus excisional biopsy and irradiation. *Breast Cancer Res Treat* 1983;3:377-82.
34. de Haes JC, Welvaart K. Quality of life after breast cancer surgery. *J Surg Oncol* 1985;28:123-5.
35. de Haes JC, van Oostrom MA, Welvaart K. The effect of radical and conservative surgery on the quality of life of early breast cancer patients. *Eur J Surg Oncol* 1986;12:337-42.
36. Fallowfield LJ, Baum M, Maguire GP. Effects of breast conservation on psychological morbidity associated with diagnosis and treatment of early breast cancer. *BMJ (Clin Res Ed)* 1986;293:1331-4.
37. Lasry JC, Margolese RG, Poisson R, Shibata H, Fleischer D, Lafleur D, *et al.* Depression and body image following mastectomy and lumpectomy. *J Chronic Dis* 1987;40:529-34.
38. Aaronson NK, Bartelink H, van Dongen JA, van Dam FS. Evaluation of breast conserving therapy: clinical, methodological and psychosocial perspectives. *Eur J Surg Oncol* 1988;14:133-40.
39. Kemeny MM, Wellisch DK, Schain WS. Psychosocial outcome in a randomized surgical trial for treatment of primary breast cancer. *Cancer* 1988;62:1231-7.
40. Levy SM, Herberman RB, Lee JK, Lippman ME, d'Angelo T. Breast conservation versus mastectomy: distress sequelae as a function of choice. *J Clin Oncol* 1989;7:367-75.
41. Ganz PA, Schag AC, Lee JJ, Polinsky ML, Tan SJ. Breast conservation versus mastectomy. Is there a difference in psychological adjustment or quality of life in the year after surgery? *Cancer* 1992;69:1729-38.
42. Lee MS, Love SB, Mitchell JB, Parker EM, Rubens RD, Watson JP, *et al.* Mastectomy or conservation for early breast cancer: psychological morbidity. *Eur J Cancer* 1992;28A(8-9):1340-4.
43. Schain WS, d'Angelo TM, Dunn ME, Lichter AS, Pierce LJ. Mastectomy versus conservative surgery and radiation therapy. Psychosocial consequences. *Cancer* 1994;73:1221-8.
44. de Haes JC, Curran D, Aaronson NK, Fentiman IS. Quality of life in breast cancer patients aged over 70 years, participating in the EORTC 10850 randomised clinical trial. *Eur J Cancer* 2003;39:945-51.
45. Moyer A. Psychosocial outcomes of breast-conserving surgery versus mastectomy: a meta-analytic review. *Health Psychol* 1997;16:284-98.
46. Fisher B, Jeong JH, Anderson S, Bryant J, Fisher ER, Wolmark N. Twenty-five-year follow-up of a randomized trial comparing radical mastectomy, total mastectomy, and total mastectomy followed by irradiation. *N Engl J Med* 2002;347:567-75.
47. Louis-Sylvestre C, Clough K, Asselain B, Vilcoq JR, Salmon RJ, Campana F, *et al.* Axillary treatment in conservative management of operable breast cancer: Dissection or radiotherapy? Results of a randomized study with 15 years of follow-up. *J Clin Oncol* 2004;22:97-101.
48. Phase III randomized study of sentinel node dissection with or without conventional axillary dissection in women with clinically node negative breast cancer. CANCERNET-PDQ [online database]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrid=66987&protocolsearchid=917856 (accessed 2005 Mar 30).
49. Phase III prognostic study of sentinel node and bone marrow micrometastases in women with stage I or IIA breast cancer. CANCERNET-PDQ [online database]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrid=67



- 017&protocolsearchid=917814 (accessed 2005 Mar 30).
50. Phase III randomized study of axillary lymph node dissection in women with stage I or IIA breast cancer who have a positive sentinel node. CANCERNETPDQ [online database]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrid=67018&protocolsearchid=917762 (accessed 2005 Mar 30).
51. Phase III randomized study of surgical resection with or without axillary lymph node dissection in women with clinically node-negative breast cancer with sentinel node micrometastases. CANCERNETPDQ [online database]. Available www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrid=339581&protocolsearchid=917850 (accessed 2005 Mar 30).
52. Phase III randomized study of complete axillary lymph node dissection versus axillary radiotherapy in sentinel lymph node positive women with operable invasive breast cancer. CANCERNET-PDQ [database online]. Available: www.nci.nih.gov/clinicaltrials/view_clinicaltrials.aspx?version=healthprofessional&cdrid=68566&protocolsearchid=917889 (accessed 2005 Mar 30).
53. Cantin J, Scarth H, Levine M, Hugi M. Clinical practice guidelines for the care and treatment of breast cancer: 13. Sentinel lymph node biopsy. *CMAJ* 2001;165:166-73.
54. Powles TJ, Hickish TF, Makris A, Ashley SE, O'Brien ME, Tidy VA, *et al.* Randomized trial of chemoendocrine therapy started before or after surgery for treatment of primary breast cancer. *J Clin Oncol* 1995;13:547-52.
55. Wolmark N, Wang J, Mamounas E, Bryant J, Fisher B. Preoperative chemotherapy in patients with operable breast cancer: nine-year results from National Surgical Adjuvant Breast and Bowel Project B-18. *J Natl Cancer Inst Monogr* 2001;(30):96-102.
56. van der Hage JA, van de Velde CJ, Julien JP, Tubiana-Hulin M, Vandervelden C, Duchateau L. Preoperative chemotherapy in primary operable breast cancer: results from the European Organization for Research and Treatment of Cancer trial 10902. *J Clin Oncol* 2001;19:4224-37.
57. Gianni L, Baselga J, Eirmann W, Porta VG, Semiglazov V, Garcia-Conde J, *et al.* First report of the European Cooperative Trial in operable breast cancer (ECTO): effects of primary systemic therapy (PST) on local-regional disease [abstract]. *Proc Am Soc Clin Oncol* 2002;21(1):34a.
58. Phase III randomized study of surgery with or without axillary node clearance followed by adjuvant tamoxifen in elderly women with breast cancer. CANCERNET-PDQ [online database]. Available: www.cancer.gov/search/ViewClinicalTrials.aspx?cdrid=78383&version=HealthProfessional&protocolsearchid=1008021 (accessed 2005 Mar 30).
59. Harris SR, Hugi MR, Olivetto IA, Levine M. Clinical practice guidelines for the care and treatment of breast cancer: 11. Lymphedema. *CMAJ* 2001;164:191-9.
60. Tengrup I, Tennvall-Nittby L, Christiansson I, Laurin M. Arm morbidity after breast-conserving therapy for breast cancer. *Acta Oncol* 2000;39:393-7.
61. McCreedy DR, Bodurtha AJ, Davis NL, Meterissian S, Robert J, Temple WJ. Sentinel lymph-node biopsy in breast cancer. *Can J Surg* 1999;42:406-7.