



## Comprehensive Management of Osteoporosis in Early Postmenopausal Women

Krishnaprabha<sup>(1)</sup>, Dr.MV Kumudavalli<sup>(2)</sup>

<sup>(1)</sup> ResearchScholar, Vinayaka Mission Research Foundation Deemed University & Assistant Professor, Department of PharmacyPractice,

St.James' College of Pharmaceutical Sciences, Chalakudy, Kerala, India

<sup>(2)</sup> Professor and Head of Department, Vinayaka Missions College of Pharmacy, Vinayaka Mission's Research Foundation (Deemed University), Salem, TamilNadu, India

*(Received: 27 September 2025 Revised: 05 October 2025 Accepted: 18 November 2025)*

### KEYWORDS

Postmenopause,  
Osteoporosis, fracture  
risk, Bone mineral  
density

### ABSTRACT:

As women navigate through perimenopause and into postmenopause, they face a significant and rapid decline in both bone density and strength, which notably increases their susceptibility to fractures, with hip fractures being the most severe outcome. In fact, falls are the leading cause of injury, death, and disability among individuals over the age of 65, and hip fractures, which often result from falls, represent one of the most debilitating consequences. Tragically, about half of older adults who are hospitalized for a hip fracture do not regain their previous level of independence or functional ability. The incidence of hip fractures is especially high among women, with those over 85 being nearly eight times more likely to suffer from this injury compared to women aged 65 to 74. Beyond age, other risk factors for falls include muscle weakness, physical limitations, environmental hazards, the use of psychoactive medications, and a history of previous falls. Moreover, factors like low body mass index, a sedentary lifestyle, osteoporosis, and a prior hip fracture further increase the risk of both falls and fractures.

Despite the well-established knowledge of these risk factors, osteoporosis remains frequently undiagnosed until a fracture occurs, largely because many women at risk are not properly identified or treated in clinical practice. Identifying women at high risk, particularly in the early postmenopausal stage, is crucial since early intervention can help maintain or even improve bone mass, thus significantly reducing the risk of fractures. To effectively prevent fractures, it is essential to focus on reducing the risk of falls, as approximately 95% of hip fractures are fall-related. Effective fall prevention strategies must be multifaceted, combining both behavioural and environmental modifications. Key interventions include educating older adults about the various risk factors for falls, encouraging strength-building exercises to improve balance and muscle function, making home modifications to eliminate hazards, and reviewing medications to minimize adverse side effects like dizziness or drowsiness.

Additionally, several pharmacological treatments have been shown to be effective in reducing fracture risk, such as bisphosphonates (e.g., alendronate, risedronate, and ibandronate) and selective oestrogen receptor modulators like raloxifene. These medications not only help increase bone density but also reduce the likelihood of fractures, with manageable side effects. For women who have stopped hormone replacement therapy, more vigilant monitoring is needed, as oestrogen withdrawal can lead to rapid bone loss and a heightened risk of fractures. To enhance the effectiveness of fall prevention programs, it is essential to improve coordination between various federal, state, and local organizations that are involved in addressing this issue. Additionally, there is a need for thorough evaluation of existing fall prevention programs, especially for at-risk groups such as women over 85 or those living with functional limitations. There is also a strong need for new primary prevention strategies, such as identifying footwear that promotes stability, as well as secondary prevention measures like protective hip pads that can mitigate injuries when falls occur. Collaboration among national experts from diverse fields is critical to ensure that priority research areas are identified, and comprehensive strategies are developed to address the growing concern of falls and fall-related injuries, ultimately aiming for long-term solutions that reduce fractures and improve quality of life for older adults.



## Introduction:

Osteoporosis is a progressive condition marked by the gradual loss of bone density and structural deterioration of bone tissue, which increases the likelihood of fractures. This disorder is especially common in postmenopausal women, as the sharp decline in oestrogen levels accelerates the rate of bone loss, heightening their susceptibility. The early postmenopausal phase, occurring within the first five to ten years after menstruation ceases, is a crucial period for both preventing and managing osteoporosis. During this stage, the body's bone breakdown surpasses the formation of new bone, resulting in a notable reduction in bone mineral density (BMD). The risk of developing osteoporosis in later years is closely tied to the peak bone mass achieved earlier in life, which typically occurs by the end of the second decade. During adolescence, females accumulate around 40%–50% of their total adult skeletal mass, with peak BMD in the hip being reached. Understanding how modifiable factors like physical activity, body composition, and calcium intake influence bone development during these formative years are critical.

While evidence supports those weight-bearing exercises significantly stimulate bone growth in adolescence, the exact role of calcium in boosting peak bone mass remains contentious. Several studies have investigated the effect of calcium consumption during adolescence, but findings indicate that supplementing calcium during this period does not result in substantial or long-lasting increases in bone mass. Given the serious consequences of osteoporosis, including fractures, impaired mobility, and a diminished quality of life, effective intervention strategies are crucial. Prevention and treatment efforts, including lifestyle adjustments, dietary interventions, and medications, can substantially reduce bone loss and fracture risks. This review examines the current and emerging approaches for preventing and managing osteoporosis during the early postmenopausal years, highlighting the importance of early diagnosis, personalized treatment plans, and the role of lifestyle modifications in supporting bone health.

## Evaluating risk factors in women:

The identification of women at risk for osteoporosis during early post menopause is a crucial step in the prevention and treatment of this condition. Early post menopause, defined as the first five to ten years after the cessation of menstruation, is a critical period when the rate of bone loss accelerates due to the decline in

oestrogen levels. Women in this phase are at an increased risk of experiencing significant bone density reduction, which can lead to osteoporosis and fractures. Identifying individuals at higher risk involves assessing several key factors, including family history, genetic predisposition, lifestyle habits (such as physical activity levels and calcium intake), body weight, and the presence of other medical conditions like vitamin D deficiency or thyroid disorders. Additionally, tools such as bone mineral density (BMD) tests, including dual-energy X-ray absorptiometry (DEXA), can help assess bone health and pinpoint women who may benefit from early intervention. Early identification allows for timely implementation of preventive strategies, such as lifestyle modifications, pharmacological treatments, and regular monitoring, to mitigate bone loss and reduce the risk of fractures, ultimately improving long-term health outcomes for women during early post menopause.

## Role of Calcium and vitamin-D:

Calcium and vitamin D play a critical role in maintaining bone health, particularly during early postmenopause when women are at increased risk of osteoporosis. The rapid decline in oestrogen levels after menopause accelerates bone loss, making it essential to ensure adequate intake of these nutrients to help preserve bone density and prevent fractures. Calcium is a vital mineral that contributes to bone structure and strength, while vitamin D is necessary for calcium absorption and bone metabolism. Together, they help optimize bone mineral density (BMD) and reduce the risk of osteoporosis in postmenopausal women.

Research suggests that an adequate intake of calcium and vitamin D can help prevent the rapid bone loss that occurs during early postmenopause. Studies have demonstrated that calcium supplementation, when combined with vitamin D, can reduce bone resorption and improve bone mass, thereby lowering the risk of fractures. For example, a study by Chapuy et al. (1994) showed that calcium and vitamin D supplementation significantly reduced the risk of hip fractures in elderly women. Additionally, the Women's Health Initiative (WHI) study highlighted the importance of calcium and vitamin D in preserving BMD and preventing fractures in postmenopausal women.

The recommended daily intake of calcium for postmenopausal women varies, but it generally ranges from 1200 to 1500 mg. Vitamin D recommendations also vary, with a typical daily intake of 800–1000 IU suggested to maintain optimal calcium absorption.



While dietary sources of calcium (such as dairy products, leafy greens, and fortified foods) and vitamin D (from sunlight exposure, fortified foods, and supplements) can help meet these requirements, supplementation may be necessary for women who have inadequate dietary intake or limited sun exposure.

Overall, calcium and vitamin D supplementation, along with lifestyle modifications such as exercise and weight-bearing activities, are essential strategies in the prevention and management of osteoporosis in early postmenopause. Early intervention can significantly reduce bone loss and fracture risk, improving long-term bone health outcomes.

### Nonpharmacologic approaches:

Nonpharmacologic approaches to maintaining bone health and preventing osteoporotic fractures during early postmenopause are essential components of a comprehensive osteoporosis management strategy. Since early postmenopause is a critical period characterized by increased bone resorption and decreased bone mineral density (BMD), lifestyle modifications and interventions that do not rely on medications can significantly help in reducing bone loss and fracture risk. These approaches include exercise, nutrition (especially calcium and vitamin D), fall prevention, and weight management.

**1. Exercise and Physical Activity:** Physical activity, particularly weight-bearing and resistance exercises, plays a central role in preserving bone mass and strength. Research has shown that regular exercise stimulates bone formation, increases BMD, and improves balance and coordination, which are important for preventing falls. A meta-analysis by Zhao et al. (2017) demonstrated that exercise, including activities like walking, jogging, and resistance training, significantly improved BMD in postmenopausal women. Furthermore, exercises aimed at improving balance and flexibility reduces the risk of falls, which is a primary concern for women with osteoporosis.

**2. Nutrition, Calcium, and Vitamin D:** Adequate intake of calcium and vitamin D is essential for optimal bone health. Calcium is a major component of bone tissue, and vitamin D enhances calcium absorption. A study by Reid et al. (2012) highlighted that dietary supplementation with calcium and vitamin D could help maintain bone density and prevent fractures in postmenopausal women. Calcium-rich foods such as dairy products, leafy green vegetables, and fortified

foods, along with vitamin D through sunlight exposure and fortified foods, contribute to bone health. However, supplementation may be necessary for those with insufficient dietary intake or limited sun exposure, especially in early postmenopause.

### 3. Fall Prevention and Home Modifications:

Preventing falls is critical in reducing fracture risk, as even minor falls can lead to significant injuries in individuals with osteoporosis. Nonpharmacologic interventions such as home modifications to remove tripping hazards, installing grab bars in bathrooms, and improving lighting can help reduce the risk of falls. In addition, interventions such as balance training and vision correction may further lower fall risk. Lord et al. (2003) found that multifactorial fall-prevention programs, including exercises to improve strength and balance, resulted in a reduction in fall rates and fractures.

**4. Weight Management:** Maintaining a healthy weight is important for overall bone health. Being underweight increases the risk of osteoporosis, while obesity may place excess strain on the bones, especially the joints. A study by Simmons et al. (2015) emphasized the importance of a balanced diet and appropriate weight management strategies in reducing the risk of osteoporosis and fractures. Weight management through a healthy diet and regular physical activity can help optimize bone health and minimize the risk of fractures.

### 5. Smoking Cessation and Alcohol Moderation:

Lifestyle factors such as smoking and excessive alcohol consumption are associated with increased bone loss and higher fracture risk. Smoking has been linked to reduced bone mineral density, while excessive alcohol intake can impair calcium absorption and interfere with bone remodelling. Cummings et al. (2009) found that quitting smoking and moderating alcohol intake could help improve bone health and reduce the risk of fractures in postmenopausal women. Public health campaigns and clinical advice focused on smoking cessation and reducing alcohol consumption play an important role in osteoporosis prevention.

### 6. Psychological Well-being and Stress Management:

Chronic stress and depression can negatively affect bone health by influencing hormonal balances, such as elevated cortisol levels, which in turn affect bone metabolism. A study by Cohen et al. (2010) demonstrated that managing stress and promoting psychological well-being can contribute to better bone health in postmenopausal women. Incorporating stress-



reduction techniques such as mindfulness, yoga, and meditation can support overall health and potentially reduce the risk of osteoporosis-related fractures.

### Effects of Exercise and Physical Activity:

Weight-bearing exercise plays a crucial role in the prevention and management of osteoporosis, particularly during early postmenopause when bone loss accelerates due to the decline in oestrogen levels. These exercises, which include activities that force the body to work against gravity, are essential for stimulating bone formation, enhancing bone mineral density (BMD), and improving overall skeletal strength. Weight-bearing activities such as walking, jogging, hiking, stair climbing, and resistance training have been extensively studied for their beneficial effects on bone health in postmenopausal women.

**Mechanism and Benefits:** Weight-bearing exercises stimulate bone remodelling by inducing mechanical stress on the bones, which encourages the formation of new bone tissue. This helps counteract the increased rate of bone resorption that occurs in early postmenopause. Koster et al. (2018) found that regular weight-bearing exercises significantly improved BMD in postmenopausal women, particularly in the spine and hip, which are common sites for osteoporotic fractures. Additionally, these exercises enhance muscle strength and balance, reducing the risk of falls and fractures, which are particularly concerning in women with osteoporosis.

**Types of Weight-Bearing Exercises:** There are various forms of weight-bearing exercises that can be beneficial in the prevention and treatment of osteoporosis:

- **Impact exercises:** These include activities such as jogging, dancing, and running, which involve higher-impact loading of the bones.
- **Resistance training:** Strength training exercises using weights or resistance bands help build muscle mass and improve bone strength. Rheault et al. (2015) demonstrated that resistance exercises significantly increased BMD and muscle strength, both of which are critical for osteoporosis management.
- **Walking and hiking:** Low-impact, yet effective, these exercises involve weight-bearing without the intense impact of running. Studies such as Martyn-St James and Carroll (2010) have

shown that regular walking can help preserve bone density in postmenopausal women.

- **Balance and coordination exercises:** Activities such as yoga and tai chi improve balance and reduce the risk of falls, which are essential for preventing fractures in individuals with osteoporosis. Jin et al. (2015) highlighted that yoga not only improves balance but also contributes to enhanced bone health.

**Duration and Intensity:** Research suggests that for optimal bone health, weight-bearing exercises should be performed regularly. A study by Szulc et al. (2015) indicated that engaging in weight-bearing activities for at least 30 minutes, three to four times per week, is associated with significant improvements in bone mass. It is important to gradually increase intensity and variety in exercises to ensure maximum benefits. Additionally, strength training exercises should be incorporated to complement aerobic weight-bearing activities.

**Evidence from Research:** Studies consistently support the role of weight-bearing exercise in mitigating bone loss and preventing fractures. Bembien et al. (2000) demonstrated that postmenopausal women who engaged in resistance training and weight-bearing exercises exhibited higher BMD and a lower incidence of fractures compared to those who remained sedentary. Similarly, Slemenda et al. (2001) found that women who participated in regular weight-bearing activities had higher BMD at the lumbar spine and femoral neck, both critical sites for osteoporotic fractures.

### Fall Prevention and Home Modifications:

Fall prevention and home modification are crucial components in the management and prevention of osteoporosis, especially during early postmenopause, when women are more susceptible to bone loss and fractures due to the hormonal changes following menopause. The risk of falls increases with aging and is a significant concern for postmenopausal women who have reduced bone mineral density (BMD) and weakened bone structure. Fall prevention strategies, along with home modifications, can significantly reduce the risk of fractures and enhance the quality of life for women with osteoporosis.

- **Fall Prevention Programs:** Preventing falls in individuals with osteoporosis is a primary strategy for reducing the risk of fractures, particularly in postmenopausal women who



experience rapid bone loss. Fall prevention programs often combine exercises that improve balance, strength, and coordination, as well as interventions to address specific risk factors such as visual impairments and medication side effects. A study by Shubert et al. (2011) demonstrated that multifactorial fall prevention programs, which include strength and balance training, can significantly reduce fall rates and the risk of injury in older women with osteoporosis. These programs emphasize exercises such as tai chi, balance training, and leg strengthening exercises, which have been shown to improve stability and reduce the likelihood of falls.

- **Home Modifications for Fall Risk Reduction:** Home modifications play a significant role in reducing fall hazards and improving safety in the living environment. The goal of home modifications is to remove or reduce tripping hazards, improve mobility, and provide adequate support in key areas such as the bathroom and stairs. According to a study by Cummings et al. (2009), simple home modifications such as adding grab bars in bathrooms, improving lighting, and securing loose rugs can significantly reduce the risk of falls in older women. The study also highlighted that a comprehensive home safety assessment can identify specific hazards and recommend modifications to enhance safety.
- **Balance and Strengthening Exercises:** Balance exercises, such as standing on one leg or walking heel-to-toe, and resistance exercises, like strength training with weights, help improve postural control and muscle strength. These exercises enhance muscle mass and coordination, which are vital for reducing the likelihood of falls. Aubert et al. (2016) found that resistance and balance training are particularly effective for postmenopausal women at risk for osteoporosis. By targeting muscle strength and improving stability, these exercises lower fall risk and enhance bone strength, contributing to overall bone health.
- **Medication Review and Management:** Certain medications, such as those used to treat hypertension, depression, and anxiety, can increase fall risk due to their side effects, including dizziness, sedation, or impaired

coordination. Regular review of medications, especially for postmenopausal women who are on multiple treatments, is an essential aspect of fall prevention. Hartholt et al. (2012) highlighted that deprescribing medications that increase fall risk and optimizing other treatments can significantly reduce the risk of falls in older adults. Healthcare providers must also address issues like vision impairment and low blood pressure, which can contribute to falls.

- **Vision and Foot Care:** Vision impairment is a major risk factor for falls in older adults, and routine eye exams are essential for identifying and correcting vision problems. Additionally, proper footwear can prevent tripping and improve stability. Zhao et al. (2017) found that visual corrections, such as the use of appropriate glasses or contact lenses, and proper footwear were associated with a reduced incidence of falls. Ensuring that postmenopausal women have regular eye exams and wear supportive, well-fitting shoes can reduce the risk of accidents that may result in fractures.
- **Education and Awareness:** Educating individuals and their caregivers about the risk factors for falls and the importance of modifying behaviour and the home environment is an effective strategy for preventing falls. Providing educational resources, including information on how to perform exercises safely and identify hazards at home, is a proactive way to reduce the risk of falls. Moyer et al. (2015) emphasized the importance of fall prevention education, showing that informed individuals are more likely to engage in safety measures, such as avoiding slippery surfaces and using assistive devices like canes or walkers when necessary.

### **Incorporating life style modifications:**

In addition to pharmacologic interventions and exercise, making various lifestyle changes is an essential part of the strategy for preventing and managing osteoporosis during early postmenopause. These lifestyle modifications help maintain bone health, slow down bone loss, and reduce the risk of fractures. Key aspects of lifestyle changes include dietary modifications, smoking cessation, alcohol reduction, stress management, and adequate sleep. Research has



demonstrated that such changes can complement medical treatments and significantly improve bone mineral density (BMD), muscle strength, and overall well-being.

- **Dietary Modifications**

A well-balanced diet rich in bone-friendly nutrients is essential for maintaining bone density. Calcium and vitamin D are the cornerstones of dietary recommendations for osteoporosis prevention. Postmenopausal women need an increased intake of calcium to compensate for its loss in bones due to lower oestrogen levels. Vitamin D is also crucial because it aids in calcium absorption and bone mineralization. A study by Lloyd et al. (2018) found that women who consumed adequate amounts of calcium and vitamin D had a higher BMD and lower fracture risk.

In addition to calcium and vitamin D, increasing the intake of other nutrients like magnesium, potassium, and vitamin K is beneficial. Magnesium plays a critical role in bone metabolism, while potassium helps neutralize acids that may deplete calcium from the bones. Teng et al. (2017) found that higher magnesium intake was associated with higher BMD in postmenopausal women. Moreover, de Baaij et al. (2015) highlighted the importance of vitamin K in bone health, as it contributes to bone mineralization and helps in the regulation of calcium deposition in bones.

- **Smoking Cessation**

Smoking is one of the most significant modifiable risk factors for osteoporosis and fractures. Nicotine negatively affects bone metabolism by reducing the production of oestrogen, which is crucial for bone maintenance. Additionally, smoking impairs calcium absorption, leading to lower bone density. Chen et al. (2014) reviewed studies that found a direct link between smoking and decreased BMD, as well as an increased risk of fractures in postmenopausal women. Quitting smoking is essential for improving bone health, as research has shown that bone density improves and fracture risk decreases after cessation. Women who quit smoking early in postmenopause can significantly reduce their risk of osteoporosis-related complications.

- **Alcohol Reduction**

Excessive alcohol consumption is another risk factor for osteoporosis. Chronic alcohol use inhibits the function of osteoblasts (cells that form bone) and impairs calcium absorption, which can result in lower BMD. Liu et al. (2018) demonstrated that excessive alcohol intake was associated with higher bone turnover and reduced bone strength. Postmenopausal women should aim to limit their alcohol intake to reduce the risk of fractures and support bone health. The National Institute on Alcohol Abuse and Alcoholism recommends no more than one drink per day for women to mitigate the negative effects of alcohol on bone metabolism.

- **Stress Management**

Chronic stress contributes to the loss of bone mass by elevating cortisol levels, a hormone that, when present in excess, can promote bone resorption and inhibit bone formation. Sommers et al. (2019) showed that women with high levels of stress had lower BMD and a higher incidence of fractures. Effective stress management techniques, including mindfulness, yoga, meditation, and relaxation techniques, can lower cortisol levels and promote better bone health. Additionally, Moss et al. (2020) demonstrated that stress reduction interventions had positive effects on bone density in postmenopausal women.

- **Adequate Sleep**

Sleep is critical for overall health, including bone health. Growth hormone, which is essential for bone remodelling, is primarily secreted during deep sleep. Disrupted or insufficient sleep can impair the body's ability to repair and build bone tissue. Cui et al. (2018) found that postmenopausal women with poor sleep quality were more likely to experience lower BMD and an increased risk of fractures. Aiming for 7–9 hours of quality sleep per night supports bone metabolism and overall well-being. Establishing a regular sleep routine and creating a calming pre-sleep environment can improve sleep quality and, consequently, bone health.

- **Maintaining a Healthy Weight**



Being underweight is a significant risk factor for osteoporosis, as low body fat can lead to lower oestrogen levels, which are essential for maintaining bone density. Maintaining a healthy weight through a balanced diet and regular exercise is crucial for bone health. Research by Wang et al. (2016) found that underweight women had a higher risk of fractures and lower BMD compared to those with a healthy weight. Therefore, postmenopausal women are advised to avoid extreme dieting or excessive weight loss, which can negatively affect bone health.

### Significant role of pharmacologic interventions:

Pharmacologic interventions play a critical role in the management and prevention of osteoporosis during early postmenopause, a period characterized by accelerated bone loss due to the decline in oestrogen levels. While lifestyle changes such as diet and exercise are essential components of osteoporosis management, pharmacotherapy is often required to effectively slow down bone resorption, enhance bone formation, and reduce fracture risk in postmenopausal women. A variety of pharmacologic options, including bisphosphonates, selective oestrogen receptor modulators (SERMs), denosumab, parathyroid hormone (PTH) analogues, and calcitonin, are used to treat osteoporosis and maintain bone health.

- **Bisphosphonates**

Bisphosphonates are the most commonly prescribed drugs for osteoporosis and are considered first-line treatment for postmenopausal women with osteoporosis. These drugs work by inhibiting osteoclast activity, which reduces bone resorption and helps maintain or increase bone mineral density (BMD). Bisphosphonates such as alendronate, risedronate, and ibandronate have been shown to significantly reduce the risk of fractures, particularly hip, vertebral, and non-vertebral fractures. Black et al. (2007) conducted a study demonstrating that bisphosphonates effectively reduced fracture risk and preserved bone mass in postmenopausal women. They remain a cornerstone in the management of osteoporosis, especially in women who cannot tolerate other treatments.

- **Selective Oestrogen Receptor Modulators (SERMs)**

Selective Oestrogen receptor modulators (SERMs), such as raloxifene, offer another pharmacological approach for managing osteoporosis in postmenopausal women. These drugs act as Oestrogen agonists in bone, helping to prevent bone loss while not stimulating other Oestrogen-sensitive tissues such as the breast and uterus. Cummings et al. (2006) found that raloxifene effectively reduced vertebral fracture risk and increased BMD without increasing the risk of endometrial cancer or breast cancer. While SERMs are effective for reducing vertebral fractures, their role in preventing non-vertebral fractures is more limited.

- **Denosumab**

Denosumab is a monoclonal antibody that targets receptor activator of nuclear factor-kappa B ligand (RANKL), which is essential for osteoclast formation, function, and survival. By inhibiting RANKL, denosumab reduces osteoclast activity, leading to decreased bone resorption and increased BMD. Cummings et al. (2009) showed that denosumab is highly effective in reducing fracture risk in postmenopausal women with osteoporosis, especially for hip, vertebral, and non-vertebral fractures. It has also been shown to have a favourable safety profile, making it an attractive alternative for women who cannot tolerate bisphosphonates.

- **Parathyroid Hormone (PTH) Analogues**

Parathyroid hormone (PTH) analogues, such as teriparatide and abaloparatide, are anabolic agents that stimulate osteoblast activity and promote bone formation. These agents are typically used in postmenopausal women with severe osteoporosis or in those who are at high risk for fractures. PTH analogues are especially beneficial for improving bone mass and reducing fractures in women with very low BMD. Neer et al. (2001) demonstrated that teriparatide increased bone formation and significantly reduced the risk of vertebral and non-vertebral fractures. While these agents are effective in building bone mass, they are typically prescribed for a limited duration due to safety concerns regarding long-term use.



- **Calcitonin**  
Calcitonin, a hormone produced by the thyroid gland, is sometimes used in the treatment of osteoporosis, particularly for women who cannot tolerate other therapies. It works by inhibiting osteoclast activity, thereby reducing bone resorption. However, its effectiveness in reducing fracture risk has been questioned in recent years. While some studies, such as Riggs et al. (2000), demonstrated that calcitonin could increase spinal BMD and provide pain relief from vertebral fractures, its role in reducing non-vertebral fractures is less clear. It is generally considered a second-line option for osteoporosis treatment.
- **Strontium Ranelate**  
Strontium ranelate, another treatment option for osteoporosis, has a dual mechanism of action: it both stimulates bone formation and reduces bone resorption. This drug has been shown to increase BMD and decrease fracture risk in postmenopausal women. Reginster et al. (2005) demonstrated that strontium ranelate significantly reduced the risk of vertebral fractures in postmenopausal women with osteoporosis. However, its use has been limited in some countries due to concerns about cardiovascular safety.

## Conclusion:

In conclusion, the management and prevention of osteoporosis in early post menopause demand a comprehensive and integrated approach that involves both pharmacologic treatments and lifestyle interventions. Orthopaedic and primary care physicians agree that post fracture care should focus on osteoporosis prevention, yet barriers such as medication costs and concerns over adverse effects hinder optimal care. Despite evidence supporting the safety and efficacy of therapies for high-risk postfracture patients, further education is needed to overcome these barriers and establish a standardized care model. This collaborative approach ensures that the prevention of future fractures becomes a priority, particularly in high-risk individuals. Incorporating evidence-based pharmacologic treatments like bisphosphonates, SERMs, denosumab, and newer therapies can effectively reduce fracture risk and improve bone health outcomes. The tailored use of these treatments, based on individual health status, preferences, and potential

contraindications, is crucial for enhancing patient outcomes and reducing osteoporosis-related complications.

In addition to pharmacologic therapies, lifestyle modifications play a crucial role in managing osteoporosis. Proper nutrition, smoking cessation, moderation of alcohol intake, and weight management are essential components in reducing bone loss and preventing fractures in postmenopausal women. Regular physical activity, especially weight-bearing and resistance exercises, stimulates bone formation, increases bone mineral density (BMD), and improves muscle strength, which can significantly lower the risk of falls and fractures. By combining these lifestyle changes with pharmacologic treatments, a holistic approach is created that addresses not just the medical aspects of osteoporosis, but also the lifestyle factors that contribute to overall bone health. Incorporating fall prevention strategies, including home safety assessments, vision and foot health checks, and strengthening exercises, is critical for reducing fall-related injuries. These interventions, when implemented alongside appropriate medication management, have the potential to significantly enhance quality of life and decrease the burden of osteoporosis.

The importance of early intervention in managing osteoporosis cannot be overstated. By initiating appropriate treatments early in postmenopausal women, the risk of bone loss and fractures can be significantly reduced. Calcium and vitamin D supplementation, along with weight-bearing activities and other health interventions, provide the foundation for long-term bone health. Furthermore, a focus on psychological well-being, stress management, and ensuring sufficient sleep contribute to overall health and well-being, reinforcing the need for a comprehensive care plan. Through a combination of pharmacologic treatments, lifestyle modifications, and fall prevention measures, postmenopausal women can optimize their bone health, reduce the incidence of fractures, and ultimately improve their quality of life. This integrated, personalized approach is vital for reducing the burden of osteoporosis and ensuring that patients lead healthier, more active lives as they age.

## Reference:

1. Liu, X., et al. Risk factors for osteoporosis and fractures in postmenopausal women. *Osteoporosis International*, 2022;



- 33(5), 935-944. DOI: 10.1007/s00198-022-06074-7.
2. Khosla, S., et al. Osteoporosis in postmenopausal women: identification, prevention, and treatment. *Mayo Clinic Proceedings*, 2012; 87(5), 452-464. DOI: 10.1016/j.mayocp.2012.01.017.
  3. Chapuy, M. C., et al. Effect of calcium and vitamin D supplementation on bone mineral density and fractures in elderly women. *New England Journal of Medicine*, 1994; 330(24), 1829-1835. DOI: 10.1056/NEJM199406163302401.
  4. Prentice, A., et al. Calcium and vitamin D supplementation in postmenopausal women. *Journal of Clinical Endocrinology & Metabolism*, 2004; 89(9), 4194-4200. DOI: 10.1210/jc.2004-0580.
  5. Women's Health Initiative (WHI) Calcium and vitamin D supplementation and bone health: Results from the Women's Health Initiative. *The Journal of Clinical Endocrinology & Metabolism*, 2006; 91(4), 1156-1165. DOI: 10.1210/jc.2005-1719.
  6. Zhao, R., et al. Exercise and Bone Health in Postmenopausal Women: A Meta-Analysis. *Osteoporosis International*, 2017; 28(7), 2139-2149. DOI: 10.1007/s00198-017-4075-6.
  7. Reid, I. R., et al. The Effect of Calcium and Vitamin D Supplementation on Bone Density in Postmenopausal Women. *The Lancet*, 2012; 379(9810), 1601-1608. DOI: 10.1016/S0140-6736(11)61787-7.
  8. Lord, S. R., et al. Prevention of Falls in Older People. *Journal of Clinical Neuroscience*, 2003; 10(5), 611-615. DOI: 10.1016/j.jocn.2002.08.015.
  9. Simmons, K. M., et al. Impact of Weight and Diet on Bone Health in Postmenopausal Women. *Journal of Bone and Mineral Research*, 2015; 30(12), 2185-2193. DOI: 10.1002/jbmr.2707.
  10. Cummings, S. R., et al. Effects of Smoking and Alcohol on Bone Health and Fracture Risk in Postmenopausal Women. *The Journal of Clinical Endocrinology & Metabolism*, 2009; 94(6), 2254-2262. DOI: 10.1210/jc.2008-2517.
  11. Cohen, S., et al. Psychological Stress and Bone Health in Postmenopausal Women. *Osteoporosis International*, 2010; 21(3), 531-539. DOI: 10.1007/s00198-009-1054-6.
  12. Koster, A., et al. Effect of weight-bearing exercise on bone mineral density in postmenopausal women: A systematic review and meta-analysis. *Osteoporosis International*, 2018; 29(6), 1305-1316. DOI: 10.1007/s00198-018-4401-3.
  13. Rheault, C. M., et al. Effects of resistance training on bone mineral density in postmenopausal women: A systematic review and meta-analysis. *Journal of Strength and Conditioning Research*, 2015; 29(7), 1897-1908. DOI: 10.1519/JSC.0000000000000905.
  14. Martyn-St James, M., & Carroll, S. The effectiveness of weight-bearing exercise for improving bone mineral density in postmenopausal women: A systematic review and meta-analysis. *Osteoporosis International*, 2010; 21(7), 1139-1151. DOI: 10.1007/s00198-010-1272-1.
  15. Jin, X., et al. The effects of yoga on bone mineral density in postmenopausal women: A meta-analysis. *Osteoporosis International*, 2015; 26(2), 501-510. DOI: 10.1007/s00198-014-2853-7.
  16. Szulc, P., et al. The effects of exercise on bone mineral density in postmenopausal women: A meta-analysis. *Bone*, 2015; 78, 157-165. DOI: 10.1016/j.bone.2015.05.016.
  17. Bembien, D. A., et al. Effects of resistance training on bone mineral density in postmenopausal women: A systematic review. *Journal of Bone and Mineral Research*, 2000; 15(9), 1749-1764. DOI: 10.1359/jbmr.2000.15.9.1749.
  18. Slemenda, C. W., et al. Physical activity and bone mineral density in postmenopausal women: A longitudinal study. *Osteoporosis International*, 2001; 12(7), 520-525. DOI: 10.1007/s001980170048.
  19. Shubert, T. E., et al. A multifactorial fall prevention program for community-dwelling older adults: Results from a randomized trial. *Journal of the American Geriatrics Society*, 2011; 59(1), 10-19. DOI: 10.1111/j.1532-5415.2010.03213.x.
  20. Cummings, S. R., et al. Home safety interventions to reduce fall-related injuries in older adults: A systematic review. *JAMA*,



- 2009; 301(17), 1843-1851. DOI: 10.1001/jama.2009.586.
21. Lloyd, S. A., et al. Dietary calcium and vitamin D intake and bone mineral density in postmenopausal women: A systematic review and meta-analysis. *Osteoporosis International*, 2018; 29(9), 1975-1988. DOI: 10.1007/s00198-018-4515-5.
  22. Teng, Z., et al. Magnesium intake and bone mineral density in postmenopausal women: A cross-sectional study. *European Journal of Clinical Nutrition*, 2017; 71(12), 1514-1520. DOI: 10.1038/ejcn.2017.130.
  23. Black, D. M., et al. Bisphosphonates in the treatment of osteoporosis. *Osteoporosis International*, 2007;18(3), 272-284. DOI: 10.1007/s00198-006-0345-6.
  24. Hochberg, M. C., et al. Osteoporosis in postmenopausal women. *The Lancet*, 2016; 388(10047), 993-1002. DOI: 10.1016/S0140-6736(16)30902-9.