



## Association between Index of Central Obesity and Lipid Profile in Obese Type 2 Diabetes

P Raja Reddy<sup>1</sup>, Jayani Reddy PR<sup>2</sup>

<sup>1</sup>Associate Professor, Department of Physiology, The Apollo Medical College, Chittoor, Andhra Pradesh; <sup>2</sup>M.B;B.S graduate from Vydehi Institute of Medical Sciences and Research ,Bangalore, Karnataka, India.

### Address for Correspondence:

P. Raja Reddy, Associate Professor, Department of Physiology, Apollo Medical College, Chittoor, AP.

(Received: 16 January 2025

Revised: 20 February 2025

Accepted: 31 March 2025)

### KEYWORDS

Type two diabetes mellitus, Waist hip ratio, Lipid profile

### ABSTRACT:

**Background:** The Index of central obesity (ICO) is defined as a ratio of waist circumference and height and is a better parameter of central obesity. To quantify proportion of visceral fat in the total body fat, waist circumference alone is not sufficient.

**Objective:** To observe the relation between ICO and lipid profile in obese type 2 diabetes mellitus.

**Materials and Methods:** The present study was carried out at rural based clinic located near to Apollo Medical College region, Chittoor, Andhra Pradesh, after a written consent from all the participants. The study includes 515 diabetic patients, in which 356 were males and 159 were females. The Index of central obesity were measured by taking the waist circumference and height and about 5 ml of blood was collected and used for the estimation of serum cholesterol, triglyceride and HDL-Cholesterol levels by using standard methods. Statistical analysis was done by using independent student t test and Pearson Correlation coefficient was calculated.

**Results:** The ICO and lipid profiles were not significantly correlated. In type 2 diabetes male patients showed higher triglycerides ( $177.96 \pm 100.19$ ) with waist hip ratio  $>0.9$ , whereas female patients showed less Triglycerides ( $178.19 \pm 99.52$ ) with waist hip ratio  $>0.8$ . However these differences were statistically not significant.

**Conclusion:** Multiple anthropometric parameters are required to correlate lipid profile rather than single parameter in type 2 diabetes mellitus. Along with anthropometric measurements, lipid profile is also need to be monitored in type 2 diabetics.

### INTRODUCTION

Defining central obesity, which is a single entity numerous WC cutoffs have been suggested, separately for males and females and various races. We believe that this difference is amendable to differences in their average heights. To quantify proportion of visceral fat in the total body fat, WC alone is not sufficient. Authors hereby hypothesis that, Index of central obesity defined by Rakeshet al<sup>3</sup>, as a ratio of WC and height is a better parameter of central obesity. National Health and nutrition Examination Survey data base is replacing

index of central obesity with waist circumference was found to enhance the specificity and sensitivity of definition of metabolic syndrome<sup>8</sup>. Therefore, the present study was aimed to assess index of central obesity (ICO) and glycemic control among type 2 diabetic patients.

Type 2 Diabetes mellitus (T2DM) is characterized by three pathophysiologic abnormalities: impaired insulin secretion, peripheral insulin resistance, and excessive hepatic glucose production. Obesity, particularly visceral or central (as evidenced by the waist-hip ratio),



is very common in type 2 Diabetes mellitus.<sup>1</sup> Alterations in body fat distribution are associated with changes in lipids and lipoproteins and with increased coronary heart disease (CHD).<sup>2</sup> Individuals with diabetes mellitus may have several forms of dyslipidemia. In obese patients with T2DM a distinct “diabetic dyslipidemia” is characteristic of the insulin resistance syndrome. Its features are a high serum triglyceride level (300-400 mg/dl), a low HDL-cholesterol (less than 30 mg/dl), and a qualitative change in LDL particles. Measures designed to correct the obesity and hyperglycemia, such as exercise, diet and hypoglycemic therapy, are the treatment of choice for diabetic dyslipidemia.<sup>3</sup> In type 2 diabetes mellitus patients who are centrally obese, increased lipolysis causes the liver to increase glucose and very low-density lipoprotein output, while muscle uses less. This leads to a rise in blood glucose and triglycerides, a drop in HDL cholesterol, and an increase in small, dense LDL particles.<sup>4</sup> The present study was undertaken to establish the relation between waist-hip ratio and lipid profile in type 2 diabetes mellitus patients.

## MATERIALS AND METHODS

The present study was carried out at rural based clinic located near to Apollo Medical College region, Chittoor, Andhra Pradesh, after a written consent from all the participants. The study includes 515 diabetic patients, in which 356 were males and 159 were females. The circumference of the waist and the hip were measured and the ratio was taken as Waist-Hip Ratio (WHR). Internationally accepted cut of points for WHR are  $>0.90$  for men and  $>0.80$  for women (National Cholesterol Education Program, 1994)<sup>5</sup> were considered. Male diabetic patients were divided into Group 1 with 49 patients with  $WHR < 0.90$  and Group 2 with 287 patients with  $WHR > 0.90$ . Female diabetic patients were divided into Group 1 with 7 patients with  $WHR < 0.80$  and Group 2 with 162 patients with  $WHR > 0.80$ . About 5 ml of fasting blood sample was collected and used for the estimation of serum cholesterol, triglyceride and HDL- Cholesterol levels using standard methods. Major selection criteria for diabetes included: a random plasma glucose level of 200mg/ dl or greater when the symptoms of diabetes were present and fasting plasma glucose level of 126 mg/dl or greater. Data were expressed as Mean  $\pm$  SD. Statistical analysis was done by using independent student ‘t’ test and Pearson Correlation coefficient was calculated and P value was taken as significant at 5 percent confidence level ( $P < 0.05$ ).

## RESULTS

Total 515 type 2 diabetics were studied out of which 336 male and 169 were female patients. The base line features of the patients were shown in Table 1. The mean serum cholesterol, triglyceride and HDL-cholesterol levels of male type-2 diabetes mellitus patients with WH ratio  $<0.90$  were 162.61 mg/dl, 163.12mg/dl and 37.46mg/dl respectively, and mean serum cholesterol, triglyceride and HDL-cholesterol levels of male type-2 diabetes mellitus patients with WH ratio  $>0.90$  were 161.70 mg/dl, 177.96 mg/dl and 37.50mg/dl respectively. Table 2 and Figure 1 shows mean values of cholesterol, triglycerides and HDL cholesterol in relation to WHR in male type 2 diabetics. The mean serum cholesterol, triglyceride and HDL-cholesterol levels of female type-2 diabetes mellitus patients with WH ratio  $<0.80$  were 184.14 mg/dl, 188.28 mg/dl and 35 mg/dl respectively, and mean serum cholesterol, triglyceride and HDL-cholesterol levels of female type-2 diabetes mellitus patients with WH ratio  $>0.80$  were 180.41 mg/dl, 178.19 mg/dl and 37.88 mg/dl respectively. Table 3 and Figure 2 shows mean values of cholesterol, triglycerides and HDL cholesterol in relation to WHR in female type 2 diabetics.

## DISCUSSION

Accelerated coronary and peripheral vascular atherosclerosis is one of the most common and serious complications of long term diabetes mellitus.<sup>6</sup> Along with other risk factors including hypertension, smoking and obesity, increasing importance has been given to secondary hyperlipidemia in the causation of accelerated atherosclerosis.<sup>7</sup> Obesity is a positive risk factor in the development of type 2 Diabetes mellitus, dyslipidemia, insulin resistance and hypertension.<sup>8</sup> Obesity is often expressed in terms of body mass index (BMI).<sup>9</sup> The distribution of adipose tissue in different anatomic depots also has substantial implications for morbidity. Specifically, intra-abdominal and abdominal subcutaneous fat has more significance than subcutaneous fat present in the buttocks and lower extremities. Determining the waist-to-hip ratio, most easily makes this distinction.<sup>10</sup> The risk of diabetes increases progressively with increasing body mass index and waist-hip ratio. Weight gain is associated with an increase in insulin resistance and deterioration in glucose tolerance. Mainly the centrally located adipocytes have specific metabolic roles in the pathogenesis of insulin resistance and type 2 diabetes mellitus.<sup>11</sup> NarasimhaRaiet al<sup>12</sup> found that, as Waist-Hip ratio increases, serum cholesterol and triglyceride levels increases in male type 2 diabetes mellitus patients. Haffner SM et al.<sup>13</sup>, in 1987 assessed diabetes



and cardiovascular risk factors in Mexican-Americans and found that Waist-Hip ratio was associated with type 2 diabetes mellitus rates, low HDL-cholesterol levels and high triglyceride levels. Buynes C et al.,<sup>14</sup> studied the sex differences in fat distribution, WH ratio, serum lipids, and blood pressure, in male and female patients with type 2 diabetes mellitus, and found that men had higher WH ratio and lower HDL-cholesterol. Samir B Al - Mukhtar et al<sup>15</sup> found that , obese diabetics, when compared to non- obese diabetics, had significant increase in the levels of serum total cholesterol, triglycerides, LDL-C and VLDL-C while serum HDL-C differs significantly.

### CONCLUSION

In the present study we found that waist hip ratio is not significantly associated with abnormal lipid parameters in both male and female type 2 diabetes mellitus. Multiple anthropometric parameters are required to correlate lipid profile rather than single parameter in type 2 diabetes mellitus. Along with anthropometric measurements, lipid profile is also need be monitored in type 2 diabetics.

### ACKNOWLEDGEMENT

We acknowledge the study participants cooperation.

**Table 1: Base line features of patients**

| Parameter                 | Mean ±SD     |
|---------------------------|--------------|
| Age ( years)              | 55.38±11.60  |
| Height (cm)               | 162.91±8.87  |
| Weight ( Kg)              | 65.68±11.17  |
| BMI (Kg/m <sup>2</sup> )  | 24.69±4.23   |
| Waist circumference (cm)  | 94.95±10.21  |
| ICO                       | 0.84± 0.25   |
| Hip circumference (cm)    | 94.89±10.14  |
| WHR                       | 1.01±0.12    |
| Total cholesterol (mg/dl) | 168.11±38.75 |
| Triglycerides( mg/dl)     | 176.70±98.28 |
| HDL-cholesterol( mg/dl)   | 37.59±5.34   |
| LDL-cholesterol( mg/dl)   | 94.58±36.84  |

**Table 2: WHR/ ICO and lipid profile in male type 2 diabetic patients**

| Lipid parameter (mg/dl) | WHR≤0.90 (N=49) | WHR>0.90 (N=287) | *P value |
|-------------------------|-----------------|------------------|----------|
| Cholesterol             | 162.61±33.05    | 161.70±36.08     | 0.948    |
| Triglyceride            | 163.12±75.86    | 177.96±100.19    | 0.323    |
| HDL-cholesterol         | 37.46±4.19      | 37.50±6.59       | 0.974    |

\*P= <0.05 considered significant

**Table 3: WHR and Lipid profile in female type 2 diabetic patients**

| Lipid parameter (mg/dl) | WHR≤0.80 (N=162) | WHR>0.80 (N=162) | *P value |
|-------------------------|------------------|------------------|----------|
| Cholesterol             | 184.14±39.37     | 180.41±41.80     | 0.954    |
| Triglyceride            | 188.28±139.16    | 178.19±99.52     | 0.120    |
| HDL-cholesterol         | 35.00±4.93       | 37.88±5.12       | 0.627    |

\*P= <0.05 considered significant




Fig. 1 ICO and Lipid profile in selected male subjects

Fig. 2 ICO and lipid profile in selected female subjects

### REFERENCES

1. Parikh R, Mohan v, Joshi S Should waist circumference be replaced by Index of central obesity in definition of metabolic



- syndrome?, Medical hypotheses, 2012, Jan 28(1): 3-5.
2. Powers AC. Diabetes Mellitus. In: Kasper DL (ed). Harrison's Principles of Internal medicine. 16th ed. Mc Graw-Hill companies. Inc 2005;II:2152-2180.
  3. WHO study group on diabetes mellitus. World health organization technical report series 1985;727.
  4. Masharani U. Diabetes Mellitus and Hypoglycemia. In: Tierney LM, McPhee SJ, Papadakis MA (ed). Current Medical Diagnosis and Treatment. 44th ed. Mc Graw-Hill Companies, Inc 2005;1157-1190.
  5. Pi-Sunyer Fx. The epidemiology of central fat distribution in relation to disease. Nutr Rev 2004;62(7 Pt 2):S120-S126.
  6. National Cholesterol Education Program, (1994). Second report of the expert panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult treatment panel II). Circulation, 89:1333-1445.
  7. Zargar, AH Wandroo FA, Wadhwa MB, Laway BA, Masoodi SR and Shah NA. Serum lipid profile in non-insulin dependent diabetes mellitus associated with obesity. Int J Diab Dev Countries 1995;5:9-13.
  8. Dunn FL. Treatment of lipid disorders in diabetes mellitus. Med Clin North America 1988;72(6):1379-1398.
  9. Park K. Obesity. In: Park's textbook of Preventive and Social Medicine. 18th ed. M/S Banarsidas Bhanot: Jabalpur (India): 2005;316-319.
  10. Park K. Diabetes Mellitus. In: Park's textbook of Preventive and Social Medicine. 18th ed. M/S Banarsidas Bhanot: Jabalpur (India): 2005;311-316.
  11. Flier JS, Flier EM. Obesity. In: Kasper DL (ed). Harrison's Principles of Internal medicine. 16th ed. Mc Graw-Hill Companies, INC. 2005;Vol II:422-429.
  12. Carey VJ, Walters EE, Colditz GA, Solomon CG, Willett WC, Rosner BA, et al. Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women. The Nurses' Health Study. Am J Epidemiol 1997;145:614-619.
  13. Narasimha Rai K and Jeganthan PS. Relation between Waist-Hip Ratio and Lipid Profile in Male Type 2 Diabetes Mellitus Patients. October-December 2012 RJPBCS Vol 3(4):954-958.
  14. Haffner SM, Stern MP, Hazuda HP, Pugh J and Patterson JK. Do upper-body and centralized adiposity measure different aspects of regional body-fat distribution? Relationship to non-insulin-dependent diabetes mellitus, lipids, and lipoproteins. Diabetes 1987;36(1):43-51.
  15. Baynes C, Henderson AD, Anyaoku V, Richmond W, Johnston DG and Elkeles RS. The Influence of Regional Adiposity on Atherogenic Risk Factors in Men and Women with Type 2 Diabetes. Diabet Med 1991;8(5):458-463.
  16. Samir B Al - Mukhtar, Raad Y. Al- Hamdani and Amjad H. Al-Naemi. Obesity and lipid profile in type 2 diabetics. Tikrit Medical Journal 2006;12(1):15-21.