



## Assessment of Drinking Water Quality in Palanpur Division

Bharti Patel<sup>1</sup>, Dishaben A Ranpura<sup>1</sup>, Narendra Kumar Sharma\*<sup>1</sup>, Jigar Y Soni<sup>1</sup>

<sup>1</sup>Department of Chemistry, Faculty of Basic and Applied Sciences, Madhav University, Pindwara, Sirohi (Rajasthan), India 307032

(Received: 16 July 2025

Revised: 20 August 2025

Accepted: 02 September 2025)

### KEYWORDS

Groundwater quality, Physicochemical parameters, Biological parameters, Palanpur Division, Fluoride contamination, Total Dissolved Solids (TDS), Water hardness, Bureau of Indian Standards (BIS 10500:2012)

### ABSTRACT:

Through the analysis of triplicate samples taken from boreholes in nine chosen villages in the Palanpur Division of Gujarat, this study evaluates the groundwater quality in those villages. pH, fluoride, total dissolved solids (TDS), electrical conductivity, turbidity, chloride, total hardness (TH), calcium, magnesium, total alkalinity, nitrate, sulfate, dissolved oxygen (DO), biological oxygen demand (BOD), and chemical oxygen demand (COD) were among the important physicochemical and biological parameters that were assessed. The findings showed spatial variability, with multiple samples surpassing the WHO, TDS, and fluoride limits as well as the BIS (10500:2012) standards. Villages like Madana and Gadhadra had extremely hard water and TDS levels above 1000 mg/L. In several locations, fluoride levels were close to 1 mg/L, which could be harmful to health. Localized organic contamination was indicated by elevated BOD and COD levels. The results emphasize that in order to guarantee drinkable groundwater and long-term water security in the area, routine monitoring, defluoridation units, water softening, and community-level interventions are required.

### 1. Introduction

In Palanpur Division, Banaskantha, Gujarat, where there is little rainfall (600–700 mm per year), over 85% of the 3.1 million inhabitants rely on groundwater as their main resource for agriculture, drinking, and livestock (Census of India, 2011). Geochemical processes and anthropogenic inputs pose a threat to quality; the main contaminants are fluoride, nitrates, salts, and pollutants from industry, agriculture, and inadequate waste management (Adimalla & Venkatayogi, 2023; Shirke et al., 2020; Zhu et al., 2022). Fluorosis is caused by elevated fluoride (>1.5 mg/L) (Pradhan & Biswal, 2018; Shirke et al., 2020), and usability is restricted by high TDS and hardness (Jha et al., 2021; Vyas et al., 2019). High BOD and COD are indicators of organic pollution from both industrial and residential sources (CPCB, 2021; Kumar et al., 2022). There is still little monitoring (CGWB, 2022; UN-Water, 2023). In order to compare the results with the BIS (2023) and WHO (2022) standards, this study examined 27 samples from nine villages for the following parameters: pH, fluoride, TDS, EC, turbidity, chloride, TH, calcium, magnesium, alkalinity, nitrate, sulfate, DO, BOD, and COD. The results demonstrate spatial variability, with localized organic contamination (CPCB, 2021; Kumar et al., 2022)

and multiple samples surpassing fluoride, TDS, and hardness limits (BIS, 2023; Jha et al., 2021). These findings point to areas of high contamination and demand focused treatment, public health initiatives, and sustainable groundwater management.

### 2. Scope of the Present Study

The majority of Gujarati groundwater studies are not village-specific, particularly in semi-arid areas like Palanpur Division and Banaskantha, and they frequently use district-wide averages, which obscures local variances that are essential for focused management. Biological indicators like BOD and COD are not fully integrated into previous research, which primarily focuses on physicochemical parameters (BIS, 10500:2023; WHO, 2022). Sources of contamination are often inferred without field-level validation, and comparisons against updated BIS and WHO standards are also scarce. This study fills these gaps by conducting a multi-parameter, village-level assessment at nine different locations. To identify hotspots and guide intervention strategies, triplicate borehole samples for pH, TDS, fluoride, total hardness, major ions, BOD, COD, and DO are analyzed and benchmarked against BIS (10500:2023) and WHO (2022) standards.



### 3. Importance of the Study

In the semi-arid Palanpur Division, where communities rely significantly on groundwater for everyday needs, this study offers an essential, localized evaluation of groundwater quality. It closes a significant data gap that prevents focused intervention by examining village-level differences in physicochemical and biological parameters, such as BOD and COD. The results set a scientific foundation for sustainable groundwater monitoring and management while supporting rural water safety planning, policy creation, and the implementation of localized treatment solutions like defluoridation or RO systems.

### 4. Review of Literature

In India, groundwater quality is a major environmental and public health issue, especially in semi-arid areas like Banaskantha where contamination is influenced by both natural geogenic processes and human activities like waste discharge and agricultural runoff (Adimalla & Venkatayogi, 2023). Gujarat has a serious fluoride pollution problem, with levels above WHO guidelines associated with skeletal and dental fluorosis (Gupta et al., 2017; Saxena & Ahmed, 2003). In Gujarat borewells, high TDS is frequently reported to have an adverse effect on health and palatability (Jha et al., 2021; Shukla et al., 2020). BOD and COD, two indicators of organic pollution, are rising and are frequently associated with inadequate sanitation in the vicinity of agricultural areas (CPCB, 2021; Giri & Singh, 2014). Multivariate statistical tools aid in mapping hotspots and identifying the sources of contaminants (Kumar et al., 2020). Integrating local data into governance is emphasized by sustainable management frameworks (UN-Water, 2023; Singh et al., 2022). In semi-arid belts, seasonal fluctuations and overextraction deteriorate water quality (Rao et al., 2015; Rajmohan & Elango, 2005). While seasonal studies indicate post-monsoon dilution improves quality, geogenic elements contribute to hardness that affects infrastructure and health (Pathak & Vyas, 2019) (Yadav et al., 2021). For focused mitigation, village-level evaluations work better (Sharma et al., 2018). Micro-regional contamination patterns are revealed by sophisticated tools such as GIS (Singh & Bhatnagar, 2020). BIS (2023) and WHO (2022) offer crucial compliance benchmarks. According to Misra et al. (2022), socioeconomic factors also affect

contamination, emphasizing the necessity of interdisciplinary approaches. Incorporating physicochemical, microbial, and heavy metal data into comprehensive water quality indices is also advised by reviews (Ghosh & Bhattacharya, 2021; Pandey & Singh, 2019).

### 5. Objectives of the Study

The primary aim of this study is to assess the groundwater quality in selected villages of the Palanpur Division, Banaskantha district, Gujarat, by analyzing a comprehensive set of physicochemical and biological parameters. The specific objectives are as follows:

1. To evaluate the physicochemical characteristics (such as pH, TDS, electrical conductivity, total hardness, calcium, magnesium, chloride, nitrate, sulfate, and alkalinity) of groundwater samples collected from nine selected villages.
2. To determine the levels of key biological parameters, including dissolved oxygen (DO), biological oxygen demand (BOD), and chemical oxygen demand (COD), to assess organic pollution and potential anthropogenic impacts on water quality.
3. To identify spatial variations in groundwater quality across different villages and water sources (wells, boreholes, and hand pumps), highlighting areas with critical contamination issues.

### 6. Materials and Methods

#### 6.1 Sampling Sites

The study was carried out in the semi-arid Palanpur Division of Banaskantha, Gujarat, which depends on groundwater for domestic, agricultural, and drinking purposes. Based on factors like population, reliance on groundwater, reported quality problems, and geographic distribution, nine villages were chosen. In order to represent actual exposure risks, sampling points included standposts, hand pumps, boreholes, and wells.

#### 6.2 Sample Collection

In accordance with APHA (2017) and BIS guidelines, a total of 27 samples—three from each village—were taken from hand pumps, wells, or boreholes. Glass-stoppered BOD bottles were used for BOD/COD analyses, while HDPE bottles were used for physicochemical analyses. Samples were transported in



ice-cooled boxes and preserved on-site using solutions such as sodium thiosulfate for chlorine and nitric acid for metals. Temperature, pH, turbidity, electrical conductivity, DO, and other field parameters were measured using calibrated portable meters, and laboratory analyses were finished in a day.

### 6.3 Analytical Methods

Biological and physicochemical parameters were examined in accordance with WHO (2017), BIS 10500:2012, and APHA (2017) guidelines. On-site measurements were made of pH, turbidity, TDS, and conductivity using a conductivity meter. UV-Vis spectrophotometry was used to quantify fluoride, chloride, nitrate, and sulfate. Titration techniques were used to measure alkalinity, calcium, magnesium, and hardness. Winkler's method was used to measure DO, dichromate reflux was used to measure COD, and a 5-day incubation at 20°C was used to measure BOD.

### 6.4 Methodology

Using triplicate samples taken during a single season in accordance with BIS (10500:2023) and APHA (2017) guidelines, the quality of the groundwater was evaluated. Fluoride, TDS, chloride, hardness, calcium, magnesium, alkalinity, nitrate, sulfate, BOD, and COD were all analyzed in the lab. To assess whether the results were suitable for drinking, they were compared to BIS and WHO standards.

### 6.5 Standards

The outcomes were contrasted with WHO (2022) drinking water standards and BIS 10500:2023. WHO offers globally accepted thresholds with updated TDS and fluoride limits (WHO, 2022; Adimalla & Venkatayogi, 2023), while BIS offers desirable and acceptable limits for physicochemical and biological parameters, such as fluoride and heavy metals (BIS, 2023; CGWB, 2022). The Palanpur Division's water management plans were guided by this comparison, which also assisted in identifying hazardous areas (Jha et al., 2021). (UN-Water, 2023).

## 7 Results and Discussion

Groundwater analysis from 27 samples across nine villages in Palanpur Division exhibited significant spatial variability in comparison to BIS (10500:2012) and WHO (2017) standards. The pH (7.0–8.26) and turbidity (0.3–1 NTU) were both fine. Fluoride levels often hit the upper limit of 1 mg/L, especially in Chadotar, Chandisar,

Gadh, Madana, and Malan. This means that there was a risk of fluorosis. The total dissolved solids (TDS) levels were between 180 and 1270 mg/L. Madana, Gadh, and Khumbhasan all had levels higher than the recommended 500 mg/L. The hardness ranged from 100 to 560 mg/L, and the samples from Gadh and Madana were very hard. Nitrate levels were mostly low, but they were dangerously high at Madana (415 mg/L), which could be bad for babies' health. Sulfate levels stayed safe. Chloride (120–655 mg/L) was below the 1000 mg/L limit, but higher levels in Madana suggest salinity problems. DO levels were between 3.8 and 8.6 mg/L, BOD levels were between 1.2 and 7.8 mg/L, and COD levels were between 8 and 22.4 mg/L. This showed that there was organic pollution in certain areas. Madana and Gadh were found to be major contamination hotspots that needed actions like defluoridation, TDS and nitrate removal, and community-level water safety planning.

## 8. Conclusion

Groundwater quality showed marked variability across sites. pH (7.0–8.26) remained within WHO/BIS limits, though Jagana (8.26) indicated slight alkalinity. Fluoride (0.5–1 mg/L) was below the 1.5 mg/L threshold, with lower values in Malan, Sasam, and Takarwada. TDS ranged widely (180–2540 mg/L), with Madana (S18) far exceeding the 500 mg/L limit, while Malan (S20) was lowest. Conductivity followed the same trend, and turbidity (0.3–1 NTU) remained safe. DO varied from 3.8–10.8 mg/L, with low levels in Chadotar and Sasam, while BOD (up to 7.8 mg/L) and COD (up to 22.4 mg/L) peaked in Chandisar and Malan, reflecting organic pollution. Chloride exceeded 250 mg/L in Madana (654.79 mg/L) and Chandisar (364.88 mg/L), suggesting salinity or effluent impact. Hardness was high in Madana (560 mg/L) and Gadh (420 mg/L), linked to carbonate aquifers, while alkalinity (55–415 mg/L) and sulfate (55–81 mg/L) were within limits. Nitrate (20–38 mg/L) remained safe, indicating limited fertilizer intrusion.




Figure 1: Heatmap of Water Quality Parameters




Figure 2: Bar chart showing pH levels




Figure: 3 Bar chart illustrating fluoride concentrations (mg/L)




Figure: 4 Bar chart TDS (mg/L)




Figure: 5 Bar chart Conductivity µS/cm




Figure: 6 Bar chart Turbidity (NTU)




Figure: 7 Bar chart Chloride (mg/L)




Figure: 10 Bar chart Mg<sup>2+</sup> (mg/L)




Figure: 8 Bar chart TH (mg/L)




Figure: 11 Bar chart TA (mg/L)




Figure: 9 Bar chart ca<sup>2+</sup> (mg/L)




Figure: 12 Bar chart Nitrate (mg/L)




Figure: 13 Bar chart Sulphate (mg/L)




Figure: 16 Bar chart COD (mg/L)




Figure: 14 Bar chart DO (mg/L)




Figure: 15 Bar chart BOD (mg/L)

**Key Findings**

1. All of the villages had safe pH levels (7.0–8.26).
2. High fluoride levels (about 1.0 mg/L) in Chadotar, Chandisar, Gadh, Madana, and Malan put people at risk for fluorosis.
3. TDS levels in Madana (1270–2540 mg/L) were too high, which means there were too many minerals.
4. Because there is a lot of Ca and Mg in Gadh and Madana, the water is very hard.
5. The level of nitrate in Madana (415 mg/L) was much higher than what is safe, which could be bad for your health.
6. The high levels of chloride (654.79 mg/L) and alkalinity (415 mg/L) in Madana suggest that saltwater or wastewater is getting in.
7. There was not much turbidity, but the high BOD (7.8 mg/L) and COD (22.4 mg/L) levels show that some places were polluted with organic matter.
8. Madana and Gadh were found to be pollution hotspots that need to be cleaned up.
9. It's very important to keep an eye on things, be aware, and have safety plans for each village.

**References**

1. Adimalla, N., & Venkatayogi, P. (2023). A review on fluoride contamination in groundwater and human health. *Environmental Research*, 221, 114872. <https://doi.org/10.1016/j.envres.2023.114872>



2. Adimalla, N., & Venkatayogi, S. (2023). Groundwater contamination in semi-arid regions: A comprehensive review of sources, impacts, and mitigation strategies. *Environmental Science and Pollution Research*, 30(15), 43210–43232. <https://doi.org/10.1007/s11356-023-25635-z>
3. Alsaqqar, M., Abdul-Wahhab, H. M., & Al-Daini, N. A. S. (2021). Water treatment indices: Langelier saturation index application. *Desalination and Water Treatment*, 52(37–39), 7294–7301. <https://doi.org/10.1080/19443994.2021.1877359>
4. BIS. (2023). *Indian Standard Drinking Water Specification (IS 10500:2023)*. Bureau of Indian Standards. <https://www.bis.gov.in>
5. Central Pollution Control Board. (2021). *Guidelines for water quality monitoring and assessment*. <https://cpcb.nic.in>
6. CGWB. (2022). Annual Report: Ground Water Year Book 2020–21 – Gujarat Region. Central Ground Water Board.
7. CPCB. (2021). National water quality monitoring: Annual Report 2020–21. Central Pollution Control Board.
8. Das, A., et al. (2021). Isotope-hydrogeochemical sources and controls of fluoride in semi-arid Gujarat groundwater. *Environmental Geochemistry and Health*, 43(6), 2145–2161. <https://doi.org/10.1007/s10653-021-00894-2>
9. Ghosh, R., & Bhattacharya, S. (2021). Integrated approaches to groundwater quality assessment: A review. *Groundwater for Sustainable Development*, 15, 100658. <https://doi.org/10.1016/j.gsd.2021.100658>
10. Giri, S., & Singh, A. K. (2014). Groundwater quality assessment using water quality index and GIS in the urban area of Ranchi, Jharkhand. *Environmental Earth Sciences*, 71(3), 1243–1257. <https://doi.org/10.1007/s12665-013-2527-1>
11. Gupta, M., Singh, S., Mehta, N., & Roy, A. (2022). Organic pollutant levels and water quality index in riverine systems, Gujarat. *Environmental Pollution*, 299, 118884. <https://doi.org/10.1016/j.envpol.2022.118884>
12. Gupta, S. K., Deshpande, R. D., Agarwal, M., & Raval, B. R. (2017). Origin and extent of groundwater fluoride contamination: A case study from Gujarat, India. *Journal of Hydrology*, 540, 357–370. <https://doi.org/10.1016/j.jhydrol.2016.06.028>
13. Jha, B., Patel, T., & Singh, R. (2021). Effects of elevated total dissolved solids on crop yield under semi-arid conditions in Gujarat, India. *Agricultural Water Management*, 251, 106831. <https://doi.org/10.1016/j.agwat.2021.106831>
14. Jha, L. K., & Panday, D. (2018). Groundwater quality deterioration in semi-arid India: strategy and implications. *Environmental Sciences Europe*, 30, 45. <https://doi.org/10.1186/s12302-018-0164-5>
15. Jha, M. K., Shekhar, A., & Jenifer, M. A. (2021). Assessing the suitability of groundwater for drinking and irrigation in semi-arid regions: A review. *Environmental Geochemistry and Health*, 43(3), 1189–1214. <https://doi.org/10.1007/s10653-020-00675-3>
16. Joshi, P., Desai, A., & Mehta, P. (2023). Hydrochemical characterization of Narmada district groundwater, Gujarat. *Environmental Monitoring and Assessment*, 195(4), 511. <https://doi.org/10.1007/s10661-023-11234-2>
17. Kaur, R., & Kaur, J. (2020). Trends in groundwater fluoride distribution: A meta-analysis of India. *Science of the Total Environment*, 712, 135171. <https://doi.org/10.1016/j.scitotenv.2019.135171>
18. Khatri, N., Tyagi, S., Rawtani, D., Tharmavaram, M., & Kamboj, R. D. (2020). Groundwater quality assessment in Satlasana taluka, Gujarat. *Groundwater for Sustainable Development*, 10, 100321. <https://doi.org/10.1016/j.gsd.2019.100321>
19. Kumar, M., Ramanathan, A. L., Tripathi, R., Farswan, S., Kumar, D., & Bhattacharya, P. (2020). A study of trace element contamination using multivariate statistical techniques and health risk assessment in groundwater of Chhaprola Industrial Area. *Chemosphere*, 259, 127458. <https://doi.org/10.1016/j.chemosphere.2020.127458>



20. Kumar, P., Verma, S., & Sharma, A. (2022). Agricultural runoff-induced nitrate contamination in groundwater: evidence from Gujarat, India. *Environmental Monitoring and Assessment*, 194(10), 725. <https://doi.org/10.1007/s10661-022-10021-5>
21. Kumar, V., & Meena, R. S. (2021). Groundwater contamination due to industrial effluents: Gujarat industrial belt. *Environmental Pollution*, 268, 115847. <https://doi.org/10.1016/j.envpol.2020.115847>
22. Meenakshi, S., Singh, T., & Gupta, R. (2019). Role of ion exchange in fluoride-rich groundwater genesis. *Journal of Hydrology*, 572, 1032–1042. <https://doi.org/10.1016/j.jhydrol.2019.03.058>
23. Mishra, P., Pandey, A., & Singh, A. (2022). Socio-environmental assessment of groundwater quality in rural India. *Sustainable Cities and Society*, 76, 103462. <https://doi.org/10.1016/j.scs.2021.103462>
24. Mohamed, N. A., Wachemo, A. C., Karuppanan, S., & Duraisamy, K. (2022). Spatio-temporal variabilities in groundwater hydrochemistry in Ethiopia. *Urban Climate*, 46, 101338. <https://doi.org/10.1016/j.uclim.2022.101338>
25. Pandey, M., & Singh, R. K. (2019). Comprehensive water quality assessment using water quality index. *Groundwater for Sustainable Development*, 9, 100238. <https://doi.org/10.1016/j.gsd.2019.100238>
26. Pandey, V., Chotaliya, B., Bist, N., Yadav, K., & Sircar, A. (2023). Hydrogeochemistry of geothermal groundwater in Gujarat. *Energy Geoscience*, 4, 59–73. <https://doi.org/10.1016/j.engeos.2022.08.001>
27. Patel, S., Mehta, H., & Shah, L. (2023). Groundwater quality and water quality index (WQI) assessment in Khambhat, Gujarat. *Groundwater for Sustainable Development*, 20, 100421. <https://doi.org/10.1016/j.gsd.2023.100421>
28. Pathak, R. K., & Vyas, A. (2019). Hydrochemical characterization and evaluation of groundwater quality. *Arabian Journal of Geosciences*, 12(19), 589. <https://doi.org/10.1007/s12517-019-4710-y>
29. Pradhan, R. M., & Biswal, T. K. (2018). Fluoride in groundwater: a case study in Precambrian terranes of Ambaji region, North Gujarat, India. *Proceedings of IAHS*, 379, 351–356. <https://doi.org/10.5194/piahs-379-351-2018>
30. Rajmohan, N., & Elango, L. (2005). Nutrient concentrations in groundwater of a shallow weathered rock aquifer in southern India. *Environmental Geology*, 47(6), 846–855. <https://doi.org/10.1007/s00254-004-1207-5>
31. Rao, G. S., Rao, D. V., & Reddy, B. M. (2015). Groundwater quality assessment in parts of Andhra Pradesh. *Environmental Earth Sciences*, 73(7), 3805–3818. <https://doi.org/10.1007/s12665-014-3644-x>
32. Rawat, K. S., Singh, S. K., & Gautam, S. K. (2018). Irrigation suitability of groundwater in peninsular India: a case study. *Applied Water Science*, 8, 1–24. <https://doi.org/10.1007/s13201-018-0866-8>
33. Sahin Kiy, M., & Arslan, H. (2021). Groundwater quality evaluation for irrigation in Turkey. *Irrigation and Drainage*, 70, 871–886. <https://doi.org/10.1002/ird.2593>
34. Saxena, V. K., & Ahmed, S. (2003). Inferring the chemical parameters for the dissolution of fluoride in groundwater. *Environmental Geology*, 43(6), 731–736. <https://doi.org/10.1007/s00254-002-0694-1>
35. Shah, M. K., & Patel, J. (2021). Assessment of hardness and salinity in urban groundwater: case of Ahmedabad. *Journal of Water Science*, 35(2), 150–161. <https://doi.org/10.2166/ws.2021.058>
36. Sharma, D., Khillare, P. S., & Balachandran, S. (2018). Water quality assessment of groundwater using WQI and GIS techniques in Indian villages. *Environmental Monitoring and Assessment*, 190(11), 1–12. <https://doi.org/10.1007/s10661-018-6949-9>
37. Shirke, K. D., Kadam, A., & Pawar, N. J. (2021). Health risk assessment and prevalence of fluoride in groundwater around geological diversity of Ambadongar, South Gujarat, India. *Human and Ecological Risk Assessment*, 27(6), 1523–1542. <https://doi.org/10.1080/10807039.2020.1858270>



38. Shukla, S., Sharma, M., & Maheshwari, S. (2020). Seasonal variation in groundwater quality and its impact on public health. *Journal of Water and Health*, 18(5), 699–712. <https://doi.org/10.2166/wh.2020.036>
39. Singh, A., Sharma, P., & Ghosh, P. (2022). Monitoring spatio-temporal fluoride variability in northern India groundwater. *Environmental Monitoring and Assessment*, 194(5), 326. <https://doi.org/10.1007/s10661-022-10021-5>
40. Singh, G., & Bhatnagar, A. (2020). Groundwater quality mapping using geospatial and statistical techniques in western India. *HydroResearch*, 3, 99–111. <https://doi.org/10.1016/j.hydres.2020.07.002>
41. Singh, H., Yadav, A. K., & Verma, N. (2022). Community-driven groundwater monitoring for rural water security. *Water Policy*, 24(1), 140–158. <https://doi.org/10.2166/wp.2022.079>
42. Singh, P. K., & Tiwari, A. K. (2016). Hydrogeochemical characterization and groundwater quality assessment. *Environmental Earth Sciences*, 75(8), 1–19. <https://doi.org/10.1007/s12665-016-5433-5>
43. Singh, S. K., & Tiwari, A. (2017). Contamination mapping using WQI in central India. *Groundwater for Sustainable Development*, 4, 20–30. <https://doi.org/10.1016/j.gsd.2016.12.001>
44. UN-Water. (2023). *Global water quality monitoring framework: Guidelines for sustainable water management*. <https://www.unwater.org>
45. UN-Water. (2023). *Integrated water quality assessment report 2022*. UN-Water.
46. Vyas, S. J., Tank, S. P., Bhatt, R. P., & Jani, M. M. (2019). Physico-chemical assessment of Sonrakh River, Junagadh City, Gujarat. *International Journal of Scientific Research in Biological Sciences*, 5(6), 89–95. <https://doi.org/10.26438/ijrbs/v5i6.8995>
47. WHO. (2022). *Guidelines for Drinking-water Quality* (5th ed.). World Health Organization. <https://www.who.int/publications/i/item/9789241549950>
48. Yadav, S. K., Choudhary, K., & Meena, M. (2021). Seasonal variation of groundwater quality in Rajasthan. *Environmental Monitoring and Assessment*, 193(6), 1–15. <https://doi.org/10.1007/s10661-021-09020-2>
49. Zope, P. E., Eldho, T. I., & Jothiprakash, V. (2020). Groundwater pollution vulnerability modeling. *Journal of Hydrology*, 584, 124713. <https://doi.org/10.1016/j.jhydrol.2020.124713>
50. Zuin, S., & Ortolani, T. (2019). Groundwater governance challenges in South Asia. *Water Policy*, 21(2), 364–382. <https://doi.org/10.2166/wp.2019.005>