

# The Role of Pharmacists in Combating Antimicrobial Resistance in Saudi Arabia

**Mashael Duhim Alanazi<sup>1</sup>, Maram Suwaylih Assaj Alanazi<sup>1</sup>, Mona Daham Lawwah Alenezi<sup>1</sup>, Laila Sumayhan Ayidh Alanazi<sup>1</sup>, Nawta Sayil Alanzi<sup>1</sup>, Naifah Gasham Khalif Alenezi<sup>1</sup>, Mona Audah alenazi<sup>1</sup>**

*1 Pharmacy Technician, Prince Abdulaziz Bin Musaed Hospital, The Northern Border Health Cluster, Arar, Saudi Arabia*

## **Abstract:**

Pharmacists play a critical role in mitigating antimicrobial resistance (AMR) in Saudi Arabia by promoting responsible medication use and educating patients about the risks associated with antibiotic misuse. As healthcare professionals with extensive knowledge of pharmacology, pharmacists are uniquely positioned to counsel patients on the appropriate use of antibiotics, ensuring that prescriptions are adhered to, and that unnecessary usage is minimized. They actively engage in community outreach programs, disseminating information about AMR and its implications for public health. Through these efforts, pharmacists foster a greater understanding among consumers about the importance of completing prescribed antibiotic courses and the dangers of self-medication or sharing medications. In addition to patient education, pharmacists in Saudi Arabia collaborate with healthcare teams to monitor antimicrobial prescribing patterns and enhance stewardship programs within hospitals and clinics. By conducting medication reviews, they identify potential instances of antibiotic overuse and suggest alternatives, thereby contributing to ongoing efforts to optimize treatment regimens. Their involvement in research and policy-making initiatives also helps to inform national strategies aimed at combatting AMR. Overall, pharmacists are essential in shaping a culture of responsible antibiotic use, ultimately leading to improved health outcomes and a reduction in the prevalence of resistant infections in the region.

**Keywords:** Pharmacists, antimicrobial resistance, Saudi Arabia, patient education, medication stewardship, antibiotic misuse, public health, healthcare collaboration, community outreach, treatment optimization.

## **Introduction:**

Antimicrobial resistance (AMR) has emerged as one of the most significant challenges facing global health systems today. It threatens the success of modern medicine, undermines advances in surgical procedures, and complicates the treatment of infectious diseases. The rise of AMR is attributed to various factors, including inappropriate use of antibiotics, over-prescription, and a lack of patient education regarding medication adherence. Global health organizations — including the World Health Organization (WHO) — have issued alarming warnings about the imminent threat posed by AMR, predicting that without concerted efforts, AMR could cause up to 10 million deaths annually by 2050. In Saudi Arabia, the issue of AMR is particularly pressing, given the country's rapidly growing population, healthcare system transformation, and increasing prevalence of antibiotic-resistant infections [1].

The role of pharmacists in addressing this critical public health crisis cannot be overstated. As the most accessible healthcare professionals, pharmacists are uniquely positioned to influence antibiotic use and promote rational prescribing practices. In Saudi Arabia, where antibiotic misuse

is pervasive, pharmacists can serve as frontline defenders against AMR. Their extensive knowledge of pharmacotherapy and patient interaction equips them to educate prescribers and patients alike, thereby enhancing adherence to appropriate antimicrobial stewardship practices. This multifaceted role encompasses several dimensions, including patient education, medication management, surveillance of antibiotic use, and collaboration with other healthcare professionals to develop and implement policies aimed at combating AMR [2].

In recent years, various stakeholders in the healthcare system, including the Saudi Ministry of Health, have recognized the integral role pharmacists must play in combatting AMR. Initiatives aimed at enhancing the training and responsibilities of pharmacists include expanding their scope of practice in antimicrobial stewardship, offering specialized education on infectious diseases, and promoting interdisciplinary collaboration among healthcare providers. Given the diverse and culturally rich medical landscape of Saudi Arabia, which incorporates traditional practices along with modern medicine, it is essential to integrate pharmacists into the broader framework of healthcare delivery, policy formulation, and public health campaigns addressing AMR [3].

Pharmacists also play a crucial role in promoting appropriate antibiotic use through patient and community education. Many patients are often unaware of the potential harms associated with self-medication or the importance of completing prescribed antibiotic courses. By providing guidance on when antibiotics are necessary, when they are not, and how to properly use them, pharmacists can help curb the misuse of these critical medications. Their expertise allows them to assist patients in navigating complex medication regimens, monitor for potential interactions and side effects, and encourage adherence to prescribed therapies, which are all vital in managing AMR [4].

Efforts to curb AMR in Saudi Arabia also hinge on effective surveillance and data collection, areas where pharmacists can make significant contributions. By engaging in mechanisms that track antibiotic prescribing patterns and resistance trends, pharmacists can provide valuable insights that support evidence-based policymaking. This surveillance can inform local guidelines and contribute to national AMR strategies, ensuring that interventions are timely and tailored to the unique healthcare challenges faced in the region [5].

Furthermore, the role of pharmacists extends beyond individual patient interactions to systemic advocacy for improved healthcare policies. By leveraging their knowledge and experience, pharmacists can engage in outreach initiatives aimed at hospitals, academic institutions, and regulatory bodies. Their input can help shape policies that promote antimicrobial stewardship programs, establish guidelines for appropriate prescribing practices, and support public health campaigns aimed at raising awareness about AMR [6].

### **The Pharmacist's Clinical Role in Antimicrobial Stewardship:**

Antimicrobial stewardship (AMS) has emerged as a critical component in the management of infectious diseases, primarily due to the alarming rise of antimicrobial resistance (AMR) across the globe. This resistance not only complicates treatment regimens but also leads to higher medical costs, prolonged hospital stays, and increased mortality rates. In this context, pharmacists play an indispensable role in optimizing the use of antimicrobials, ensuring both efficacy and safety [7].

Antimicrobial stewardship refers to a set of coordinated strategies designed to improve the use of antimicrobials to enhance patient health outcomes, reduce resistance, and decrease unnecessary costs. AMS encompasses several elements, including appropriate prescribing, monitoring, and educating healthcare professionals and patients regarding the proper use of antimicrobials. Effective AMS programs can lead to judicious use of antibiotics, mitigate the spread of resistance, and ultimately improve the quality of care provided to patients [8].

## **The Central Role of Pharmacists in Antimicrobial Stewardship**

Pharmacists are uniquely positioned within the healthcare framework to contribute to antimicrobial stewardship for various reasons, including their expertise in medication management, accessibility to healthcare teams and patients, and their extensive training in pharmacology. The Clinical Pharmacist's role can be divided into various categories:

### **1. Medication Review and Optimization:**

Pharmacists conduct thorough medication evaluations, ensuring that prescribed antimicrobials are appropriate for the patient's specific infection, taking into account pathogens, susceptibility profiles, and patient-specific factors such as renal function or allergies. This involves correlating clinical guidelines with local resistance patterns and assessing the need for broad-spectrum versus narrower-spectrum agents. By fostering a tailored therapeutic approach, pharmacists can help mitigate the indiscriminate use of antibiotics, hence playing a crucial role in curbing AMR [9].

### **2. Dosing Recommendations:**

Pharmacists provide invaluable input regarding the correct dosing of antimicrobials, especially in populations that may require special considerations, such as pediatrics, geriatrics, or those with renal impairment. Ensuring the pharmacokinetic and pharmacodynamic properties of antibiotics are matched with individual patient profiles is vital for achieving effective concentrations while minimizing toxicity and resistance development [10].

### **3. Monitoring and Follow-Up:**

Continuous monitoring of patients on antimicrobial therapy is essential for the early detection of potential adverse effects or therapeutic failures. Pharmacists employ their clinical skills to evaluate the progress of treatment, assess laboratory results such as cultures and sensitivities, and adjust antimicrobial therapy accordingly. Such proactive management not only encourages optimal therapeutic outcomes but also emphasizes the need for reassessment of ongoing therapy based on clinical response and microbiological data [11].

### **4. Education and Training:**

Education is a cornerstone of antimicrobial stewardship, and pharmacists are at the forefront of delivering this knowledge. They are responsible for providing both healthcare professionals and patients with comprehensive education regarding the appropriate use of antimicrobials, risks associated with misuse, and infection control measures that can reduce the reliance on antibiotics. Regular training sessions or seminars for healthcare staff can enhance the entire team's adherence to AMS principles [11].

**5. Collaboration within Healthcare Teams:**

Pharmacists act as integral members of interprofessional teams, collaborating with physicians, nurses, and infection control staff to devise and implement effective AMS strategies. By fostering an environment of interdisciplinary communication, pharmacists can facilitate consensus on best practices and clinical guidelines for antibiotic use, aligning treatment protocols with the goals of AMS [12].

**6. Research and Data Analysis:**

As experts in drug-related information, pharmacists are often involved in the collection and analysis of data related to antimicrobial use and resistance patterns. This information is crucial for refining existing AMS programs and informing future policies. By publishing their findings and insights, pharmacists contribute to the growing body of literature that supports evidence-based practices in antimicrobial stewardship [12].

### **Challenges in Implementing Antimicrobial Stewardship**

Despite the pivotal role pharmacists play in AMS, several challenges hinder their effectiveness. Barriers such as time constraints, insufficient staffing, lack of access to patient-specific data, and a fragmented healthcare system can limit the potential of pharmacists to participate fully in stewardship efforts. Moreover, there may be resistance from prescribers when pharmacists attempt to influence prescribing practices. Overcoming these obstacles requires organizational commitment, adequate resources, and the establishment of clear roles within AMS initiatives [13].

### **Patient Education and Counseling Strategies for Antibiotic Use:**

In recent years, the misuse and overuse of antibiotics have emerged as critical public health issues, contributing significantly to the rise of antibiotic resistance. This phenomenon poses a notable threat to effective treatment options for bacterial infections, leading healthcare professionals to prioritize patient education and counseling as essential strategies for optimizing antibiotic use [14].

Antibiotics are powerful medicines used to treat infections caused by bacteria. However, they are ineffective against viral infections, such as the common cold or influenza. Patients often harbor misconceptions about antibiotics, believing they can cure all types of infections. This misunderstanding can lead to inappropriate requests for antibiotics, pressure on healthcare providers to prescribe them, and ultimately, their unnecessary use. Thus, patient education is pivotal for correcting misconceptions and fostering informed decision-making regarding antibiotic therapies [14].

The consequences of inappropriate antibiotic use extend beyond individual health; they impact community health and public resources. Increased resistance diminishes the efficacy of commonly prescribed antibiotics, leading to prolonged illnesses, increased healthcare costs, and the need for more expensive or toxic alternative treatments. Therefore, patient education is not only a matter of individual responsibility but also a public health priority [15].

## Effective Counseling Strategies

To facilitate patient education regarding antibiotics, healthcare providers can employ various counseling strategies that enhance understanding and promote responsible use [15].

- 1. Clear Communication:**

The complexity of medical terminology can alienate patients and hinder comprehension. Providers should utilize layman's terms, avoid jargon, and ensure that explanations are straightforward. For example, instead of stating that a patient has a "bacterial infection," a provider might say, "You have an infection caused by bacteria, and we need to treat it with antibiotics."

- 2. Engagement Through Questions:**

Encouraging patients to voice their concerns and questions enhances their engagement in their care. Providers can use open-ended questions such as, "What brings you in today?" or "How are you feeling about your treatment plan?" This approach allows the clinician to address specific concerns regarding antibiotics while reinforcing the collaborative nature of the healthcare relationship. When patients feel heard, they are more likely to trust the information shared [16].

- 3. Utilize Visual Aids and Brochures:**

Visual aids, brochures, and handouts can be useful tools for reinforcing verbal information. Infographics that illustrate the differences between viral and bacterial infections or the consequences of antibiotic overuse can serve as quick reference materials for patients. Such resources can significantly improve retention and understanding of key messages [17].

- 4. Tailor Message to Individual Needs:**

Individual patient characteristics affect how they perceive and understand health information. Cultural beliefs, age, education level, and previous experiences with antibiotics will shape a patient's attitude toward their use. Providers should assess these factors and adapt their educational strategies accordingly, ensuring that the techniques used resonate with each patient [17].

- 5. Conveying the Importance of Completing Prescribed Courses:**

Educating patients on the significance of taking the full course of antibiotics, even if symptoms subside, is vital to curtailing resistance development. Patients may feel better and assume they no longer need the medication, which can leave residual bacteria that may develop resistance. Emphasizing this point during consultations can improve adherence and outcomes [18].

- 6. Promote Alternative Treatments for Symptomatic Relief:**

In cases where antibiotics are unnecessary, such as viral infections, healthcare providers should discuss alternative symptomatic treatments. Providing options for supportive care and self-management strategies empowers patients and reassures them that their symptoms are being taken seriously [19].

- 7. Incorporating Follow-Up Plans:**

Implementing a follow-up strategy reinforces the importance of monitoring effectiveness and side effects of antibiotic treatments. Scheduled check-ins allow healthcare providers to reassess the treatment plan and address any emerging concerns about antibiotic use.

Moreover, having a defined follow-up communicates that the patient's health is a priority and encourages them to adhere to prescribed treatments [20].

### **The Role of Healthcare Providers**

Healthcare providers play an essential role in antibiotic stewardship. They are not only responsible for diagnosing and prescribing but also for educating patients about antibiotic use. Medical education should incorporate curricula focused on the principles of antimicrobial resistance, evidence-based prescribing practices, and the social responsibilities of healthcare providers [21].

Providers must also remain vigilant in identifying signs of infection and discerning when antibiotic prescriptions are warranted versus when they can potentially cause harm. By establishing themselves as trusted advisors in antibiotic use, healthcare professionals can advocate for responsible prescribing practices and promote patient-centric care that prioritizes educational efforts to combat antibiotic misuse [22].

### **Pharmacists' Contributions to Policy Development and Implementation:**

Antimicrobial resistance (AMR) has emerged as one of the most pressing public health challenges of the 21st century. As pathogens develop resistance to the very medications designed to eradicate them, the implications for global health are profound; they include increased morbidity and mortality, prolonged hospital stays, and heightened healthcare costs. In response to this urgent threat, policies aimed at the prudent use of antimicrobials have become essential [23].

Pharmacists are medication experts who possess extensive training in pharmacology, therapeutics, and the mechanisms of action of drugs, including antibiotics. Their deep understanding of how antibiotics work enables them to make informed decisions regarding their use, promoting both efficacy and safety. In settings such as hospitals and community pharmacies, pharmacists regularly review medication regimens to ensure appropriate antibiotic selection, dosing, and duration of therapy. This attention to detail is critical to minimizing the risks of ineffective treatment and the development of resistance [24].

Moreover, pharmacists routinely engage in medication reconciliation, particularly for patients transitioning between care settings. This process helps to identify and resolve discrepancies that may occur due to the complex nature of antimicrobial therapy. By monitoring patients for adverse drug reactions and interactions, pharmacists can help to safeguard patient safety while also ensuring that antibiotics are used judiciously [25].

Antimicrobial Stewardship Programs (ASPs) are institutional initiatives aimed at optimizing the use of antimicrobials to enhance patient outcomes while minimizing unintended consequences, such as the emergence of resistant strains. Pharmacists are integral to the development and implementation of these programs. Their role typically includes designing protocols for appropriate prescribing, conducting audits of antimicrobial usage, and providing recommendations based on current guideline adherence [25].

In addition, pharmacists often lead educational initiatives targeting both healthcare providers and patients. By initiating discussions about the importance of appropriate antibiotic use and the dangers of overprescribing, pharmacists foster a culture of stewardship within healthcare settings. They also develop educational materials that empower patients to understand their treatments, emphasizing the need to complete prescribed courses and the dangers associated with misuse [26].

The collection, analysis, and interpretation of data regarding antibiotic usage and resistance patterns are crucial to guiding policymakers in their efforts to combat AMR. Pharmacists contribute significantly to this aspect by monitoring drug utilization and resistance trends, thereby providing critical data that informs antimicrobial policy development [27].

By participating in antimicrobial resistance surveillance programs, pharmacists can identify emerging resistance patterns in their local communities, which are essential for adapting guidelines and practices to current realities. Their input helps to shape institutional and regional policies, ensuring that local data drives the development of targeted antimicrobial strategies. In this way, pharmacists play a vital role in customizing stewardship efforts to meet the unique needs of their populations [28].

Effective antimicrobial policy development relies heavily on interdisciplinary collaboration. Pharmacists serve as essential partners in multidisciplinary teams comprised of physicians, nurses, infection control specialists, and public health professionals. Their involvement facilitates a more comprehensive understanding of the complexities of antimicrobial use and stewardship [29].

For instance, during rounds or clinical decision-making meetings, pharmacists can offer real-time consultation based on their expertise in pharmacotherapy. This collaborative interaction enhances the quality of care and can lead to more informed prescribing practices. Additionally, pharmacists bring a unique perspective to interdisciplinary discussions, often advocating for evidence-based approaches that balance the benefits of antimicrobial therapy against the risks of resistance [30].

Furthermore, pharmacists' familiarity with health informatics tools allows them to engage in data-sharing initiatives with other healthcare providers, streamlining communication and enhancing overall team effectiveness. This collaborative approach not only improves patient care but also supports the successful implementation of antimicrobial policies [31].

Beyond clinical settings, pharmacists advocate for policies at local, national, and international levels aimed at addressing AMR more broadly. Engaging in outreach campaigns that inform the public about the importance of responsible antibiotic use is a key aspect of this advocacy. Through community engagement efforts, pharmacists can disseminate information about AMR, dispel myths surrounding antibiotic therapy, and underscore the importance of vaccination and infection prevention strategies [31].

Pharmacists also often collaborate with public health organizations to devise educational materials and community programs focused on antimicrobial stewardship. Their role in these initiatives is critical in shaping public perception and behavior regarding antibiotic use, fostering a culture of responsibility that transcends healthcare settings [32].

Pharmacists are pivotal in educating both healthcare professionals and the public about the importance of antimicrobial stewardship. As educators, they are often involved in training programs aimed at enhancing the competencies of their colleagues regarding appropriate antimicrobial prescribing practices. By offering continuing education programs and workshops, pharmacists ensure that healthcare providers remain current with the latest research and guidelines [33].

Moreover, pharmacists educate patients about their medications, including correct usage, potential side effects, and the dangers associated with improper use. This direct interaction is fundamental in promoting adherence to antimicrobial therapy, helping to prevent treatment failures and reduce the risk of resistance development [34].

### **Interprofessional Collaboration in the Fight Against Antimicrobial Resistance:**

Antimicrobial resistance (AMR) has emerged as one of the most pressing global health challenges of the 21st century. The rise of resistant strains of bacteria, viruses, fungi, and parasites has led to increased morbidity, mortality, and healthcare costs, challenging the efficacy of antimicrobial therapies that have been the cornerstone of modern medicine. Addressing this complex issue requires a coordinated approach involving multiple disciplines across the healthcare continuum, emphasizing the necessity of interprofessional collaboration [35].

Before delving into the mechanisms of interprofessional collaboration, it is imperative to understand the landscape of AMR. Antimicrobial resistance occurs when microorganisms evolve and develop the ability to resist the effects of drugs, rendering standard treatments ineffective. This phenomenon can result from various factors, including misuse and overuse of antimicrobial agents, inadequate sanitary conditions, and lack of patient education. The World Health Organization (WHO) has identified AMR as a critical public health threat, contributing to an estimated 1.27 million deaths in 2019 alone [36].

The insidious nature of AMR underscores its multifactorial origins, which necessitate a holistic approach to management. An effective response requires not only medical expertise but also insights from microbiology, pharmacy, nursing, public health, and social sciences. By fostering interprofessional collaboration, healthcare teams can leverage their diverse skills and perspectives to develop comprehensive strategies to combat AMR [37].

### **The Importance of Interprofessional Collaboration**

Interprofessional collaboration (IPC) is defined as the process of individuals from different professional backgrounds working together towards a common goal. In the context of AMR, IPC is vital for several reasons:

1. **Comprehensive Patient Care:** Effective management of infections requires input from various specialists. For example, infectious disease specialists can provide insight into the most effective antibiotic therapy, while pharmacists can offer guidance on appropriate dosing and identify potential drug interactions. Nurses play a critical role in monitoring

patient responses and educating them about completing the prescribed treatment regimen [38].

2. **Improved Stewardship Programs:** Antimicrobial stewardship programs aim to optimize the treatment of infections while minimizing the adverse effects associated with inappropriate use of antibiotics. Interprofessional teams can design and implement these programs, integrating strategies from different professions to enhance adherence to guidelines, monitor outcomes, and adjust protocols as necessary [38].
3. **Education and Awareness:** Public awareness campaigns about AMR can directly benefit from interprofessional efforts. Health educators, pharmacists, and physicians can collaborate to create educational materials that inform patients about the importance of adhering to prescribed treatment regimens, the dangers of self-medication, and the role of vaccinations in preventing infections [39].
4. **Research and Development:** Addressing AMR requires ongoing research and innovation. Interprofessional teams are better equipped to conduct comprehensive studies that consider the biological, clinical, and sociological aspects of AMR, leading to more robust and effective solutions [40].

### Challenges to Interprofessional Collaboration

Despite the clear benefits of IPC, several challenges can hinder effective collaboration among healthcare professionals:

1. **Siloed Practices:** A longstanding tradition of isolated practice within healthcare professions often results in communication barriers and a lack of understanding of each other's expertise. These silos can exacerbate issues related to AMR management and ultimately compromise patient care [41].
2. **Differing Professional Cultures:** Each healthcare profession has its own culture, language, and practices. Differences in perspectives and priorities can lead to misunderstandings or conflict within interprofessional teams. Developing mutual respect and understanding among team members is essential for fostering collaboration [41].
3. **Lack of Training:** Many healthcare professionals receive insufficient training in IPC during their education. This lack of exposure can result in apprehension or reluctance to engage in collaboration, limiting the potential for comprehensive care in combating AMR [41].
4. **Resource Constraints:** Effective interprofessional collaboration may require additional staffing, time, and financial resources. In settings with limited resources, such as underfunded healthcare systems, prioritizing IPC can be difficult [41].

### Community Outreach Initiatives to Raise Awareness of AMR:

Antimicrobial Resistance (AMR) represents one of the most pressing public health challenges of the 21st century. Characterized by the ability of microbes to resist the effects of medications—particularly antibiotics—AMR complicates the treatment of infectious diseases, leading to prolonged illnesses, increased healthcare costs, and a heightened risk of death. According to the World Health Organization (WHO), AMR is responsible for an estimated 700,000 deaths annually worldwide, a figure that could escalate to 10 million by 2050 if no effective interventions are

implemented. Given the profound implications of AMR, community outreach initiatives aimed at raising awareness are critical to mobilizing public action and encouraging responsible behaviors towards antimicrobial usage [42].

Before delving into outreach initiatives, it is vital to comprehend the mechanics of AMR. The origin of AMR can primarily be attributed to the misuse and overuse of antimicrobial agents across various settings, including healthcare, agriculture, and veterinary practices. When antibiotics are excessively prescribed, taken incorrectly, or used to promote growth in livestock, they can foster the emergence of resistant strains of bacteria [42].

Moreover, AMR is exacerbated by global factors, such as increased travel, trade, and urbanization, which enable the rapid spread of resistant microbes. Inadequate sanitation, insufficient infection prevention measures, and lack of access to quality healthcare are additional contributors to the escalation of AMR. As communities become more aware of these underlying factors, outreach initiatives can play a pivotal role in altering behavior at both individual and collective levels [43].

### **The Role and Importance of Community Outreach Initiatives**

Community outreach initiatives serve as essential platforms for disseminating knowledge about AMR and mobilizing local populations toward collective action. Their effectiveness is rooted in several key components:

1. **Education and Awareness:** Outreach efforts focus on educating communities about the causes and consequences of AMR. This education is critical in debunking common myths about antimicrobials, such as the belief that antibiotics can treat viral infections like the common cold or flu. Accurate information empowers individuals to make informed choices about their health and that of their families [44].
2. **Behavioral Change:** By increasing awareness, outreach initiatives can foster behavioral change. For instance, educating patients about the importance of completing prescribed courses of antibiotics can significantly diminish the chances of resistance development. Additionally, campaigns that emphasize preventive measures, such as vaccination and proper hygiene practices, can reduce the need for antimicrobials in the first place.
3. **Community Engagement:** Successful outreach initiatives often involve local stakeholders, including healthcare providers, schools, religious organizations, and community leaders. By leveraging existing community structures, initiatives can effectively communicate the importance of AMR awareness and encourage participation in prevention strategies [44].
4. **Policy Advocacy:** Grassroots movements can drive policy changes at local, regional, and national levels. By mobilizing communities to express their concerns about AMR, outreach initiatives can influence decision-makers to prioritize AMR in public health agendas, allocate funding for research, and promote responsible antibiotic use [45].

### **Examples of Community Outreach Initiatives**

Numerous innovative outreach initiatives have successfully raised awareness of AMR across various communities globally. Here are some notable examples:

1. **The Antibiotic Guardian Campaign:** Launched by Public Health England, this campaign calls upon individuals to take pledges on how they can reduce antibiotic resistance. Participants commit to actions such as not demanding antibiotics for viral infections and completing their prescribed medication. This national initiative effectively engages communities and generates substantial public commitment towards responsible antibiotic use [46].
2. **School-based Programs:** Many outreach initiatives target school-age children to instill awareness about AMR early on. Programs that combine classroom education with interactive activities—such as games, discussions, and science fairs—teach children about the dangers of antibiotic misuse. Through these programs, children learn to advocate for proper hygiene practices, which help them combat infections without unnecessary reliance on antimicrobials [47].
3. **Community Health Workshops:** Local health departments and non-governmental organizations often host workshops aimed at educating community members about AMR. These workshops typically feature expert speakers, informational brochures, and Q&A sessions to clarify misinformation about antibiotics. By creating a safe and open environment for discussion, these initiatives foster trust and enhance community engagement [47].
4. **Social Media Campaigns:** In the digital age, social media has emerged as a powerful tool for raising awareness. Campaigns designed for platforms such as Facebook, Instagram, and Twitter can reach diverse age groups and demographics. Using eye-catching infographics, educational videos, and testimonials from healthcare professionals, social media campaigns can rapidly disseminate information and mobilize communities towards action [48].
5. **Collaborative Initiatives:** Another effective model for community outreach involves partnerships among various stakeholders, including public health agencies, educational institutions, and local businesses. For instance, local pharmacies might collaborate with health departments to offer free consultations regarding antimicrobial use, while local schools might provide materials for distribution during health fairs [48].

## Challenges Ahead

While community outreach initiatives are vital in raising awareness of AMR, they face several challenges. One significant hurdle is the pervasive misinformation regarding antibiotics and their use. Mistrust in public health messages can lead individuals to rely on anecdotal evidence rather than scientific recommendations. Moreover, limited access to healthcare and education in some communities can inhibit the effectiveness of outreach efforts [49].

Furthermore, sustaining momentum for AMR initiatives can be challenging, as community interest may wane over time. Therefore, it is critical for outreach initiatives to create a framework that ensures ongoing engagement and education, adapting to changing community dynamics and emerging research on AMR [50].

## **Challenges Faced by Pharmacists in Addressing Antimicrobial Resistance:**

Antimicrobial resistance (AMR) has emerged as a formidable public health challenge globally, threatening the effectiveness of antimicrobial agents such as antibiotics, antifungals, and antivirals. According to the World Health Organization (WHO), AMR leads to 700,000 deaths annually, and without significant action, this number is projected to rise sharply within the coming years. As healthcare professionals at the forefront of patient care and medication management, pharmacists play a pivotal role in combating AMR. However, they face numerous challenges in addressing this pressing issue, ranging from misinformation about antimicrobial usage to systemic healthcare barriers [51].

One of the most significant challenges pharmacists face in addressing AMR is the widespread misinformation surrounding antimicrobial use. Many patients harbor misconceptions about antibiotics, believing they are effective for viral infections such as colds and flu or that they can demand antibiotics without a proper diagnosis. Such misunderstandings often lead to misuse, which is a primary contributor to AMR [52].

Pharmacists are uniquely positioned to bridge this knowledge gap. However, their ability to educate patients is often hindered by time constraints in practice settings. Busy pharmacy environments may limit the interaction time pharmacists have with patients, constraining their capacity to provide detailed educational counseling. Furthermore, pharmacists are sometimes perceived as “just dispensers” of medication rather than healthcare providers, which can diminish their authority in advocating for appropriate antimicrobial stewardship [53].

Another significant barrier pharmacists encounter is the lack of voice and visibility within the broader healthcare system regarding AMR initiatives. While many healthcare settings recognize the importance of antimicrobial stewardship programs (ASPs), pharmacists may not always be included in decision-making processes or program planning. This oversight can stifle the potential impact pharmacists could have in formulating effective strategies for decreasing AMR [54].

Additionally, limitations in the scope of practice can impede pharmacists' ability to implement antimicrobial stewardship effectively. In some regions, pharmacists lack the authority to initiate or modify antimicrobial therapy based on clinical guidelines, which curtails their ability to respond rapidly to emerging resistance patterns. A re-evaluation of pharmacy practice legislation and a move toward more collaborative healthcare models that leverage the expertise of pharmacists could enhance their role in tackling AMR [55].

Access to current and comprehensive resistance data poses another challenge for pharmacists. Information regarding local antimicrobial resistance patterns is crucial for evidence-based clinical decision-making. However, inadequate surveillance systems can hinder pharmacists' ability to choose the most effective antimicrobials or educate patients and healthcare colleagues on updated resistance trends [56].

Moreover, pharmacists often lack access to real-time data concerning patient-specific characteristics that could inform therapeutic decisions, such as microbiological culture results.

This gap is especially notable in community pharmacy settings, where pharmacists may not have the resources to consult laboratory data as readily as those in hospital environments [57].

Effective collaboration among healthcare professionals is crucial in addressing AMR, but interprofessional barriers often stand in the way. Communication gaps between pharmacists and other members of the healthcare team can lead to fragmentation of care, where pharmacists may be unaware of a patient's full drug regimen, underlying conditions, or clinical status. This lack of synergy can result in inappropriate antimicrobial prescriptions, ultimately contributing to increased levels of resistance [58].

Pharmacists also face numerous regulatory challenges in addressing AMR within their practice. Policies governing antimicrobial prescribing are often not well articulated, making it difficult for pharmacists to navigate their roles in stewardship. In some jurisdictions, restrictions surrounding the dispensing of specific antimicrobials may conflict with the pharmacist's best judgment, limiting their involvement in optimizing therapy [59].

Effective advocacy for sound antimicrobial policies is essential. Pharmacists must engage with policymakers to promote legislation that empowers pharmacists in their stewardship roles, improves access to antimicrobial stewardship programs, and enhances funding for AMR initiatives [59].

Despite these challenges, numerous strategies can be implemented to bolster the role of pharmacists in combating AMR. Enhancing educational programs that focus on antimicrobial stewardship during pharmacy education could cultivate a generation of pharmacists adept at addressing AMR issues. Ongoing professional training and development also play a critical role in ensuring pharmacists remain informed about emerging resistance trends and appropriate prescribing practices [60].

Furthermore, pharmacy practice models should evolve to allow greater pharmacist involvement in patient care, such as expanding the scope of practice to permit pharmacists to initiate and modify therapies based on resistance patterns. Enhanced collaboration with other healthcare professionals through structured interprofessional training can also help improve communication and teamwork [61].

Developing robust antimicrobial surveillance systems that track resistance patterns can empower pharmacists with valuable data, enabling more informed clinical decision-making. That, combined with a strong advocacy effort aimed at shaping policy and expanding pharmacists' roles in healthcare teams, can facilitate significant progress toward addressing AMR [61].

### **Future Perspectives and Recommendations for Pharmacists in Saudi Arabia:**

Antimicrobial resistance (AMR) has emerged as a pressing public health concern globally, significantly impacting the effectiveness of medical treatments and posing challenges to healthcare systems. Recent reports from the World Health Organization (WHO) estimate that AMR could lead to approximately 10 million deaths annually by 2050, surpassing deaths from cancer and diabetes combined. In the Kingdom of Saudi Arabia (KSA), like many other countries, the rise of

AMR necessitates urgent action. Pharmacists, being accessible healthcare professionals at the frontline of medication management, possess a unique position to contribute significantly to the fight against AMR [62].

Saudi Arabia has seen a gradual increase in AMR, influenced by factors such as over-prescription of antibiotics, inadequate patient education, and a rise in self-medication practices. Studies indicate that significant percentages of pathogens in the country display resistance to first-line antibiotics, complicating the treatment landscape for various infections. The government has recognized this challenge, and national strategies such as the Saudi National Action Plan to Combat Antimicrobial Resistance (SNAP-AMR) have been established. However, increased awareness among healthcare professionals, particularly pharmacists, is pivotal in conjunction with these governmental initiatives [62].

### **Vision for Pharmacists' Role in Combating AMR**

Pharmacists in KSA can serve as a crucial link between patients and the healthcare system to combat AMR. In the future, pharmacists should envision themselves as:

1. **Educators:** Pharmacists must take an active role in educating patients about the responsible use of antibiotics, emphasizing the dangers of misuse and overuse. This includes informing patients about the importance of adherence to prescribed treatments and the risks associated with self-medication [63].
2. **Consultants:** They should act as essential members of healthcare teams, collaborating with physicians to promote appropriate antimicrobial prescribing practices. By assessing and reviewing prescriptions, pharmacists can provide recommendations to mitigate unnecessary antibiotic usage [63].
3. **Advocates:** Pharmacists should champion AMR awareness campaigns within their communities, utilizing social media, workshops, and other informal forums to disseminate information effectively.
4. **Researchers:** Engaging in research related to AMR will help pharmacists identify trends and contribute valuable data that can inform local health policies and strategies [64].

### **Recommendations for Raising Awareness of AMR**

To empower pharmacists in their roles and enhance their contributions toward raising AMR awareness, several recommendations can be proposed:

1. **Continuing Education and Training:** Mandatory continuing education programs on AMR should be instituted for pharmacists. These programs should focus on the mechanisms of resistance, local resistance patterns, and effective communication strategies to counsel patients regarding antibiotic use [65].
2. **Collaboration with Health Authorities:** Pharmacists should work closely with the Saudi Ministry of Health and local health authorities to develop standardized guidelines that outline best practices for antibiotic dispensing. This additional framework will help pharmacists provide consistent and accurate information to patients [66].

3. **Public Awareness Campaigns:** Pharmacists can collaborate with community organizations to launch widespread campaigns that educate the public about AMR. Utilizing various media channels—such as television, social media, and workshops—can significantly enhance outreach. Tailoring messages to align with culturally relevant practices will ensure higher acceptance and adherence [67].
4. **Integration of Technology:** The use of digital platforms to provide educational resources about AMR could be instrumental. Pharmacists can develop mobile applications or websites that offer information on AMR, responsible antibiotic use, and promote easy access to professional advice [68].
5. **Pharmacy Practice Models:** Implementing advanced practice roles for pharmacists, such as antimicrobial stewardship programs within hospitals and community settings, can create more structured approaches to monitoring and advising on antibiotic use.
6. **Reporting Systems for Resistance Patterns:** Establishing and maintaining a database to track and report antimicrobial resistance patterns in Saudi Arabia can allow pharmacists to provide data-driven recommendations. This effort would involve collaboration with hospitals, clinics, and research institutions [69].
7. **Legislative Support:** Advocacy for legislative changes that restrict over-the-counter sales of antibiotics and enhance prescription drug monitoring is critical. Pharmacists can play a vital role in lobbying for policies that encourage responsible use and dispensing practices [69].

### **Conclusion:**

In conclusion, pharmacists are vital allies in the battle against antimicrobial resistance (AMR) in Saudi Arabia. Their extensive training in pharmacotherapy equips them to provide essential services, including patient education, medication management, and active participation in antimicrobial stewardship programs. By raising awareness about the dangers of antibiotic misuse and promoting adherence to prescribed therapies, pharmacists can significantly contribute to reducing the incidence of resistant infections. Furthermore, their involvement in policy-making and interprofessional collaboration strengthens healthcare strategies aimed at combating AMR at both community and national levels.

As the landscape of healthcare continues to evolve, it is crucial for pharmacists to embrace their role as leaders and educators in the fight against AMR. This necessitates ongoing training, increased access to resources, and fostering collaboration with other healthcare professionals. By enhancing their engagement and visibility within the healthcare system, pharmacists can help ensure a more effective response to AMR, ultimately leading to improved patient outcomes and the preservation of effective antibiotic therapies for future generations.

### **References:**

1. Alomi, Y. A. J. O. A. J. S., 2017. National antimicrobial stewardship program in Saudi Arabia; initiative and the future. 4, 1-7.
2. Hadi M.A., Karami N.A., Al-Muwalid A.S., et al. Community pharmacists' knowledge, attitude, and practices towards dispensing antibiotics without prescription (DAWP): a cross-sectional survey

- in Makkah Province, Saudi Arabia. *Int. J. Infect. Dis.* 2016;47:95–100. doi: 10.1016/j.ijid.2016.06.003.
3. Alrasheedy A.A., Alsalloum M.A., Almuqbil F.A., et al. The impact of law enforcement on dispensing antibiotics without prescription: a multi-methods study from Saudi Arabia. *Expert Rev. Anti-infective Ther.* 2020;18:87–97. doi: 10.1080/14787210.2020.1705156.
  4. AlRukban M., AlRuthia Y., Almasaoud M., et al. Community pharmacists' views of the enforced antibiotics dispensing law and its impact on oral antibiotics sales in Saudi Arabia. *Risk Manag. Healthc. Policy.* 2020;13:2899. doi: 10.2147/RMHP.S278404.
  5. Dadgostar P.J.I., Resistance D. Antimicrobial resistance: implications and costs. *Infect. Drug Resist.* 2019;12:3903. doi: 10.2147/IDR.S234610.
  6. Feng, Z., K. Hayat, Z. Huang, et al., 2020. Knowledge, attitude, and practices of community pharmacy staff toward antimicrobial stewardship programs: a cross-sectional study from Northeastern China. 1-8.
  7. Al-Omari A., Al Mutair A., Alhumaid S., et al. The impact of antimicrobial stewardship program implementation at four tertiary private hospitals: results of a five-years pre-post analysis. *Comparative Study.* 2020;9:1–9. doi: 10.1186/s13756-020-00751-4.
  8. Howard P., Ashiru-Oredope D., Gilchrist M.J.E. Time for pharmacy to unite in the fight against antimicrobial resistance. *Pharma. J.* 2020;14:34.
  9. Franco B.E., Martínez M.A., Rodríguez M.A.S., et al. The determinants of the antibiotic resistance process. *Infect. Drug Resist.* 2009;2:1.
  10. Almaghaslah D., Alsayari A., Asiri R., et al. Pharmacy workforce in Saudi Arabia: challenges and opportunities: a cross-sectional study. *Int. J. Health Plann. Manage.* 2019;34:e583–e593. doi: 10.1002/hpm.2674.
  11. Hayat, K., M. Rosenthal, A. H. Gillani, et al., 2019. Perspective of key healthcare professionals on antimicrobial resistance and stewardship programs: a multicenter cross-sectional study from Pakistan. 10, 1520.
  12. Amer M.R., Akhras N.S., Mahmood W.A., et al. Antimicrobial stewardship program implementation in a medical intensive care unit at a tertiary care hospital in Saudi Arabia. *Comparative Study.* 2013;33:547–554. doi: 10.5144/0256-4947.2013.547.
  13. Hoxha, I., A. Malaj, B. Kraja, et al., 2018. Are pharmacists' good knowledge and awareness on antibiotics taken for granted? The situation in Albania and future implications across countries. 13, 240-245.
  14. Erku, D. A. J. I. p. o. i. d., 2016. Antimicrobial stewardship: a cross-sectional survey assessing the perceptions and practices of community pharmacists in Ethiopia. 2016.
  15. Abdulhak A.A.B., Al Tannir M.A., Almansor M.A., et al. Non prescribed sale of antibiotics in Riyadh, Saudi Arabia: a cross sectional study. *BMC Public Health.* 2011;11:1–5. doi: 10.1186/1471-2458-11-538.
  16. Hanafi, S., F. Poormalek, H. Torkamandi, et al., 2013. Evaluation of Community Pharmacists' Knowledge, Attitude and Practice towards Good Pharmacy Practice in Iran. 19-24.
  17. Hsu, J. J. B., 2020. How covid-19 is accelerating the threat of antimicrobial resistance. 369.
  18. Kapadia, S. N., E. L. Abramson, E. J. Carter, et al., 2018. The expanding role of antimicrobial stewardship programs in hospitals in the United States: lessons learned from a multisite qualitative study. 44, 68-74.
  19. Saleem Z., Hassali M.A.A., Godman B., et al. Sale of WHO AWaRe groups antibiotics without a prescription in Pakistan: a simulated client study. *J. Pharm. Policy Pract.* 2020 doi: 10.1186/s40545-020-00233-3.
  20. Khan, F. U., F. U. Khan, K. Hayat, et al., 2021. Knowledge, Attitude and Practice on Antibiotics and Its Resistance: Two Phases Mixed-Methods Online Study Among Pakistani Community Pharmacists to Promote Rational Antibiotics Use.

21. Phiri, M. N., M. Banda, S. Mudenda, et al., 2020. Coronavirus Disease 2019 (COVID-19): the role of pharmacists in the Fight against COVID-19 Pandemic.
22. Kotwani, A., K. J. T. M. Holloway and I. Health, 2014. Antibiotic prescribing practice for acute, uncomplicated respiratory tract infections in primary care settings in New Delhi, India. 19, 761-768.
23. Majumder M.A.A., Rahman S., Cohall D., et al. Antimicrobial stewardship: fighting antimicrobial resistance and protecting global public health. *Infect. Drug Resist.* 2020;13:4713.
24. Rehman I.U., Asad M.M., Bukhsh A., et al. Knowledge and practice of pharmacists toward antimicrobial stewardship in Pakistan. *Pharmacy (Basel)* 2018;6:116.
25. Saha, S. K., D. Kong, K. Thursky, et al., 2020. A nationwide survey of Australian general practitioners on antimicrobial stewardship: Awareness, uptake, collaboration with pharmacists and improvement strategies. 9, 310.
26. Saleem, Z., M. A. Hassali, F. K. Hashmi, et al., 2019. Antimicrobial dispensing practices and determinants of antimicrobial resistance: a qualitative study among community pharmacists in Pakistan.
27. Saha, S. K., L. Hawes and D. J. J. o. A. C. Mazza, 2019. Effectiveness of interventions involving pharmacists on antibiotic prescribing by general practitioners: a systematic review and meta-analysis. 74, 1173-1181.
28. Khan M.U., Hassali M.A.A., Ahmad A., et al. Perceptions and practices of community pharmacists towards antimicrobial stewardship in the state of Selangor, Malaysia. *PLOS ONE.* 2016;11:e0149623.
29. Tegagn G., Yadesa T., Ahmed Y.J.J.B.B. Knowledge, attitudes and practices of healthcare professionals towards antimicrobial stewardship and their predictors in Fitch Hospital. *J. Bioanal. Biomed.* 2017;9:91–97.
30. Mudenda, S., M. Hankombo, Z. Saleem, et al., 2020. Knowledge, Attitude, and Practices of Community Pharmacists on Antibiotic Resistance and Antimicrobial Stewardship in Lusaka, Zambia.
31. Palmer H.R., Weston J., Gentry L., et al. Improving patient care through implementation of an antimicrobial stewardship program. *Am. J. Health Syst. Pharm.* 2011;68:2170–2174.
32. Nasrin, D., P. J. Collignon, L. Roberts, et al., 2002. Effect of  $\beta$ -lactam antibiotic use in children on pneumococcal resistance to penicillin: prospective cohort study. 324, 28.
33. Klepser, M. E., A. J. Adams and D. G. J. H. s. Klepser, 2015. Antimicrobial stewardship in outpatient settings: leveraging innovative physician-pharmacist collaborations to reduce antibiotic resistance. 13, 166-173.
34. Saleem Z., Saeed H., Ahmad M., et al. Antibiotic self-prescribing trends, experiences and attitudes in upper respiratory tract infection among pharmacy and non-pharmacy students: a study from Lahore. *PLoS One.* 2016;11:e0149929.

35. Sarwar, M. R., A. Saqib, S. Iftikhar, et al., 2018. Knowledge of community pharmacists about antibiotics, and their perceptions and practices regarding antimicrobial stewardship: a cross-sectional study in Punjab, Pakistan. 11, 133.
36. Organization, W. H., 2014. The role of a pharmacist in encouraging prudent use of antibiotic medicines and averting antimicrobial resistance—a review of current policies and experiences in Europe, Copenhagen: WHO Regional Office for Europe.
37. Wong JR, Bauer KA, Mangino JE, Goff DA. Antimicrobial stewardship pharmacist interventions for coagulase-negative staphylococci positive blood cultures using rapid polymerase chain reaction. *Ann Pharmacother* 2012;46:1484–1490.
38. Rattanaumpawan P, Upanan P, Thamlikitkul V. A noninferiority cluster-randomized controlled trial on antibiotic postprescription review and authorization by trained general pharmacists and infectious disease clinical fellows. *Infect Control Hosp Epidemiol* 2018;39:1154–1162.
39. Garau J, Bassetti M. Role of pharmacists in antimicrobial stewardship programmes. *Int J Clin Pharm* 2018;40:948–952.
40. Jantarathaneewat K, Montakantikul P, Weber DJ, Nanthapisal S, Rutjanawech S, Apisarnthanarak A. Impact of an infectious diseases pharmacist-led intervention on antimicrobial stewardship program guideline adherence at a Thai medical center. *Am J Health Syst Pharm* 2022;79:1266–1272.
41. Antimicrobial resistance surveillance in Europe 2020–2022. European Centre for Disease Control and Prevention website. Published January 26, 2022. Accessed October 7, 2022.
42. McCormick JZ, Cardwell SM, Wheelock C, Wong CM, Vander Weide LA. Impact of ambulatory antimicrobial stewardship on prescribing patterns for urinary tract infections. *J Clin Pharm Ther* 2020;45:1312–1319.
43. Saha SK, Hawes L, Mazza D. Effectiveness of interventions involving pharmacists on antibiotic prescribing by general practitioners: a systematic review and meta-analysis. *J Antimicrob Chemother* 2019;74:1173–1181.
44. Saleh D, Abu Farha R, Alefishat E. Impact of educational intervention to promote Jordanian community pharmacists' knowledge and perception towards antimicrobial stewardship: pre–post interventional study. *Infect Drug Resist* 2021;14:3019–3027.
45. Apisarnthanarak A, Kwa AL, Chiu CH, et al. Antimicrobial stewardship for acute-care hospitals: an Asian perspective. *Infect Control Hosp Epidemiol* 2018;39:1237–1245.
46. US Department of Health and Human Services. Antibiotic resistance threats in the United States 2019. Centers for Disease Control and Prevention website. Published 2019. Accessed October 7, 2022.
47. Devchand M, Kirkpatrick CMJ, Stevenson W, et al. Evaluation of a pharmacist-led penicillin allergy de-labelling ward round: a novel antimicrobial stewardship intervention. *J Antimicrob Chemother* 2019;74:1725–1730.

48. ASHP statement on the pharmacist's role in antimicrobial stewardship and infection prevention and control. *Am J Health Syst Pharm* 2010;67:575–577.
49. Bishop PA, Isache C, McCarter YS, Smotherman C, Gautam S, Jankowski CA. Clinical impact of a pharmacist-led antimicrobial stewardship initiative evaluating patients with *Clostridioides difficile* colitis. *J Investig Med* 2020;68:888–892.
50. Barlam TF, Cosgrove SE, Abbo LM, et al. Implementing an antibiotic stewardship program: guidelines by the Infectious Diseases Society of America and the Society for Healthcare Epidemiology of America. *Clin Infect Dis* 2016;62:e51–e77.
51. Yam ELY, Hsu LY, Yap EP, et al. Antimicrobial resistance in the Asia Pacific region: a meeting report. *Antimicrob Resist Infect Control* 2019;8:202.
52. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* 2022;399:629–655.
53. Apisarnthanarak A, Lapcharoen P, Vanichkul P, Srisaeng-Ngoen T, Mundy LM. Design and analysis of a pharmacist-enhanced antimicrobial stewardship program in Thailand. *Am J Infect Control* 2015;43:956–959.
54. Arensman K, Dela-Pena J, Miller JL, et al. Impact of mandatory infectious diseases consultation and real-time antimicrobial stewardship pharmacist intervention on *Staphylococcus aureus* bacteremia bundle adherence. *Open Forum Infect Dis* 2020;7:ofaa184.
55. Yu K, Rho J, Morcos M, et al. Evaluation of dedicated infectious diseases pharmacists on antimicrobial stewardship teams. *Am J Health Syst Pharm* 2014;71:1019–1028.
56. Xu S, Wang X, Song Z, Han F, Zhang C. Impact and barriers of a pharmacist-led practice with computerized reminders on intravenous to oral antibiotic conversion for community-acquired pneumonia inpatients. *J Clin Pharm Ther* 2021;46:1055–1061.
57. Hasegawa S, Tagashira Y, Murakami S, et al. Antimicrobial time-out for vancomycin by infectious disease physicians versus clinical pharmacists: a before–after crossover trial. *Open Forum Infect Dis* 2021;8:ofab125.
58. Zhou Y, Ma LY, Zhao X, Tian SH, Sun LY, Cui YM. Impact of pharmacist intervention on antibiotic use and prophylactic antibiotic use in urology clean operations. *J Clin Pharm Ther* 2015;40:404–408.
59. Jantarathaneewat K, Apisarnthanarak A, Limvorapitak W, Weber DJ, Montakantikul P. Pharmacist-driven antibiotic stewardship program in febrile neutropenic patients: a single-site prospective study in Thailand. *Antibiotics (Basel)* 2021;10:456.
60. Zhang J, Li X, He R, et al. The effectiveness of clinical pharmacist-led consultation in the treatment of infectious diseases: a prospective, multicenter, cohort study. *Front Pharmacol* 2020;11:575022.
61. Xu J, Huang J, Yu Y, et al. The impact of a multifaceted pharmacist-led antimicrobial stewardship program on antibiotic use: evidence from a quasi-experimental study in the department of vascular and interventional radiology in a Chinese tertiary hospital. *Front Pharmacol* 2022;13:832078.

62. Chaononghin S, Jantarathaneewat K, Weber DJ, Warren DK, Apisarnthanarak A. Impact of antibiotic heterogeneity by periodic antibiotic monitoring and supervision strategy at two units with different prevalences of multidrug-resistant organisms. *Infect Control Hosp Epidemiol* 2021. doi:10.1017/ice.2021.231.
63. Du Y, Li J, Wang X, et al. Impact of a multifaceted pharmacist-led intervention on antimicrobial stewardship in a gastroenterology ward: a segmented regression analysis. *Front Pharmacol* 2020;11:442.
64. Takito S, Kusama Y, Fukuda H, Kutsuna S. Pharmacist-supported antimicrobial stewardship in a retirement home. *J Infect Chemother* 2020;26:858–861.
65. Nampoothiri V, Sudhir AS, Joseph MV, et al. Mapping the implementation of a clinical pharmacist-driven antimicrobial stewardship programme at a tertiary-care centre in South India. *Antibiotics (Basel)* 2021;10(2):220.
66. Zhou H, Liu L, Sun X, et al. The impact of pharmacist intervention on prophylactic antibiotics use in orthopedic surgery at a hospital in China. *Medicine (Baltimore)* 2021;100:e28458.
67. Suh Y, Ah YM, Chun HJ, et al. Potential impact of the involvement of clinical pharmacists in antimicrobial stewardship programs on the incidence of antimicrobial-related adverse events in hospitalized patients: a multicenter retrospective study. *Antibiotics (Basel)* 2021;10:853.
68. Song YJ, Kim M, Huh S, et al. Impact of an antimicrobial stewardship program on unnecessary double anaerobic coverage prescription. *Infect Chemother* 2015;47:111–116.
69. Li Z, Cheng B, Zhang K, et al. Pharmacist-driven antimicrobial stewardship in intensive care units in East China: a multicenter prospective cohort study. *Am J Infect Control* 2017;45:983–989.