

Self-care Practices of Saudi Diabetic Foot Patients in 2024

Ebtihal Othman Barnawi¹, Ghofran Ahmed Hosein¹, Nada Abdulrhman Alsaab¹, Abdualmajed Muhyi Alqarni², Thamer abdullah Alotaibi², Salem Naji Alharbi², Abdulaziz Hasan Almalki², Abrar Othman Hawsawi³, Aysha Mohd Said Alomari⁴, Mashaal Ali Najmi⁵

1. Nursing Specialist, maternity and children hospital, Saudi Arabia.
2. Health Education Specialist, King Faisal Hospital, Saudi Arabia.
3. NURSE, KING ABDULAZIZ HOSPITAL, Saudi Arabia.
4. Nursing technician, Makhwah1 Primary Health Care Center, Saudi Arabia.
5. Nursing technician, maternity and children hospital, Saudi Arabia.

ABSTRACT

Background: Diabetic foot is a foot that shows any pathology that consequence in a straight line from diabetes mellitus or any long -term complications of DM which lead to diabetic foot amputation when delaying in treatment. **Aim of the study:** Was to assess self-care practices of Saudi diabetic foot patients. **Research design:** A descriptive research design was utilized to conduct this study. **Setting:** This study was conducted at Diabetic Patient Clinic of Taif hospital, Saudi Arabia. **The sample:** A systematic sample of 200 diabetic foot patients over 50 years old. **Tools:** One tool was used in this study. **1):** A structured interviewing questionnaire which consists of four parts. **Part (1):** Demographic characteristics of diabetic foot patients, **Part (2):** Medical history of diabetic foot patients. **Part (3):** Patients' knowledge about diabetes Mellitus and diabetic foot. **Part (4):** Diabetic patients' reported practices toward diabetic foot self- care. **Results:** 46.5% of studied patients aged from 60<70 years old, 47.5% of them had basic education, and 50% of them were male. 52% of studied patients had diabetes mellitus from 1 to less than 5 years and 35% of them were visiting The Diabetic Clinic every three weeks Also 49.5% of studied patients had poor total knowledge level, while; 51% of them had satisfactory total reported self-care practices regarding diabetic foot care. **Conclusion:** There was highly statistically positive correlation between total knowledge of studied patients, their total reported practices regarding care of diabetic foot. **Recommendations:** Develop health educational program for diabetic foot patients to increase their knowledge and self -care practices regarding diabetic foot.

KEYWORDS: Diabetic foot, Patient practices, Self- care

Introduction

Diabetes Mellitus (DM), is a cluster of metabolic disease which characterized by high glucose levels, resulting in defect in insulin secretion, insulinaction or both and disturbance of carbohydrate, fats and protein metabolism. Diabetes can be classified into type 1 diabetes, type 2 diabetes, gestational diabetes and specific types of diabetes due to other causes (Monogenetic diabetes syndromes, disease of exocrine pancreas and drug or chemical induced diabetes). Over the long- term high glucose levels are associated with damage to the body and failure of various organs and tissues (**Chakraborty et al., 2021**).

The most common known consequence of diabetes mellitus is a diabetic foot ulcer, which is characterized by deep tissue lesions linked to peripheral vascular disease and neurological abnormalities in the lower extremities. It is a full-thickness wound beneath the ankle in a diabetic patient that penetrates the dermis, the deep vascular and collagenous inner layer of the skin. Diabetic foot ulcers are divided into six wound classes primarily according to the depth of the wound. Grade 0 has no ulcer but the foot is susceptible to developing one; Grade 1 has a superficial ulcer; Grade 2 has a deep infection but does not involve the bone; Grade 3 has osteomyelitis; Grade 4 has localized gangrene; and Grade 5 has gangrene of the entire foot (**Tuha et al.,2021**).

In their lifetime, 15% of diabetes individuals will get foot ulcers, which, if left untreated, can progress to chronic ulcers, chronic osteomyelitis, and in 85% of instances, amputation. Every 20 seconds, someone in the world loses a limb due to complications from diabetes, and more than half of those individuals will pass away within five years following the procedure, according to the International Working Group on Diabetic Foot. By practicing foot self-care at home, the majority of foot or leg amputations can be avoided or at least postponed (**Mekonen et al., 2020**).

General diet, foot care, blood glucose monitoring, diabetes medication, and physical activity are specific items for measuring the self-care behaviors of diabetic patients. Self-care practices are linked to the quality of life of patients with diabetes, and it is important to know related factors to improve it. Economic status, work, the presence of caregivers, self-efficacy, coping strategy, and family involvement all have an impact. Since type 2 diabetes is a degenerative condition and most patients are elderly, family members help them manage their condition. Patients' self-care practices were positively impacted by their perceptions of family support. (**Kim et al., 2020**).

Health care professionals who actively participate in the prevention and early identification of diabetes and its complications include public health members, particularly nurses. They may play a part in patient care, health care, community education, health systems management, and enhancing people's quality of life. They carry out their educational function in the areas of diabetic foot prevention, foot care, and injury prevention. In the care dimension, nurses are in charge of foot care, dressing, using new technologies, and identifying any changes in skin and foot sensation early (**Doenges et al., 2019**).

Significance of the Study:

World health organization has reported that Saudi Arabia ranks the second highest in the middle east and the seventh in the world for the rate of diabetes. In fact, diabetes has approximately registered a ten-fold increase in the past three eras in Saudi Arabia. Diabetic foot ulcer was found to be the main cause of lower limb amputation in Saudi Arabia

Aim of the Study:

This study aimed to assess self-care practices of Saudi diabetic foot patients

Research Questions:

- 1- What is the knowledge of diabetic patient regarding diabetic foot?
- 2- What are the self-care practices of diabetic patients regarding diabetic foot?
- 3-Is there a relation between socio- demographic characteristics of diabetic patients and their self-care practices regarding diabetic foot?
- 4- Is there a relation between diabetic patients' self-care practices regarding diabetic foot, and their knowledge?

Subjects and Methods:

Research design:

A descriptive study design was utilized to conduct this study.

Setting:

This study was conducted at Diabetic Foot Outpatient Clinic at Taif Hospital, Saudi Arabia.

Sampling:

A systemic random sample was used in the study, the total number of patients at the previous mentioned setting were 960 patients. Patients were placed in a list with a serial number, and then the starting point was randomly selected, after that each patient was selected every 5 patients to be in the sample. The total number of the sample was 200 patients with inclusion criteria, patients age was over 50 years old.

Tools of Data Collection:

Tool (I): A structured interviewing questionnaire:

It was developed by the researchers based on reviewing related literatures and it was written in simple clear Arabic language. It is comprised of four parts.

First part: Was concerned with demographic characteristics of diabetic foot patients. It comprised of 8 questions (Sex, age, marital status, educational level, occupation,

monthly income, family type).

Second part: Was concerned with diabetic patients' medical history. It comprised of 9 questions (onset of diabetes, frequency of visiting diabetic clinic, number of hospitalization due to diabetes mellites, medical intervention to diabetic foot, current complains, types of treatment, presence of other chronic diseases, factors that precipitate of diabetic foot, stages of diabetic foot).

Third part: It consisted of two sections: section (1): Was concerned with studied patients' knowledge regarding diabetes mellitus. It comprised of 7 questions (meaning of diabetes, types, causes, risk factors, manifestation, diagnosis, complications).

Section (2): Was concerned with studied patients' knowledge regarding diabetic foot. It comprised of 9 questions (meaning of diabetic foot, causes, manifestations, stages, complications, treatment, prevention, medical care, importance of performance of complete and periodic medical care for diabetic foot).

Scoring System:

The scoring system for studied patients' knowledge was calculated as follows: (2) score for correct complete answer, and (1) score for correct in complete answer, while (0) for don't know. For each area of knowledge, the score of items was summed-up and the total divided by the number of the items, giving a mean score for the part. These scores were converted into a present score. The total score of knowledge

=32points. The total score was considered good when score of total knowledge $\geq 75\%$ (≥ 24 point), and considered average if it equals $50 < 75\%$ ($16 < 24$ points), while considered poor when the total score was $< 50\%$ (< 16 point).

Fourth part: Was concerned with diabetic patients' reported practices toward diabetic foot self-care through asking questions regarding care of foot. which divided into six sections: **Personal hygiene:** It comprised of seven questions, **Foot care:** It comprised of eleven questions, **Nutrition:** It included eleven questions, **Exercise:** It included twelve questions, **dealing with difficult walking:** It included four questions and **Therapeutic regimen and diabetes follow up:** It included eleven questions.

Scoring System:

The reported self-care practices was scored by two level of answers: done or not done. These were respectively 1.0. The scores of the items were summed up and the total divided by the number of the items, giving a mean score for the part. These were converted into a percent score. The total practice score =56

The total practices scores were satisfactory if the score of the total practices $\geq 60\%$ (≥ 33 points), while considered unsatisfactory if it is $< 60\%$ (< 33 points).

Reliability of the Tool:

Reliability of the tool was applied by the researchers for test the internal consistency of the tool by administration of the same tools to the same subjects under similar condition on one or more occasion. Answers from repeated testing were compared (test-re-test reliability). The reliability was done Cronbach Alpha coefficients test which revealed that the tool consisted of relatively homogenous items as indicated by

Ebtihal Othman Barnawi, Ghofran Ahmed Hosein, Nada Abdulrhman Alsaab, Abdualmajed Muhyi Alqarni, Thamer abdullah Alotaibi, Salem Naji Alharbi, Abdulaziz Hasan Almalki, Abrar Othman Hawsawi, Aysha Mohd Said Alomari, Mashael Ali Najmi

moderate to high reliability of each tool. The internal consistency of the knowledge was 0.721, while: reliability of practices was 0.834.

Content Validity of the Tool:

Content validity of the tool was done by experts in the field who reviewed the tools for clarity, relevance, comprehensiveness, applicability and give their opinion.

Ethical Considerations:

All ethical issues were assured; oral consent has been obtained from each patient with diabetic foot before conducting the interview and give them a brief orientation to the purpose of the study. They were also reassured that all information will be treated confidentially and used only for the purpose of the study. The patients had right to withdraw from the study at any time without giving any reasons.

Pilot Study:

The pilot study was carried out on (20) patients with diabetic foot which represented 10 % of the total sample size (200). The pilot study was aimed to assess the tool clarity, applicability and time needed to fill each sheet, completing the sheet consumed about 30 minutes. No modifications were done, so the pilot study sample was included in the total sample.

Field Work:

The actual field was carried out over a period of 6 months from the beginning of February 2024 to the end of July 2024.

Statistical Analysis:

All data collected were organized, tabulated and analyzed by using the statistical test, The data were analyzed by using the Statistical Package for Social Science (SPSS), which was applied to calculate frequencies and percentages for qualitative descriptive data and chi-square coefficient χ^2 was used for relation tests, mean and Standard Deviation (SD) was used for qualitative data, person correlation coefficient (r) was used for correlation analysis and degree of significance was identified.

Statistically significance was considered at:

- Highly statistically significant when P-value >0.001 .
- Statistically significant result when P-value <0.05 .
- Not significant result when P-value >0.05 .

Results:

Table (1): Shows that; 46.5% of studied patients aged from 60 to less than 70 years old with Mean \pm SD 61.52 \pm 6.34 and 48% of them were married. Concerning educational level; 47.5% of them had basic education, 49% of them were house wives / not working and 63% of them had enough income. In addition, 57% of them had extended family

Demographic characteristics	No	%
Gender:		
Male	100	50.0
Female	100	50.0
Age:		
50<60	87	43.5
60<70	93	46.5
70<80	20	10.0
Mean \pmSD	61.52\pm6.34	
Marital status:		
Married	96	48.0
Widow	87	43.5
Divorced	17	8.5
Educational level:		
Can't read or write	47	23.5
Basic education	95	47.5
Intermediate education	42	21.0
University education or more	16	8.0
Occupation:		
Employee	38	19.0
Free works	43	21.5
Housewives / Not working	98	49.0
Retired	21	10.5
Monthly income:		
Enough and save	26	13.0
Enough	126	63.0
Not enough	48	24.0
Family Type:		
Nuclear family	86	43.0
Extended family	114	57.0

Table (2): Reveals that; 52% of studied patients had diabetes mellitus from 1 to less than 5 years and 35% of them were visiting the diabetic clinic every three weeks while; 43% of them never hospitalized due to Diabetes Mellitus. Concerning the medical intervention to the diabetic foot, 40% of studied patients had dressing for diabetic foot wound and 50% of them complained from sores on the foot. In addition, 60.5% of them treated with subcutaneous insulin and 53.5% of them had chronic diseases as hypertension.

Medical history	No	%
Onset of DM:		
1<5 years	104	52.0
5<10 years	86	43.0
>10 years	10	5.0
Frequency of visiting to diabetic clinic:		

Ebtihal Othman Barnawi, Ghofran Ahmed Hosein, Nada Abdulrhman Alsaab, Abdualmajed Muhyi Alqarni, Thamer abdullah Alotaibi, Salem Naji Alharbi, Abdulaziz Hasan Almalki, Abrar Othman Hawsawi, Aysha Mohd Said Alomari, Mashael Ali Najmi

Every week	24	12.0
Every two weeks	62	31.0
Every three weeks	70	35.0
Every month or more	44	22.0
Number of hospitalization due to DM:		
Never	86	43.0
Once	77	38.5
Twice ore more	35	17.5
Medical intervention to diabetic foot:		
Dressing for diabetic foot wound	80	40.0
Deep cleaning to diabetic foot	70	35.0
Surgical intervention for diabetic foot at the operating room	41	20.5
Don't need medical intervention for diabetic foot	9	4.5
*Current complains:		
Sores on the foot	100	50.0
Foot dryness that results in cracks	47	23.5
Loss of sensation with pain or cold	83	41.5
Change in the shape of the foot due to wearing tight shoes for long period	40	20.0
Types of treatment:		
Oral tablets drugs	79	39.5
Subcutaneous Insulin	121	60.5
*Presecence of other chronic diseases:		
Hypertension.	107	53.5
High cholesterol level	37	18.5
Peripheral atherosclerosis.	4	2.0
Cardiac disease	55	27.5
Nothing	18	9.0
*Factors that precipitate diabetic foot:		
High blood glucose level for a long time.	92	46.0
Presence of a medical history of the patient or family for diabetic foot.	47	23.5
Appearance of foot ulcers.	45	22.5
Weight gain leads to excessive pressure on the foot.	30	13.0
Lack of foot care	95	23.5
Stages of diabetic foot:		

Natural diabetic foot	68	34.0
Diabetic foot with an increase in the thickness of the skin and beginning of a superficial ulcer	86	43.0
Ulcerated diabetic foot with fingers and edges of the foot	44	18.5
Diabetic foot with deep ulcers and osteomyelitis	40	4.5

Figure (1): Shows that; 49.5% of studied patients had poor total knowledge level, and 37% of them had average total knowledge level. While ;13.5% of them had good total knowledge level regarding diabetic foot.





Figure (2): shows that 51% of studied patients had satisfactory total practices regarding diabetic foot care. While; 49% of them had unsatisfactory total practices regarding diabetic foot care.



Ebtihal Othman Barnawi, Ghofran Ahmed Hosein, Nada Abdulrhman Alsaab, Abdualmajed Muhyi Alqarni, Thamer abdullah Alotaibi, Salem Naji Alharbi, Abdulaziz Hasan Almalki, Abrar Othman Hawsawi, Aysha Mohd Said Alomari, Mashael Ali Najmi

Table (3) Demonstrates that, there were no statistically significant relation between patients' total reported practices about diabetic foot care and all their socio-demographic characteristics except for their educational level, there was statistically significant relation ($p < 0.05^*$).

Socio-demographic characteristics	Total self-care practices				X	p-value
	Unsatisfactory (n=98)		Satisfactory (n=102)			
Gender:						
Male	48	49.0	52	51.0	0.08	0.777
Female	50	51.0	50	49.0		
Age:						
50<60	44	44.9	43	42.2	3.23	0.199
60<70	48	49.0	45	44.1		
70<80	6	6.1	14	13.7		
Marital status:						
Married	46	46.9	50	49.0	1.845	0.397
Widow	46	46.9	41	40.2		
Divorced	6	6.1	11	10.8		
Educational level:						
Can't read or write	25	25.5	22	21.6 12	.163	0.007*
Basic education	55	56.1	40	39.2		
Intermediate education	15	15.3	27	26.5		
University education or more	3	3.1	13	12.7		
Occupation:						
Employee	12	12.2	26	25.5	6.742	0.15
Free works	23	23.5	20	19.6		
Housewives /not working	42	42.9	34	33.3		
Retired	9	9.2	12	11.8		
Monthly income						
Enough and save	13	13.3	13	12.7	0.253	0.881
Enough	63	64.3	63	61.8		
Not enough	22	22.4	26	25.5		

Family Type						
Nuclear family	43	43.9	43	42.2	0.06	0.806
Extended family	55	56.1	59	57.8		

Table (4) Reveals that there was highly statistically positive correlation between total knowledge of studied patients and their total self-care practices about diabetic foot ($p < 0.001^{**}$)

Total knowledge	Total self-care				χ^2	p-alue
	Unsatisfactory (n=98)		Satisfactory (n=102)			
Poor (n=99)	55	56.1	44	43.	14.735	.001**
Average (n=74)	39	39.8	35	34.		
Good (n=27)	4	4.1	23	22.		

Discussion:

Diabetes mellitus (DM) is one of the major diseases leading to death worldwide. It is a condition of high sugar or glucose concentration in blood resulting from abnormal insulin secretion, insulin action, or both. The chronic hyperglycemia of diabetes is linked to long-term damage, dysfunction, and failure of various organs, particularly the eyes, kidneys, nerves, heart, and blood vessels. Diabetic foot ulcer is one of the major complications caused by diabetes mellitus (DM). Foot ulcers occur due to damaged skin tissues under the big toes and on the feet plantar. Foot ulcers cause the underlying skin layer to be exposed, affecting the feet to the bone as a result of the complication of uncontrolled diabetes. Improper and late handling of patients with a diabetic foot can result in an amputation of the foot (**Munadi et al., 2022**).

The findings of the current study were discussed under the following consequences: demographic characteristics of the studied patients, medical history of them, knowledge regarding diabetes mellitus and diabetic foot, reported practices of studied patients regarding diabetic foot care.

According to demographic characteristics of the studied patients with diabetic foot, the present study findings showed that nearly half of studied patients age 60 to less than 70 years old with mean age 61.52+6.34 years. This finding was inconsistent with **Al-Jarallah et al. (2018)**, who studied "Knowledge and practice of diabetic foot care among diabetic patients in Aseer region, Saudi Arabia"(n=351) and founded that more than two fifth (42.2%) of studied patients aged more than 60 years old. Also, this finding disagreed with **Dedefo et al., (2019)**, who reported that more than tenth (12.7%) aged >60 years old.

Regarding patients' gender, this current study showed that half of patients were males. This finding agreed with **Mekonen & Demssie (2022)**, who found that half (50.5%) of the studied patients were male.

Concerning onset of diabetes mellitus, the present study showed that half of studied

Ebtihal Othman Barnawi, Ghofran Ahmed Hosein, Nada Abdulrhman Alsaab, Abdualmajed Muhyi Alqarni, Thamer abdullah Alotaibi, Salem Naji Alharbi, Abdulaziz Hasan Almalki, Abrar Othman Hawsawi, Aysha Mohd Said Alomari, Mashael Ali Najmi

patient had diabetes from 1 to less than 5 years. This finding was consistent with **Sari et al., (2020)**, who founded that half (49.8%) of studied patients had diabetes from 1 to less than 5 years. However, this finding was supported by **Mahmood et al., (2019)**, who found that nearly two fifth (38.2%) of them had diabetes mellites from 1 to less than 5 years.

According to stages of diabetic foot, the current study revealed that more than majority of studied patients had beginning of superficial ulcer. The finding disagreed with **pourkazemi et al., (2020)**, who found that more than three quarters (76.8%) of the studied patients had no diabetic foot ulcers.

In relation to patients' knowledge regarding diabetic foot, the results of current study clarified that half of studied patients had poor total knowledge level, and more than one third of them had average total knowledge level. While 13.5% of them had good total knowledge level regarding diabetes mellitus and diabetic foot. This might be due to the low schooling levels of the patients that contributed to the poor knowledge. This result agreed with **Wazqar et al., (2021)**, who conducted a study "Assessment of knowledge and foot self-care practices among diabetes mellitus patients in a tertiary care Centre in Makkah, Saudi Arabia"(n=409) and founded that nearly three quarters (72.4%) of the participants had a poor level of knowledge whereas only 4.2% expressed a good level of knowledge. Furthermore, our study findings were similar to the results of **Khan et al., (2020)**, who showed that most people with diabetes had poor knowledge and negative behaviors toward foot -care.

In contrast, this result disagrees with **Asmelash et al, 2019**, and **Alsous et al., (2019)** who showed that more than three fifth (62%) of the participants, respectively had a good knowledge about DM. This might be due to the fact that there are sociodemographic variations across the countries, the difference in health education, sample size and access to sources of information like television, radio and newspaper.

According to total self-care reported practices, the current study showed that more than half of studied patients had satisfactory total self-care reported practices. These findings were disagreed with **Abu-Elenin et al., (2018)**, who mentioned that more than three fifth of studied patients (62.2%) had inadequate self-foot care practice level. This might be due to patients have an elementary role in promoting proper diabetic foot self-care.

Regarding to relation between demographic characteristic of diabetic patients and their self-care reported practices regarding diabetic foot. the results of present study clarified that, there were no statistically significant relation between patient's self-care reported practices about diabetic foot care and all their demographic characteristic except their educational level there was statistically significant relation ($p>0.05$),

The findings of current study disagreed with **Raj et al., (2021)**, who showed that there is a significant association between practice regarding foot self-care behavior among diabetic patients and their demographic variables such as educational status, family history, monthly income and type of anti-diabetic medication.

Owing to correlation between total knowledge and total practices regarding diabetic

foot care. the finding of current study reported that there was highly statistically positive correlation between total knowledge of studied patients and their total self-care practices about diabetic foot ($p < 0.001^{**}$).

This result was consistent with **Jing et al., 2022**, who showed that there is a significant positive correlation between foot care knowledge and foot care practices ($p < 0.001$). Increasingly, the result of present study agreed with (**Alharbi and Sulaiman, 2022**) who demonstrated that a strong statistical association was detected between patients' good knowledge of diabetic foot self-care and good practice ($P\text{-value} = 0.001$).

Conclusion:

Approximately less than half of studied patients had poor total knowledge level about diabetic foot. While; more than half of them had satisfactory total reported practices regarding foot care. occupation. There was no statistically significant relation between total patients' self-care reported practices about diabetic foot care and all their demographic characteristics except for their educational level, there was statistically significant relation. There was highly statistically positive correlation between total knowledge of studied patients and their total self-care reported practices regarding foot care.

Recommendations:

- 1- Develop health educational program for diabetic foot patients to increase their knowledge and self-care practices regarding diabetic foot.
- 2- Regular follow up care for diabetic foot patients in outpatient's clinics by specialized team to prevent complication of diabetic foot. 3- Further researches are required to study diabetes, diabetic foot and prevention of diabetic foot amputation on a large sample.
- 4- Colored illustrated booklet should be available and distributed to each patient with diabetic foot about self-care management practices regarding diabetic
- 5- Encourage the use of different modalities of Telemedicine (TM) as a communication tool between caregivers across the health care sectors and help patients with DFUs opens up for multi-sectoral and interdisciplinary close follow-up .

References:

- Abu-Elenin M, Elshoura A, and Alghazaly M. (2018).** Knowledge, practice and barriers of foot self-care among diabetic patients at Tanta university hospitals, Egypt. The Egyptian journal of community medicine; 36(4):94-102.
- Alharbi, O., and Sulaiman, A. (2022).** Foot care knowledge, attitude and practices of diabetic patients: A survey in Diabetes health care facility. Journal family med prim care; 11:3816-23.
- Al-Jarallah, M., Alqahtani, M., Alshahrani, S., Alshahrani, N., and Alshehri, A. (2020).** Knowledge and practice of diabetic foot care among diabetic patients in Aseer region, Saudi

Ebtihal Othman Barnawi, Ghofran Ahmed Hosein, Nada Abdulrhman Alsaab, Abdualmajed Muhyi Alqarni, Thamer abdullah Alotaibi, Salem Naji Alharbi, Abdulaziz Hasan Almalki, Abrar Othman Hawsawi, Aysa Mohd Said Alomari, Mashael Ali Najmi

Arabia. *International journal medecin Developing countries*; 4(1): 95-100.

Alsous, M., Abdel Jalil, M., Odeh, M., Al Kurdi, R., and Alnan M. (2019). Public knowledge, attitudes and practices toward diabetes mellitus: A cross-sectional study from Jordan. *Plos one journal*:143: 0214479.

Asmelash, D., Abdu, N., Tefera, S., Baynes, W., and Derbew, C. (2019). Knowledge, attitude, and practice towards glycemic control and its associated factors among diabetes mellitus patients. *Journal of diabetes research*:2019. Available at <https://doi.org/10.1155/2019/2593684>.

Chakraborty, R., Sen, S., Kalita, P., and Pathak, P. (2021). Diabetes mellitus and COVID-19 understanding the association in light of current evidence. *World journal of clinical cases*; 9(28): 8327.

Dedefo, G., Ejeta, M., Wakjira, B., Mekonen, F., and Labata, B. (2019). Self-care practices regarding diabetes among diabetic patients in West Ethiopia. *BMC research notes*; 12(1):1-7.

Doenges, E., Moorhouse, F., and Murr, C. (2019). Nursing care plans: Guidelines for individualizing client care across the life span. **Jing, J., Azmil, S., Sean, C., Fuen, S., Choo, G., Patel, A., and Mawardi, M. (2022).** Foot care knowledge and self-care practices among diabetic patients in Penang: A primary care study. *Med journal malaysia*; 77 (2):224-231.

Khan, L., Al-Nami, Q., Al-Gaseer, H., and Al-Neami, A. (2020). Foot self-care knowledge and practice evaluation among patients with diabetes. *Journal of diabetology*; 11:169-74.

Kim, J., and Han, S. (2020). Factors related to self-care behaviors among patients with diabetic foot ulcers. *Journal of Clinical Nursing*;29(9-10): 1712-1722.

Mahmood, K., Parkar, Z., Mustafa, T., Mustafa, S., Makin, A., and Alawadi, F., (2019). Diabetic foot self care: Awareness and practice among type 2 diabetic patients in primary healthcare centers, Dubai Health Authority. *Journal Community Med Public Health*; 6:1 7.

Mekonen, G., and Demssie, G. (2022). Preventive foot self care practice and associated factors among diabetic patients attending the university of Gondar comprehensive specialized referral hospital, Northwest Ethiopia, 2021. *Journal of BMC Endocrine Disorders*, 22(124):1-11.

Mekonnen, K., Abate, K., and Tegegne, T. (2020). Knowledge, Attitude, and practice toward lifestyle modification among diabetes mellitus patients attending the university of Gondar comprehensive specialized hospital northwest, Ethiopia. *Diabetes metabolic syndrome obesity*; 13:1969-1977.

Munadi, K., Saddami, K., Oktiana, M., Roslidar, R., Muchtar, K., Melinda, M., and Arnia, F. (2022). A deep learning method for early detection of diabetic foot using decision fusion and thermal images. *Applied Sciences*; 12(15): 7524. *Oxford Learner's Dictionaries (2023):* Definition of compliance noun from the *Oxford Advanced Learner's Dictionary*.

Pourkazemi, A., Ghanbari, A., Khojamli, M., Balo, H., Hemmati, H., Jafaryparvar, Z., and Motamed, B. (2020). Diabetic foot care: knowledge and practice. *BMC endocrine disorders*;20: 1-8.

Raj, K., Aishwarya., Mathew, S., Sunil, R., Mol, A., Amrutha, P. (2021). Knowledge and practice regarding foot self-care behavior among patients with diabetes mellitus. *International journal of evidence-based nursing*; 4:1.

Sari, Y., Upoyo, S., Isworo1, A., Taufik, A., Sumeru, A., Anandari, D., and Sutrisna, E. (2020). Foot self-care behavior and its predictors in diabetic patients in Indonesia. *BMC research notes*; 13(38):1-6.

Tuha, A., Getie Faris, A., Andualem, A., and Mohammed, S. (2021). Knowledge and practice

on diabetic foot self-care and associated factors among diabetic patients at dessie referral hospital, northeast Ethiopia: mixed method. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*; 1203:1214.

Wazqar, A., Baatya, M., Lodhi and Khan, A. (2021). Assessment of knowledge and foot self-care practices among diabetes mellitus patients in a tertiary care center in Makkah, Saudi Arabia: a cross-sectional analytical study. *Pan African Medical Journal*;40(123): 10.11604.