

# Epilepsy and Seizures Management in Emergency Department: A Nursing Perspective

**ABDULLAH AHMED MOHAMMED ALKHALDI <sup>1</sup>, MAITHA KHALIL LAFI ALKHAIBARI <sup>2</sup>, sultan Salem Sakhani Alruwaili <sup>3</sup>, Abdullah Fihat Ghadeer Alanazi <sup>4</sup>, Sultan Marji Amash Alanazi <sup>5</sup>, Amani Saud Al-Daghmani <sup>6</sup>, Tahani Saud Al-Daghmani <sup>7</sup>, Jamal Salamah Hamoud Albalawi <sup>8</sup>, Ahmed Nasser Ahmed Alasiri <sup>9</sup>, Abdulkarim Mulhi Mulhid Alanazi <sup>10</sup>, Omar Nadaa Huosein Alanazi <sup>11</sup>, NAIF KHALAF AWAD ALSWINA <sup>12</sup>, Dhafer Mesfer Ali Al Awn <sup>13</sup>**

- 1- *Emergency Medical Services, Emergency Management and Medical Coordination in Arar, Saudi Arabia*
- 2- *Emergency Medical Services, Emergency Management and Medical Coordination in Arar, Saudi Arabia*
- 3- *Emergency Medical Services, Emergency Management and Medical Coordination in Arar, Saudi Arabia*
- 4- *Emergency Medical Services, Emergency Management and Medical Coordination in Arar, Saudi Arabia*
- 5- *Emergency Medical Services, Emergency Management and Medical Coordination in Arar, Saudi Arabia*
- 6- *Nursing Technician Prince Miteb Bin Abdulaziz Hospital Sakaka Kingdom of Saudi Arabia*
- 7- *Nursing Specialist, King Abdulaziz Specialist Hospital, Sakaka, Kingdom of Saudi Arabia*
- 8- *Nursing and Midwifery, Mental Health Hospital, Tabuk, Saudi Arabia*
- 9- *Emergency Medical Services, Psychiatric Hospital in Tabouk, Saudi Arabia*
- 10- *Emergency Medical Services, Medical Transport Emergency Management Arar, Saudi Arabia*
- 11- *Paramedic, Emergency, maternity and children Hospital, Arar, Saudi Arabia*
- 12- *Emergency Medical Services, Medical Transport Emergency Management Arar, Saudi Arabia*
- 13- *Emergency Medical Services, Psychiatric Hospital in Tabouk, Saudi Arabia*

## **Abstract:**

Effective management of seizure disorders requires a comprehensive approach that combines pharmacological treatment with patient education and support. Nurses play a crucial role in administering anti-seizure medications, monitoring patients for side effects, and ensuring adherence to prescribed treatment regimens. Understanding the different types of seizures—such as focal, generalized, and absence seizures—enables nurses to tailor their care strategies to individual patient needs. Additionally, nurses must provide patients and their families with education on seizure triggers, the importance of medication compliance, and the proper response to seizures, which can significantly improve the quality of life for those affected by these disorders. In the nursing perspective, creating a safe environment for patients with seizure disorders is essential. This includes implementing safety precautions like padded bedrails and educating patients about avoiding known triggers. Nurses often collaborate with interdisciplinary teams to develop comprehensive care plans that encompass not only medical

management but also psychological support, as seizure disorders can lead to anxiety and stigma. Regular assessments and open communication with patients about their concerns are vital to ensure that care remains patient-centered and effective. By fostering a supportive environment and advocating for the needs of patients, nurses can significantly contribute to the overall management and well-being of individuals living with seizure disorders.

**Keywords:** Seizure disorders, Nursing care, Anti-seizure medications, Patient education, Safety precautions, Triggers, Interdisciplinary collaboration, Psychological support, Patient-centered care, Quality of life.

### **Introduction:**

Seizure disorders, commonly known as epilepsy, represent a significant public health challenge across the globe. Characterized by recurrent, unprovoked seizures, these neurological conditions can profoundly affect the quality of life for individuals living with the disorder, as well as for their families. With an estimated prevalence of 1-2% of the population worldwide, epilepsy necessitates comprehensive management strategies that encompass not only medical but also therapeutic and psychosocial support interventions. Nurses, often serving as frontline healthcare providers, play a pivotal role in the management of seizure disorders, demonstrating the critical intersection of clinical expertise, patient education, and holistic care in this field [1].

In the past few decades, there has been a paradigm shift in the understanding and management of seizure disorders that emphasizes a patient-centered approach. Historically, epilepsy was shrouded in stigma and often poorly managed due to misconceptions about its etiology and implications. However, the integration of advanced diagnostics, evolving pharmacological treatments, and robust support networks has propelled the field toward more effective management strategies. Nursing professionals possess a unique vantage point in the healthcare continuum. They are integral to developing care plans, providing education, advocating for patients, and ensuring the continuity of care—all of which are essential in managing seizure disorders [2].

The complexity of seizure disorders, which can include a broad spectrum ranging from focal seizures to generalized tonic-clonic seizures, necessitates an understanding of the neurological underpinnings that inform treatment modalities. Nurses must not only be well-versed in the pharmacotherapy used to manage these conditions—such as antiepileptic drugs—but also aware of the importance of adherence to treatment regimens, potential side effects, and the psychosocial impacts on patients and their families. This multifaceted care approach encapsulates the need for an interprofessional collaboration model, wherein nurses, physicians, neurologists, mental health professionals, and social workers work as a cohesive unit to address the biomedical and psychosocial aspects of epilepsy [3].

Moreover, nursing perspectives on managing seizure disorders extend beyond immediate clinical interventions. Education is a primary pillar in promoting the self-management of seizure disorders. Patients and caregivers must not only learn about the nature of the disorder, triggers, and how to manage seizures when they occur but also understand the importance of lifestyle modifications and routine follow-ups with healthcare providers. The nurse's role in health education encapsulates facilitating discussions surrounding potential lifestyle changes—such as sleep hygiene, stress management, and adherence to prescribed therapy—that are critical in reducing seizure frequency and improving overall quality of life [4].

In addition, the support of nurses in emergencies, such as during a seizure episode, cannot be understated. The training they receive equips them to provide immediate care and to alleviate the fears of patients and bystanders while promoting a safe environment. Education about seizure first aid is also vital, as many individuals may encounter situations where they need to assist someone experiencing a seizure. This proactive education helps to dismantle the stigma and fear often

associated with seizures, thereby enhancing community support for individuals living with seizure disorders [5].

Finally, it is essential to acknowledge the evolving role of nursing research and advocacy in the field of epilepsy care. There exists a critical need for ongoing research to elucidate the underlying mechanisms of seizure disorders, develop novel therapeutic approaches, evaluate the effectiveness of current interventions, and address disparities in access to care. Nurses, as part of this research landscape, contribute their insights and experiences, advocating for patients by participating in clinical trials, feedback mechanisms, and health service research that can inform policy and clinical practice [6].

### **Types of Seizures and Their Clinical Implications:**

Seizures are neurological events characterized by sudden, abnormal electrical activity in the brain, leading to various physical and psychological manifestations. Understanding the different types of seizures is crucial for accurate diagnosis, effective treatment, and appropriate management of individuals who experience them [7].

#### **Classification of Seizures**

Seizures are primarily classified into two broad categories: focal (or partial) seizures and generalized seizures. This classification is rooted in the origin of the seizure activity within the brain and has significant implications for clinical diagnosis and treatment [8].

##### **1. Focal Seizures**

Focal seizures originate in a specific area of the brain and can manifest with sensory, motor, or autonomic symptoms. They are further divided into two categories: focal seizures without impaired awareness and focal seizures with impaired awareness [9].

- **Focal Seizures Without Impaired Awareness:** These seizures often do not alter consciousness. Patients may exhibit motor symptoms like jerking of a limb or experience sensory changes such as tingling or visual disturbances. The clinical implication of these seizures is that patients may remain aware of their surroundings, which allows for a rapid description of the episode that can aid in diagnosis [10].
- **Focal Seizures With Impaired Awareness:** In these cases, consciousness is altered. Patients may appear confused, unresponsive, or engage in automatic movements, such as lip-smacking or hand rubbing. The clinical implication here is significant: since the patient may not recall the event, gathering eyewitness accounts becomes essential for accurate diagnosis and treatment planning [10].

##### **2. Generalized Seizures**

Generalized seizures result from widespread abnormal electrical activity across both hemispheres of the brain. These seizures can be further subdivided into several types, including:

- **Tonic-Clonic Seizures (Grand Mal):** Characterized by a combination of muscle rigidity (tonic phase) followed by rhythmic, jerking seizures (clonic phase). These seizures often involve loss of consciousness and can result in physical injuries, making it critical for immediate medical intervention post-episode [11].
- **Absence Seizures (Petit Mal):** Typically seen in children, absence seizures feature brief lapses in awareness and may include subtle body movements such as eye blinking. These seizures can be misinterpreted as daydreaming, emphasizing the need for awareness and

correct diagnosis to prevent misunderstandings regarding the individual's attentiveness [11].

- **Atonic Seizures:** These seizures result in a sudden loss of muscle tone, often causing the person to collapse. The clinical implications are serious, as they carry a high risk of falls and injuries, necessitating safety measures and possibly lifestyle adjustments to avoid hazardous situations [12].
- **Myoclonic Seizures:** Involves sudden, brief muscle jerks. Patients may experience one or multiple jerks, and even though consciousness is usually preserved, these seizures can be indicative of underlying conditions such as myoclonic epilepsies, which may require specific management approaches and long-term care [13].

### **Clinical Implications of Seizure Types**

The typology of seizures directly impacts clinical decision-making. For instance, understanding the nature of the seizure informs the choice of antiepileptic medications, both in terms of effectiveness and potential side effects [13].

1. **Medication Management:** Focal seizures may respond better to specific medications such as lamotrigine or carbamazepine, whereas generalized seizures often require drugs like valproate or levetiracetam. The selection process is guided not only by seizure type but also by underlying conditions, patient age, and other variables [14].
2. **Diagnosis of Underlying Conditions:** The classification and observed clinical manifestations can help healthcare providers identify the underlying cause of seizures (e.g., structural brain abnormalities, metabolic issues, genetic predispositions). Further diagnostic testing, including neuroimaging (like MRI or CT scans) or electroencephalograms (EEGs), can illuminate the etiology [14].
3. **Psychosocial Considerations:** Seizures can have profound psychosocial implications. Focal seizures with impaired awareness, for instance, may lead to misunderstandings in social scenarios or workplaces. Educating patients, their families, and even co-workers about their condition and how to respond during a seizure episode is vital to ensuring safety and reducing stigma [15].
4. **Emergency Management and Safety:** Certain seizure types, particularly tonic-clonic and atonic seizures, pose risks of injury during a seizure. Proper planning and education on how to respond to seizures, including where to safely position the person and when to seek medical assistance, are critical [15].
5. **Long-term Consequences and Care:** The long-term relationship with epilepsy can vary significantly based on seizure type. While some individuals may achieve remission with treatment, others may require ongoing management. Recognizing the type of seizures can guide the long-term care plan, including lifestyle adjustments and continuous monitoring for co-morbid conditions such as anxiety, depression, or cognitive impairment [15].

### **Pharmacological Management of Seizure Disorders:**

Seizure disorders, also known as epilepsy, encompass a wide range of medical conditions characterized by recurrent seizures, which are sudden and uncontrolled electrical disturbances in the brain. These disorders have significant implications for the affected individual's quality of life and often necessitate long-term pharmacological management. The treatment landscape of seizure disorders is marked by a diverse array of anticonvulsant medications, each with distinct mechanisms of action, side effects, and considerations for use [16].

### **Classification of Antiepileptic Drugs**

Antiepileptic drugs are generally classified based on their mechanism of action and their chemical structure. The two primary categories are:

1. **Traditional AEDs:** This group includes older agents such as phenytoin, carbamazepine, valproate, and phenobarbital. These medications have been used for decades and are well-researched, providing physicians with a solid understanding of their efficacy and safety profiles [17].
2. **Newer AEDs:** Introduced in recent years, these medications include lamotrigine, levetiracetam, topiramate, and zonisamide. They are associated with improved tolerability and fewer drug interactions, making them attractive options for many patients [17].

Understanding the classification helps clinicians choose the appropriate therapy based on the seizure type, patient age, comorbidities, and potential drug interactions. Additionally, it is crucial to note that some AEDs may work better for certain types of seizures than others, highlighting the importance of a personalized treatment approach [17].

### **Mechanisms of Action**

The pharmacological effects of AEDs stem from their interactions with various neurotransmitter systems and ion channels within the brain. The proposed mechanisms of action include:

1. **Sodium Channel Modulation:** Drugs like phenytoin and carbamazepine stabilize the neuronal membrane by reducing sodium influx during depolarization, which prevents the propagation of abnormal electrical activity [18].
2. **Calcium Channel Blockade:** Medications such as ethosuximide are effective in treating absence seizures by inhibiting T-type calcium channels, thereby reducing the influx of calcium ions that contributes to seizure activity [18].
3. **Increase in GABAergic Activity:** Benzodiazepines and valproate are known to enhance the effects of gamma-aminobutyric acid (GABA), the primary inhibitory neurotransmitter in the brain. This enhancement helps to dampen excessive neuronal excitability [18].
4. **Glutamate Receptor Antagonism:** Some newer AEDs, such as topiramate, inhibit the action of glutamate, an excitatory neurotransmitter, further contributing to the stabilization of neural activity [18].

Understanding these mechanisms enables healthcare professionals to tailor therapy to individual patients based on their seizure type and neurological condition [18].

### **Considerations for Pharmacological Management**

The pharmacological management of seizure disorders involves several important considerations, including dosage adjustments, monitoring for side effects, and potential drug interactions [19].

1. **Initial Therapy and Dosage Adjustments:** Initiating treatment usually starts with a single AED at a low dose, which is gradually increased based on the patient's response and tolerance. Therapeutic drug monitoring is vital for certain medications, such as phenytoin and valproate, to maintain plasma levels within a therapeutic range [19].
2. **Side Effects and Tolerability:** Each AED comes with potential side effects. For instance, valproate may cause weight gain and liver toxicity, while lamotrigine can lead to skin rashes. The side effects often dictate medication compliance; hence, patients should be educated about the importance of adherence to their medication regimen [19].
3. **Drug Interactions:** AEDs can interact with other medications, resulting in altered drug efficacy or increased toxicity. For example, enzyme-inducing AEDs like phenytoin may decrease the effectiveness of contraceptive agents. Therefore, a comprehensive review of a patient's medication history is essential to mitigate the risk of adverse interactions [19].

### **Personalized Medicine**

With the advances in pharmacogenomics, the field of personalized medicine is gaining traction in the management of seizure disorders. Genetic testing can identify variations in genes that affect drug metabolism and response. For instance, the UGT1A1 gene polymorphism impacts the metabolism of lamotrigine, and individuals with specific alleles may require dose adjustments to avoid toxicity. Moreover, the development of biomarkers to predict treatment response can lead to more tailored therapeutic strategies, improving efficacy and minimizing side effects [20].

### **Emerging Therapeutic Strategies**

While pharmacological treatments remain the cornerstone of management for seizure disorders, innovative strategies are emerging on the horizon. These include:

1. **Non-pharmacological Interventions:** Techniques such as ketogenic diets, which shift the body's metabolism to utilize fats instead of carbohydrates, have shown efficacy in managing refractory epilepsy, particularly in children [20].
2. **Neurostimulation Devices:** Vagus nerve stimulation (VNS) and responsive neurostimulation (RNS) are surgical options designed to modulate brain activity and reduce seizure frequency. These approaches promote the idea that not all patients will respond to pharmacotherapy, necessitating alternative approaches [20].
3. **Research into Novel Agents:** Ongoing research is dedicated to discovering new AEDs with less severe side effects and improved efficacy. For example, cannabidiol, a non-psychoactive component of cannabis, has been approved for treating certain refractory seizure syndromes, showcasing the potential of alternative treatments [21].

### **Nursing Assessment and Monitoring Strategies:**

Seizure disorders are among the most common neurological conditions affecting individuals of all ages, from infants to the elderly. These disorders are characterized by abnormal electrical activity in the brain, leading to diverse manifestations, including convulsions, loss of consciousness, and altered sensations and behaviors. The ongoing management of seizure disorders requires a comprehensive nursing assessment and vigilant monitoring [22].

Seizure disorders, commonly referred to as epilepsy, can arise from various etiologies, including genetic predispositions, structural brain abnormalities, metabolic imbalances, and traumatic injuries. Clinically, seizures can be classified into several types, including focal seizures (which originate in one hemisphere of the brain) and generalized seizures (which affect both hemispheres). The variability in seizure types necessitates tailored nursing assessments and monitoring strategies to effectively address the unique needs of each patient [23].

### **Nursing Assessment**

A thorough nursing assessment for patients with seizure disorders should encompass several key components. The goals of this assessment are to establish a baseline for the patient's condition, identify triggers and patterns of seizure activity, and provide essential information to guide treatment decisions [24].

1. **Health History:** Collecting a comprehensive health history involves documenting the patient's previous seizure episodes, including the frequency, duration, and character of the seizures, as well as any accompanying symptoms. It is also vital to inquire about family history of epilepsy, concurrent medical conditions, medications (including over-the-counter and herbal supplements), and substance use [24].

2. **Physical Examination:** A thorough physical examination should assess neurological function and identify any potential complications related to seizure activity, such as injuries from past seizures. During the examination, nurses should evaluate cognitive function, motor skills, and the patient's awareness and orientation level [24].
3. **Seizure Diary:** Encouraging patients and their families to maintain a seizure diary can be instrumental in tracking seizure frequency, duration, and any identifiable triggers such as stress, sleep deprivation, medication adherence, and environmental factors. This diary can also provide valuable insights into postictal states (the period following a seizure), which can help in the management and adjustment of treatment plans.
4. **Diagnostic Testing:** Nurses often facilitate diagnostic testing, including electroencephalography (EEG) and neuroimaging studies (CT or MRI scans). EEG testing helps to visualize electrical activity in the brain and can identify specific patterns of abnormal activity indicative of seizures. Neuroimaging can help detect structural abnormalities that may be contributing to seizure disorders [24].

### Monitoring Strategies

Effective monitoring of patients with seizure disorders is crucial for ensuring patient safety, preventing complications, and optimizing treatment outcomes. The following strategies are essential for nurses in this regard:

1. **Continuous Observation:** For patients hospitalized or at risk of experiencing seizures, continuous observation is necessary to monitor for seizure activity and its potential complications. In some cases, using video-EEG monitoring systems allows for real-time observation of seizures, providing enhanced data for diagnosis and treatment planning [25].
2. **Vital Signs Monitoring:** Nurses should routinely monitor vital signs (including heart rate, blood pressure, and oxygen saturation) before, during, and after seizure activity. Changes in vital signs can indicate autonomic dysregulation, and careful monitoring can help identify potential complications, such as status epilepticus, a prolonged seizure that requires immediate intervention [25].
3. **Postictal Assessment:** After a seizure, patients often experience a transitional phase known as the postictal state. Nurses should perform a thorough postictal assessment to evaluate the patient's level of consciousness, any neurological deficits, and vital signs. Understanding a patient's postictal symptoms is essential for differentiating between various seizure types and planning subsequent care [25].
4. **Medication Management:** Since pharmacotherapy is a cornerstone of seizure management, monitoring the patient's medication adherence, effectiveness, side effects, and serum drug levels (where applicable) is imperative. Educating patients about the importance of consistent medication use, potential side effects, and interactions with other drugs is also vital for optimal management [25].
5. **Patient and Family Education:** Educating patients and their families about seizure disorders and how to respond during a seizure episode is crucial. Nurses should provide

clear instructions on safety measures, emergency contacts, and how to use rescue medications if applicable. Empowering patients and families with knowledge enhances their ability to manage seizures at home and reduces anxiety about the condition [26].

6. **Emotional and Psychological Monitoring:** Seizure disorders can impact the emotional and psychological well-being of patients, often resulting in anxiety, depression, and social isolation. Regular assessment of the patient's emotional health is necessary. Nurses should initiate conversations about mental health and refer patients to counselors or support groups when needed [26].

### **Patient Education and Counseling Techniques:**

Seizure disorders, commonly referred to as epilepsy, encompass a range of neurological conditions characterized by recurrent seizures due to abnormal electrical activity in the brain. The prevalence of seizure disorders in the general population necessitates effective patient education and counseling. Understanding the complexities surrounding these conditions can greatly empower patients and their families, promoting adherence to treatment plans and improving quality of life [27].

Before delving into educational techniques, it is crucial to establish a foundational understanding of what seizure disorders entail. Seizures manifest in numerous forms, categorized broadly into focal seizures (originating from a specific area of the brain) and generalized seizures (affecting the entire brain). Symptoms can vary widely from uncontrolled jerking movements, to periods of unconsciousness, or subtle changes in behavior. Recognizing these symptoms is fundamental not only for diagnosis but also for patient and caregiver education [28].

A successful educational and counseling approach begins with the establishment of a therapeutic relationship between healthcare providers and patients. Trust and rapport form the bedrock upon which effective communication can occur. Clinicians should actively listen to patients' concerns, validate their feelings, and ensure that they feel understood. This can be achieved through open-ended questions that promote dialogue, allowing patients to express their experiences and worries about their conditions. A collaborative approach reassures patients that they are active participants in their own care [29].

Each patient's experience with seizures is unique; hence, educational techniques should be tailored to meet individual needs. Factors such as age, cognitive ability, cultural background, and personal experiences with seizures play a significant role in how information is received. For example, educating young children about seizures may involve using simplified language, visual aids, or interactive methods, while older adults may benefit from more comprehensive discussions that address lifestyle changes, medication management, and the importance of routine follow-up care [30].

The use of visual aids such as diagrams, charts, and videos can significantly enhance understanding for patients and their families. Visual representations of the brain, the types of seizures, and mechanisms of action of anti-seizure medications can make complex information more digestible. Moreover, instructional videos demonstrating seizure first-aid techniques are invaluable for caregivers, allowing them to be prepared in emergencies. Modern technological solutions, such as mobile applications that track seizures, remind patients of medication schedules, and provide educational resources, can further support self-management and patient autonomy [31].

Giving patients educational pamphlets, booklets, or access to reliable online resources can reinforce their understanding. These materials should cover essential topics such as seizure recognition, management strategies, lifestyle modifications, medication adherence, and potential

social implications of living with a seizure disorder. Simple, clear language alongside bullet points and headings can make these resources more accessible. However, it's essential that patients are guided on how to use these materials effectively, as overwhelming information can lead to confusion [32].

Education should not be limited to the patients alone; it must also include families and caregivers who play a crucial role in the patient's support system. Inviting family members to attend educational sessions fosters a collaborative approach and ensures that caregivers understand how to respond during a seizure, help manage medications, and provide ongoing support. Family dynamics, beliefs, and experiences shape a patient's journey, making it essential for the care team to engage with these networks. Additionally, support groups and community resources offer invaluable opportunities for patients and families to connect with others facing similar challenges, thereby reducing feelings of isolation [33].

Educating patients about the importance of medication adherence is paramount in managing seizure disorders. Many patients may struggle with side effects, the complexity of medication regimens, or forgetfulness. Healthcare professionals can utilize various strategies, such as teaching the patient about their specific medications—how they work, the reasons for taking them, and understanding side effects—to engender a sense of ownership. Employing medication management tools, such as pill organizers, reminders, and apps, can help patients maintain adherence. Additionally, regular follow-ups to review treatment effectiveness and make necessary adjustments can reinforce the importance of staying consistent with medications [34].

Living with a seizure disorder can be emotionally taxing due to the uncertainty and societal stigma associated with seizures. It is crucial to incorporate psychological counseling and mental health screenings into routine care. Patients may benefit from techniques such as cognitive-behavioral therapy (CBT) to help them cope with anxiety and depression often linked to their condition. Encouraging mindfulness practices and relaxation techniques can also reduce stress, which is known to be a seizure trigger for many individuals. By addressing these intertwined psychological factors, healthcare providers can significantly improve overall patient well-being.

Another critical facet of managing seizure disorders involves lifestyle and dietary modifications. Educating patients on the impact of sleep, stress, alcohol, and recreational drugs on seizure control is essential. For some individuals, specific diets—such as the ketogenic diet—may be effective in reducing seizure frequency. Providing individualized dietary plans and resources can bolster the patient's ability to make informed choices regarding their health. Exercise regimens should also be discussed, balancing physical activity with safety considerations specific to their condition [35].

### **Safety Considerations and Emergency Protocols:**

Seizure disorders, also known as epilepsy, encompass a range of neurological conditions characterized by recurrent, unprovoked seizures. These disturbances in brain function can vary in intensity and duration, presenting unique safety challenges for individuals and caregivers alike. Given the unpredictable nature of seizures, understanding safety considerations and emergency protocols is crucial for minimizing the risk of injury and managing crises effectively [36].

Seizures occur due to abnormal electrical activity in the brain, which can stem from various causes including genetic factors, head injuries, infections, and brain tumors, among others. The manifestations of seizures are diverse; some individuals may experience generalized tonic-clonic seizures, which involve loss of consciousness and rhythmic muscle contractions, while others may have focal seizures, characterized by localized symptoms such as twitching or sensory alterations. Each type brings distinct safety concerns that warrant specific attention [37].

## Safety Considerations for Individuals with Seizure Disorders

- 1. Mobility and Environment:**  
Individuals with seizure disorders should assess their living environments for potential hazards. This includes removing sharp objects, installing safety gates, and ensuring that medications are stored securely. Individuals prone to seizures, particularly if they exhibit loss of consciousness, should avoid situations that could increase their risk of injury, such as climbing ladders, swimming unsupervised, or riding bicycles without helmets [38].
- 2. Driving Regulations:**  
In many regions, guidelines restrict individuals with seizure disorders from driving until they have been seizure-free for a legal period, often ranging from six months to two years. Strict adherence to these regulations is fundamental to ensure not only the driver's safety but also that of others on the road [38].
- 3. Supervision:**  
Individuals who experience seizures may benefit from having a caregiver, particularly in situations where seizures could lead to danger. This is especially important for children and individuals with significant disabilities. Caregivers should be trained to recognize seizure triggers and understand how to provide immediate assistance if a seizure occurs [38].
- 4. Adequate Medical Management:**  
Regular consultations with healthcare providers for medication management are vital. Medications can help control seizures, but they must be taken consistently and adjusted as necessary. Patients must also be educated about potential side effects and the importance of medication adherence to maintain their health and prevent breakthrough seizures [39].
- 5. Creating a Seizure Response Plan:**  
Individuals with seizure disorders, along with their families and caregivers, should establish a seizure response plan. This plan should detail step by step procedures for how to respond during a seizure, including the positioning of the individual to prevent aspiration, how to time the duration of the seizure, and when to seek emergency help [39].

## Emergency Protocols for Seizure Disorders

Despite preventive measures, emergencies can still arise. Knowing how to react appropriately is crucial for ensuring the safety of individuals experiencing a seizure [40].

- 1. Immediate Response:**  
When a seizure occurs, the first step is to remain calm. Shouting or panicking can create undue stress for both the person experiencing the seizure and those nearby. Caregivers should ensure that the environment is safe by moving away potential hazards, such as furniture or sharp objects [40].
- 2. Timing the Seizure:**  
It is essential to note the duration of the seizure. Most seizures last between a few seconds to a couple of minutes. If a seizure extends beyond five minutes or if a second seizure follows immediately, emergency medical assistance should be sought [40].
- 3. Positioning:**  
During a seizure, individuals should be gently laid on their side to prevent choking and facilitate breathing. Placing something soft under their head can help to reduce the risk of injury. It is crucial to avoid restraining the individual or placing anything in their mouth, as this can cause harm [41].
- 4. Postictal Care:**  
After the seizure, the person may experience confusion, drowsiness, or memory loss, a state known as the postictal phase. It is advisable to offer reassurance and stay with the

individual until they regain full awareness. Hydration and rest are important during this recovery period [42].

5. **Educating**

**Bystanders:**

Bystanders play an important role in seizure management. Educational efforts should include teaching those around the individual how to respond to seizures effectively. Good practices include clearing the area, timing the seizure, and knowing when to call for emergency services [42].

6. **Emergency**

**Contacts:**

Keeping a list of emergency contacts readily available is prudent. This list should include medical professionals, family members, and any healthcare proxies who can provide immediate assistance or vital information to responders [43].

### **Interdisciplinary Collaboration in Seizure Management:**

Seizure disorders, primarily characterized by abnormal electrical activity in the brain, pose significant challenges for both patients and healthcare providers. The complexity of epilepsy, the most prevalent of these disorders, necessitates an approach that transcends individual medical specialties. Interdisciplinary collaboration is not just beneficial but essential for the effective management of seizure disorders [44].

Seizures can manifest in various forms, impacting individuals differently based on the type and frequency of seizures experienced. Epilepsy affects approximately 1% of the global population, with around 50 million people worldwide living with this condition, making it one of the most common neurological disorders. Seizure management typically involves medical, psychological, educational, and social considerations. Therefore, effective management requires contributions from multiple healthcare disciplines including neurology, psychiatry, nursing, social work, and education [45].

### **Components of Interdisciplinary Collaboration**

At its core, interdisciplinary collaboration involves the integration of knowledge, skills, and perspectives from various healthcare professionals to provide comprehensive care. In the context of seizure management, key components of this collaboration include:

1. **Team Composition:** A well-rounded team might include neurologists, epileptologists, neuropsychologists, psychiatric clinicians, nurse practitioners, social workers, and occupational or physical therapists [46].
2. **Shared Goals:** Every member of the team must understand the primary objective—achieving the best outcomes for the patient. This may include controlling seizure activity, minimizing side effects of treatment, improving quality of life, and providing education about the disorder [46].
3. **Effective Communication:** Regular and clear communication among team members ensures that all aspects of a patient's condition are addressed. This communication can take place through team meetings, electronic health records, and patient conferences, where multidisciplinary perspectives enhance decision-making [46].
4. **Patient-Centric Approach:** Central to interdisciplinary collaboration is a focus on the patient and their family. Engaging patients in discussions about their treatment options ensures that care is tailored to their individual needs, preferences, and lifestyles [46].

### **Benefits of Interdisciplinary Collaboration**

The integration of diverse skill sets and knowledge areas results in numerous benefits for seizure management:

1. **Comprehensive Assessment:** Patients with seizure disorders often experience comorbidities, including anxiety, depression, and cognitive impairment. An interdisciplinary approach allows for thorough assessments that take into account all factors affecting the patient's health, leading to more accurate diagnoses and effective treatments [47].
2. **Holistic Treatment Plans:** Collaborative teams can develop a multi-faceted treatment plan that addresses not only neurological aspects but also psychosocial dimensions of the disorder. For example, a neurologist may prescribe antiepileptic drugs while a psychologist addresses mental health needs through therapy [48].
3. **Enhanced Support and Resources:** Collaborating with social workers or case managers helps families navigate complex healthcare systems and connects them with community resources, educational opportunities, and support networks [48].
4. **Informed Decision-Making:** Leveraging the expertise of various specialists allows for evidence-based recommendations tailored to the patient's unique situation. This might include considering alternative treatments such as dietary modifications or advanced therapies while weighing potential risks and benefits [48].
5. **Improved Quality of Life:** When diverse healthcare professionals collaborate effectively, patients are more likely to experience a reduction in the frequency and severity of seizures, enhancing their overall quality of life. In addition, supportive therapeutic services, such as occupational therapy, can address daily functioning challenges related to seizure disorders [49].

### **Challenges to Interdisciplinary Collaboration**

While the benefits of interdisciplinary collaboration in seizure management are significant, several challenges persist:

1. **Professional Silos:** Healthcare professionals often work within established disciplinary boundaries, leading to fragmentation in care. Breaking down these silos requires a cultural shift in health services towards embracing interdisciplinary approaches [50].
2. **Communication Barriers:** Differences in professional language, priorities, and workflows can impede effective communication. Team members must develop mutual understanding and establish clear communication pathways to facilitate collaboration [51].
3. **Resource Allocation:** Integrating multiple services may lead to resource strain, particularly in healthcare systems with limited funding. Patients may struggle to access the full range of services available due to financial constraints or administrative complexities [52].
4. **Time Constraints:** Coordinating schedules for team meetings and patient consultations can be difficult, particularly in busy clinical settings. Effective collaboration demands dedicated time and commitment from all team members [52].
5. **Training and Education:** Interdisciplinary collaboration is not typically emphasized in healthcare education programs, resulting in professionals who may feel unprepared to work in a team-centric environment. Programs focused on teaching collaborative skills are necessary to bridge this gap [52].

### **Moving Forward: Strategies for Improvement**

For the effective implementation of interdisciplinary collaboration in seizure management, several strategies can be pursued:

1. **Interprofessional Education (IPE):** Incorporating IPE into medical and allied health programs fosters teamwork skills among future healthcare providers. This education can enhance understanding of the roles and contributions of various professionals.

2. **Development of Collaborative Protocols:** Establishing standardized interdisciplinary protocols can streamline care, ensuring that all team members are aligned in their approach to managing seizure disorders [53].
3. **Utilization of Technology:** Advanced electronic health records can facilitate communication among team members, allowing real-time updates on a patient's status and shared access to treatment plans [54].
4. **Research and Evidence-Based Practice:** Engaging in collaborative research initiatives exploring best practices for interdisciplinary seizure management can contribute to the development of effective treatment models [55].
5. **Policy and Advocacy:** Stakeholders can advocate for policies supporting collaborative healthcare models, emphasizing funding for interdisciplinary initiatives and promoting public awareness about the importance of comprehensive care for patients with seizure disorders [55].

### **Impact of Seizure Disorders on Mental Health and Quality of Life:**

Seizure disorders, commonly referred to as epilepsy, encompass a range of neurological conditions that are characterized by recurrent seizures. These episodic disruptions of electrical activity in the brain can significantly affect the lives of patients, not only through the physical manifestations of the seizures themselves but also in their profound impact on mental health and overall quality of life. Understanding the relationship between seizure disorders and these broader life domains is crucial for healthcare providers, patients, and families, as it can guide effective management strategies and improve outcomes for those affected [56].

Seizure disorders manifest in various forms, with epilepsy being the most recognized. The types of seizures can vary widely, encompassing generalized seizures, which affect both hemispheres of the brain, and partial seizures, which originate in a specific area of the brain. The unpredictability of seizures can lead to alarming consequences, including physical injuries, cognitive impairments, and social stigmatization. While medical treatments, including anticonvulsant medications and surgical interventions, are widely used to manage seizure disorders, the indirect consequences on the mental health of patients cannot be overlooked [57].

Research has shown a strong correlation between seizure disorders and mental health issues, particularly anxiety and depression. The fear of experiencing a seizure, especially in public or during critical moments, can lead to increased anxiety, which may evolve into a chronic anxiety disorder. Patients may find themselves in a constant state of hypervigilance, monitoring their bodies for any signs that a seizure might be imminent. This ongoing anxiety can severely limit a person's willingness to engage in everyday activities, such as driving, socializing, or even going to work, consequently leading to increased feelings of isolation [58].

Depression is another prevalent mental health issue among individuals with seizure disorders. The chronic nature of the condition, coupled with the stress of managing seizures and potential side effects of medication, can create a sense of hopelessness. Additionally, the societal stigma attached to epilepsy often exacerbates feelings of worthlessness or shame. Research indicates that individuals with epilepsy are more likely to experience depressive episodes compared to the general population, with studies suggesting that the prevalence of major depressive disorder in individuals with epilepsy may be as high as 30-50% [59].

Quality of life (QoL) for individuals with seizure disorders is multifaceted and influenced by both the direct effects of seizures and the secondary outcomes associated with mental health. Several

factors contribute to the perception of QoL, including physical health, mental well-being, social relationships, and the ability to participate in daily activities. Patients experiencing frequent seizures often report lower QoL due to the limitations imposed on their lifestyles. For instance, many individuals with seizure disorders struggle to maintain employment, pursue education, or engage in leisure activities due to their condition, leading to socioeconomic disadvantages [60].

The stigma surrounding seizure disorders also plays a significant role in QoL. Individuals may face discrimination or social isolation, which can lead to a decrease in meaningful relationships. This loss of social support can further exacerbate feelings of depression and anxiety, creating a vicious cycle that can be difficult to break. Moreover, the fear of future seizures can lead to a decrease in self-efficacy, where individuals may feel powerless to control their condition and its consequences [61].

It is important to note that while seizures may occur unpredictably and are often uncontrollable, the management of a seizure disorder can significantly enhance an individual's quality of life. Utilizing a comprehensive treatment approach that includes medication, lifestyle modifications, and psychological support can ameliorate both the mental health and QoL challenges faced by patients. Personalized care that addresses the individual's specific experiences with seizure disorders can foster a sense of agency and hope [62].

The role of support systems cannot be understated when discussing the impact of seizure disorders on mental health and QoL. Family members, friends, and support groups can provide vital emotional support, helping individuals navigate the complexities of their condition. Educational resources that inform loved ones about seizure disorders can facilitate a better understanding and foster empathy, minimizing stigma and isolation [63].

Coping strategies are also essential for individuals managing seizure disorders. Mindfulness practices, such as yoga and meditation, have been shown to reduce anxiety levels and support emotional regulation. Cognitive-behavioral therapy (CBT) can be particularly beneficial for addressing distorted thought patterns associated with seizures and their repercussions. Therapeutic interventions can empower patients to develop resilience and adaptive coping techniques, augmenting their ability to manage both seizures and the accompanying mental health challenges [64].

### **Conclusion:**

In conclusion, managing seizure disorders from a nursing perspective encompasses a multifaceted approach that prioritizes patient safety, effective pharmacological intervention, and comprehensive education. Nurses play a pivotal role in the assessment, monitoring, and coordination of care for individuals living with these complex conditions. By understanding the various types of seizures and their specific implications, nurses can tailor care plans that address both medical and psychosocial needs, ultimately enhancing the patient's quality of life.

ABDULLAH AHMED MOHAMMED ALKHALDI <sup>1</sup>, MAITHA KHALIL LAFI ALKHAIBARI <sup>2</sup>, Sultan Salem Sakhani Alruwaili <sup>3</sup>, Abdullah Fihat Ghadeer Alanazi <sup>4</sup>, Sultan Marji Amash Alanazi <sup>5</sup>, Amani Saud Al-Daghmani <sup>6</sup>, Tahani Saud Al-Daghmani <sup>7</sup>, Jamal Salamah Hamoud Albalawi <sup>8</sup>, Ahmed Nasser Ahmed Alasiri <sup>9</sup>, Abdulkarim Mulhi Mulhid Alanazi <sup>10</sup>, Omar Nadaa Huosein Alanazi <sup>11</sup>, NAIF KHALAF AWAD ALSWINA <sup>12</sup>, Dhafer Mesfer Ali Al Awn <sup>13</sup>

Moreover, the importance of patient and family education cannot be overstated, as it empowers them to recognize triggers, adhere to treatment regimens, and effectively respond to seizure episodes. Through collaborative efforts with interdisciplinary teams, nurses can ensure holistic care that considers not only the physiological aspects of seizure management but also the emotional and psychological support required for patients. As the landscape of healthcare evolves, continued education, research, and advocacy in the management of seizure disorders remain essential, reinforcing the critical role of nursing in improving the lives of those affected.

## Introduction:

1. Camfield P., Camfield C. Incidence, prevalence and aetiology of seizures and epilepsy in children. *Epileptic Disord.* 2015;17(2):117–123. doi: 10.1684/epd.2015.0736.
2. Idro R., Gwer S., Kahindi M. The incidence, aetiology and outcome of acute seizures in children admitted to a rural Kenyan district hospital. *BMC Pediatr.* 2008;8:5. doi: 10.1186/1471-2431-8-5.
3. Govender R., Eley B., Walker K. Neurologic and neurobehavioral sequelae in children with human immunodeficiency virus (HIV-1) infection. *J Child Neurol.* 2011;26(11):1355–1364. doi: 10.1177/0883073811405203.
4. Verity C.M., Greenwood R., Golding J. Long-term intellectual and behavioural outcomes of children with febrile convulsions. *N Engl J Med.* 1998;338(24):1723–1728. doi: 10.1056/NEJM199806113382403.
5. Wagner R.G., Ibinda F., Tollman S. Differing methods and definitions influence DALY estimates: using population-based data to calculate the burden of convulsive epilepsy in rural South Africa. *PLoS One.* 2015;10(12) doi: 10.1371/journal.pone.0145300.
6. Baskind R., Birbeck G.L. Epilepsy-associated stigma in sub-Saharan Africa: the social landscape of a disease. *Epilepsy Behav.* 2005;7(1):68–73. doi: 10.1016/j.yebeh.2005.04.009.
7. Chomba E., Taylor T.E., Hauser W. Seizure recurrence in rural Zambian children admitted with febrile seizures. *Open J Trop Med.* 2008:1101–1107.
8. Edwards T., Scott A.G., Munyoki G. Active convulsive epilepsy in a rural district of Kenya: a study of prevalence and possible risk factors. *Lancet Neurol.* 2008;7(1):50–56. doi: 10.1016/S1474-4422(07)70292-2.
9. Glauser T., Shinnar S., Gloss D. Evidence-based guideline: treatment of convulsive status epilepticus in children and adults: report of the Guideline Committee of the American Epilepsy Society. *Epilepsy Curr.* 2016;16(1):48–61. doi: 10.5698/1535-7597-16.1.48.
10. Mung'ala-Odera V., White S., Meehan R. Prevalence, incidence and risk factors of epilepsy in older children in rural Kenya. *Seizure.* 2008;17(5):396–404. doi: 10.1016/j.seizure.2007.11.028.
11. Atadzhanov M., Haworth A., Chomba E.N. Epilepsy-associated stigma in Zambia: what factors predict greater felt stigma in a highly stigmatized population? *Epilepsy Behav.* 2010;19(3):414–418. doi: 10.1016/j.yebeh.2010.08.017.
12. Wagner R.G., Bottomley C., Ngugi A.K. Incidence, remission and mortality of convulsive epilepsy in rural northeast South Africa. *PLoS One.* 2015;10(6). doi: 10.1371/journal.pone.0129097.
13. Wilmshurst J.M., Burgess J., Hartley P. Specific neurologic complications of human immunodeficiency virus type 1 (HIV-1) infection in children. *J Child Neurol.* 2006;21(9):788–794. doi: 10.1177/08830738060210091901.
14. WHO, Emergency Triage, Assessment and Treatment. 2005: Geneva. p. 83.
15. Wilmshurst J.M., Donald K.A., Eley B. Update on the key developments of the neurologic complications in children infected with HIV. *Curr Opin HIV AIDS.* 2014;9(6):533–538. doi: 10.1097/COH.0000000000000101.
16. Ba-Diop A., Marin B., Druet-Cabanac M. Epidemiology, causes, and treatment of epilepsy in sub-Saharan Africa. *Lancet Neurol.* 2014;13(10):1029–1044. doi: 10.1016/S1474-4422(14)70114-0.
17. Shohet C., Yelloly J., Bingham P. The association between the quality of epilepsy management in primary care, general practice population deprivation status and epilepsy-

- related emergency hospitalisations. *Seizure*. 2007;16(4):351–355. doi: 10.1016/j.seizure.2007.02.005.
18. Birbeck G.L., Molyneux M.E., Kaplan P.W. Blantyre Malaria Project Epilepsy Study (BMPES) of neurological outcomes in retinopathy-positive paediatric cerebral malaria survivors: a prospective cohort study. *Lancet Neurol*. 2010;9(12):1173–1181. doi: 10.1016/S1474-4422(10)70270-2.
  19. Mwipopo E.E., Akhtar S., Fan P. Profile and clinical characterization of seizures in hospitalized children. *Pan Afr Med J*. 2016;24:313. doi: 10.11604/pamj.2016.24.313.9275.
  20. WHO. MHGAP intervention guideline 2.0 for mental, neurological and substance use disorders in non-specialized settings. Geneva: WHO; 2016. p. 174.
  21. Glauser T., Shinnar S., Gloss D., et al. Evidence-based guideline: treatment of convulsive status epilepticus in children and adults: report of the guideline committee of the American Epilepsy Society. *Epilepsy Curr*. 2016;16:48–61. doi: 10.5698/1535-7597-16.1.48.
  22. Task Force for the Diagnosis and Management of Syncope, European Society of Cardiology (ESC), European Heart Rhythm Association (EHRA), Heart Failure Association (HFA) et al. Guidelines for the diagnosis and management of syncope (version 2009). *Eur Heart J*. 2009;30:2631–71. doi: 10.1093/eurheartj/ehp298.
  23. Reuber M., Chen M., Jamnadas-Khoda J., et al. Value of patient-reported symptoms in the diagnosis of transient loss of consciousness. *Neurology*. 2016;87:625–33. doi: 10.1212/WNL.0000000000002948.
  24. Bleck T., Cock H., Chamberlain J., et al. The Established Status Epilepticus Trial 2013. *Epilepsia*. 2013;54:89–92. doi: 10.1111/epi.12288.
  25. Ferguson M., Bianchi M.T., Sutter R., et al. Calculating the risk benefit equation for aggressive treatment of non-convulsive status epilepticus. *Neurocrit Care*. 2013;18:216–27. doi: 10.1007/s12028-012-9785-y.
  26. Yasiry Z., Shorvon S.D. The relative effectiveness of five antiepileptic drugs in treatment of benzodiazepine-resistant convulsive status epilepticus: a meta-analysis of published studies. *Seizure*. 2014;23:167–74. doi: 10.1016/j.seizure.2013.12.007.
  27. Sheldon R., Rose S., Ritchie D., et al. Historical criteria that distinguish syncope from seizures. *J Am Coll Cardiol*. 2002;40:142–8. doi: 10.1016/s0735-1097(02)01940-x.
  28. Chen D.K., So Y.T., Fisher R.S. Use of serum prolactin in diagnosing epileptic seizures: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology*. 2005;65:668–75. doi: 10.1212/01.wnl.0000178391.96957.d0.
  29. Dunn M.J.G., Breen D.P., Davenport R.J., Gray A.J. Early management of adults with an uncomplicated first generalised seizure. *Emerg Med J*. 2005;22:237–42. doi: 10.1136/emj.2004.015651.

30. van Dijk N., Boer K.R., Colman N., et al. High diagnostic yield and accuracy of history, physical examination, and ECG in patients with transient loss of consciousness in FAST: the Fainting Assessment study. *J Cardiovasc Electrophysiol.* 2008;19:48–55. doi: 10.1111/j.1540-8167.2007.00984.x.
31. Matz O., Zdebek C., Zechbauer S., et al. Lactate as a diagnostic marker in transient loss of consciousness. *Seizure.* 2016;40:71–5. doi: 10.1016/j.seizure.2016.06.014.
32. Trinka E., Cock H., Hesdorffer D., et al. A definition and classification of status epilepticus – Report of the ILAE task force on classification of status epilepticus. *Epilepsia.* 2015;56:1515–23. doi: 10.1111/epi.13121.
33. Lempert T., Bauer M., Schmidt D. Syncope: a videometric analysis of 56 episodes of transient cerebral hypoxia. *Ann Neurol.* 1994;36:233–7. doi: 10.1002/ana.410360217.
34. Marson A., Jacoby A., Johnson A., et al. Immediate versus deferred antiepileptic drug treatment for early epilepsy and single seizures: a randomised controlled trial. *Lancet Lond Engl.* 2005;365:2007–13. doi: 10.1016/S0140-6736(05)66694-9.
35. Soteriades E.S., Evans J.C., Larson M.G., et al. Incidence and prognosis of syncope. *N Engl J Med.* 2002;347:878–85. doi: 10.1056/NEJMoa012407.
36. National Institute for Health and Clinical Excellence. *Epilepsies: diagnosis and management [CG137].* London: NICE; 2012.
37. Joint Epilepsy Council. *Prevalence, incidence and other statistics.* JEC. 2011.
38. Fisher R.S., Cross J.H., French J.A., et al. Operational classification of seizure types by the International League Against Epilepsy: position paper of the ILAE Commission for Classification and Terminology. *Epilepsia.* 2017;58:522–30. doi: 10.1111/epi.13670.
39. Briggs R., Coughlan T., Doherty J., et al. Investigation and diagnostic formulation in patients admitted with transient loss of consciousness. *Ir Med J.* 2017;110:563.
40. Beghi E., Carpio A., Forsgren L., et al. Recommendation for a definition of acute symptomatic seizure. *Epilepsia.* 2010;51:671–5. doi: 10.1111/j.1528-1167.2009.02285.x.
41. Westover M.B., Shafi M.M., Bianchi M.T., Moura L.M.V.R., O'Rourke D., Rosenthal E.S., Chu C.J., Donovan S., Hoch D.B., Kilbride R.D., et al. The probability of seizures during EEG monitoring in critically ill adults. *Clin. Neurophysiol.* 2015;126:463–471. doi: 10.1016/j.clinph.2014.05.037.
42. Shorvon S., Ferlisi M. The treatment of super-refractory status epilepticus: A critical review of available therapies and a clinical treatment protocol. *Brain J. Neurol.* 2011;134:2802–2818. doi: 10.1093/brain/awr215.
43. Riazi K., Galic M.A., Pittman Q.J. Contributions of peripheral inflammation to seizure susceptibility: Cytokines and brain excitability. *Epilepsy Res.* 2010;89:34–42. doi: 10.1016/j.eplepsyres.2009.09.004.

44. Crepeau A.Z., Rabinstein A.A., Fugate J.E., Mandrekar J., Wijdicks E.F., White R.D., Britton J.W. Continuous EEG in therapeutic hypothermia after cardiac arrest: Prognostic and clinical value. *Neurology*. 2013;80:339–344. doi: 10.1212/WNL.0b013e31827f089d.
45. Pruitt P., Naidech A., Ornam J.V., Borczuk P. Seizure frequency in patients with isolated subdural hematoma and preserved consciousness. *Brain Inj.* 2019;0:1–5. doi: 10.1080/02699052.2019.1606446.
46. Dham B.S., Hunter K., Rincon F. The Epidemiology of Status Epilepticus in the United States. *Neurocrit. Care*. 2014;20:476–483. doi: 10.1007/s12028-013-9935-x.
47. Delanty N., Vaughan C.J., French J.A. Medical causes of seizures. *The Lancet*. 1998;352:383–390. doi: 10.1016/S0140-6736(98)02158-8.
48. Liu X., Yan B., Wang R., Li C., Chen C., Zhou D., Hong Z. Seizure outcomes in patients with anti-NMDAR encephalitis: A follow-up study. *Epilepsia*. 2017;58:2104–2111. doi: 10.1111/epi.13929.
49. McNett M., Moran C., Johnson H. Evidence-Based Review of Clinical Trials in Neurocritical Care. *AACN Adv. Crit. Care*. 2018;29:195–203. doi: 10.4037/aacnacc2018200.
50. Nielsen N., Sunde K., Hovdenes J., Riker R.R., Rubertsson S., Stammet P., Nilsson F., Friberg H. Adverse events and their relation to mortality in out-of-hospital cardiac arrest patients treated with therapeutic hypothermia. *Crit. Care Med*. 2011;39:57–64. doi: 10.1097/CCM.0b013e3181fa4301.
51. Vezzani A., Balosso S., Ravizza T. Chapter 10—Inflammation and epilepsy. In: Stefan H., Theodore W.H., editors. *Handbook of Clinical Neurology—Epilepsy*. Volume 107. Elsevier; Maryland Heights, MO, USA: 2012. pp. 163–175.
52. Shin J.-W., Koo Y.S., Kim Y.-S., Kim D.W., Kim K.K., Lee S.-Y., Kim H.K., Moon H.-J., Lim J.-A., Byun J.-I., et al. Clinical characterization of unknown/cryptogenic status epilepticus suspected as encephalitis: A multicenter cohort study. *J. Neuroimmunol*. 2018;315:1–8. doi: 10.1016/j.jneuroim.2017.12.004.
53. Brophy G.M., Bell R., Claassen J., Alldredge B., Bleck T.P., Glauser T., LaRoche S.M., Riviello J.J., Shutter L., Sperling M.R., et al. Guidelines for the Evaluation and Management of Status Epilepticus. *Neurocrit. Care*. 2012;17:3–23. doi: 10.1007/s12028-012-9695-z.
54. Cost of Status Epilepticus: A Systematic Review—ScienceDirect (accessed on 18 December 2018).
55. Claassen J., Mayer S.A., Kowalski R.G., Emerson R.G., Hirsch L.J. Detection of electrographic seizures with continuous EEG monitoring in critically ill patients. *Neurology*. 2004;62:1743–1748. doi: 10.1212/01.WNL.0000125184.88797.62.

56. Malu C.K., Kahamba D.M., Walker T.D. Efficacy of sublingual lorazepam versus intrarectal diazepam for prolonged convulsions in Sub-Saharan Africa. *J Child Neurol.* 2014;29(7):895–902. doi: 10.1177/0883073813493501.
57. Merry C., Mulcahy F., Barry M. Saquinavir interaction with midazolam: pharmacokinetic considerations when prescribing protease inhibitors for patients with HIV disease. *AIDS.* 1997;11(2):268–269.
58. Ahmad S., Ellis J.C., Kamwendo H. Efficacy and safety of intranasal lorazepam versus intramuscular paraldehyde for protracted convulsions in children: an open randomised trial. *Lancet.* 2006;367(9522):1591–1597. doi: 10.1016/S0140-6736(06)68696-0.
59. Zeiler F.A. Early use of the NMDA receptor antagonist ketamine in refractory and superrefractory status epilepticus. *Crit Care Res Pract.* 2015;2015:831260. doi: 10.1155/2015/831260.
60. Portela J.L., Garcia P.C., Piva J.P. Intramuscular midazolam versus intravenous diazepam for treatment of seizures in the pediatric emergency department: a randomized clinical trial. *Med Intensiva.* 2015;39(3):160–166. doi: 10.1016/j.medin.2014.04.003.
61. Hsu A.J., Carson K.A., Yung R. Severe prolonged sedation associated with coadministration of protease inhibitors and intravenous midazolam during bronchoscopy. *Pharmacotherapy.* 2012;32(6):538–545. doi: 10.1002/j.1875-9114.2011.01045.x.
62. Drislane F.W. Presentation, evaluation, and treatment of nonconvulsive status epilepticus. *Epilepsy Behav.* 2000;1(5):301–314. doi: 10.1006/ebeh.2000.0100.
63. Chamberlain J.M., Altieri M.A., Futterman C. A prospective, randomized study comparing intramuscular midazolam with intravenous diazepam for the treatment of seizures in children. *Pediatr Emerg Care.* 1997;13(2):92–94. doi: 10.1097/00006565-199704000-00002.
64. Crawley J., Waruiru C., Mithwani S. Effect of phenobarbital on seizure frequency and mortality in childhood cerebral malaria: a randomised, controlled intervention study. *Lancet.* 2000;355(9205):701–706. doi: 10.1016/S0140-6736(99)07148-2.