JOURNAL OF INTERNATIONAL CRISIS AND RISK COMMUNICATION RESEARCH
ISSN: 2576-0017
2024, VOL 7, NO S8

Critical Moments: Trauma Nursing in High-Pressure Emergency Settings

Sahari Mohammed Aloufi[1]
Tahani Mubarak Fayez Alaklabi[2]
Manal Obaid Matar Alenezi[3]
Maha Abdo Mohammed Al-Somali[4]
Mona Abdo Mohammed Alsomali[5]
Bandar Hamad Shunayf[6]
Rajaa Mohammed Al Salem[7]
Jamelah Aowad Barak Almutairi[8]

1. Nursing Specialist
   Shorn Health Center
   Madinah
2. Nursing
   Specialized Dental Center
   Riyadh
3. Midwife
   Alyamammah Hospital
   Riyadh
4. Nursing Technician
   Dhahran General Hospital
   Al-Khobar
5. Nursing Technician
   Alolaya And Alsolamanyah Primary Health Care
   Riyadh
6. Nursing Technician
   Najran General Hospital
   Najran
7. Nursing Technician
   Dental Complex East Riyadh
   Riyadh
8. Nursing Specialist
   Riyadh Specialized Dental Center
   Riyadh

Abstract
Trauma nursing in high-pressure emergency settings represents a critical and evolving field that combines advanced clinical expertise, rapid decision-making, and emotional resilience to manage patients with severe injuries and life-threatening conditions. This literature review explores the multifaceted roles and challenges faced by trauma nurses, emphasizing their contributions to patient stabilization, care coordination, and holistic recovery support. Drawing from global and historical contexts, the review traces the evolution of trauma nursing, highlighting milestones such as battlefield medicine, the development of Advanced Trauma Life Support protocols, and the rise of structured education and training programs like the Trauma Nursing Core Course. The analysis delves into the unique demands of high-pressure emergency environments, where time-sensitive interventions, resource constraints, and unpredictable scenarios demand exceptional adaptability and teamwork. It examines the physical and psychological stressors that affect trauma nurses, including secondary traumatic stress and burnout, and underscores the importance of institutional support and mental health initiatives to foster resilience. Evidence-based recommendations, including continuing education, interprofessional collaboration, and mental health support, are presented as strategies to enhance both patient outcomes and nursing well-being. By synthesizing current research and best practices, this review underscores the indispensable role of trauma nurses in modern healthcare systems, advocating for ongoing advancements in training, resources, and systemic support to meet the demands of this high-stakes field.

Introduction

Trauma is a significant global health challenge, particularly affecting children and young people. The World Health Organization (WHO) identifies trauma as the leading cause of death in individuals under 30 years old (WHO, 2021). This includes injuries from road traffic accidents, falls, interpersonal violence, and other unintentional injuries.In the UK, the statistics underscore the gravity of trauma's impact. Annually, approximately 16,000 people lose their lives

due to traumatic injuries, highlighting the public health burden of this issue (Lucena-Amaro et al., 2022). Among those who survive, many endure life-altering consequences, including severe disabilities and prolonged recovery periods. These injuries often require extensive rehabilitation and may lead to a reduced quality of life, chronic pain, and psychological challenges such as post-traumatic stress disorder (PTSD) (WHO, 2021).Beyond individual suffering, trauma places a substantial strain on healthcare systems. Survivors with complex injuries often require critical care, surgeries, and long-term support, contributing to significant healthcare costs. Effective trauma care systems, including rapid response, advanced surgical interventions, and comprehensive rehabilitation services, are essential to improve survival rates and outcomes (Magruder et al., 2017).

Traumatic injuries encompass a range of severe and complex conditions, with devastating impacts on physical and neurological health. Traumatic Spinal Cord Injury (SCI) results in the loss of motor, sensory, and autonomic functions below the injury level, with causes including vehicular accidents and falls. Initial mechanical trauma often leads to progressive damage due to secondary injury mechanisms such as inflammation and neurodegeneration (Lima et al., 2022). Effective management includes surgical decompression and stabilization, but challenges persist due to the formation of cystic cavities and glial scarring that inhibit regeneration (Rupp, 2021).Traumatic Brain Injury (TBI) involves direct mechanical damage to brain tissues, often accompanied by diffuse axonal injuries. It causes a cascade of cellular disruptions, leading to permanent disabilities, cognitive impairments, or persistent vegetative states (Mckee & Daneshvar, 2018). Emerging treatments focus on minimizing damage through therapies targeting mitochondrial functions and inflammation (Hamblin et al., 2011).

Trauma nurses are the backbone of high-pressure environments, combining clinical expertise, emotional intelligence, and organizational skills to save lives and improve outcomes. Their ability to manage critical situations and provide holistic care makes them indispensable in trauma and emergency care settings.Emergency and critical care (ECC) units play a vital role in healthcare, dedicated to managing patients with urgent and life-threatening conditions. Nurses in these settings face significant stress due to the high demands of their work environment and the complexity of their responsibilities (Liu et al., 2023). During public health crises like the COVID-19 pandemic, ECC nurses took on critical duties despite the risk of exposure. Their efforts included identifying and managing cases, implementing safety protocols, collecting medical samples, providing respiratory and circulatory support such as ECMO, and delivering intensive bedside care (Han et al., 2022). As key patient caregivers, they maintain close contact and significantly contribute to the department's operations. Collaborative efforts between ECC nurses and physicians have been shown to improve outcomes by reducing emergency wait times, enhancing patient satisfaction, lowering mortality rates, and cutting healthcare costs, outperforming physician-only care models (Woo et al., 2017).Trauma nursing in high-pressure emergency settings represents a critical component of modern healthcare, blending rapid decision-making, emotional resilience, and technical expertise to manage complex patient needs. This specialized field has evolved from battlefield medicine to a structured profession influenced by advancements in emergency protocols and interdisciplinary collaboration (Henshall et al., 2022). Trauma nursing originated from battlefield medicine, where rapid response and triage were essential. Over time, it has evolved into a structured profession incorporating evidence-based emergency protocols and interdisciplinary collaboration (Baik et al., , 2024). The introduction of programs like Advanced Trauma Care for Nurses has standardized care and improved outcomes through enhanced knowledge and technical proficiency (Awwad et al., 2021). Nurses in these roles face multifaceted challenges, including emotional strain, resource constraints, and high patient volumes, necessitating a balance between clinical precision and psychological support. Through evidence-based practices, continuous education, and innovations in emergency care systems, trauma nursing remains pivotal in enhancing patient outcomes during critical moments (Van Way, 2016).

The objectives of this review are to analyze the evolution and current practices of trauma nursing within high-pressure emergency settings, emphasizing its critical role in managing severe and complex injuries. It seeks to identify the multifaceted responsibilities of trauma nurses, including patient assessment, intervention, care coordination, and psychological support, while exploring the historical context that shaped the profession. Furthermore, the review aims to highlight the challenges faced by trauma nurses, such as resource constraints, emotional strain, and secondary traumatic stress, and to evaluate strategies for addressing these issues. By integrating evidence-based practices, training advancements, and multidisciplinary collaboration, the review aspires to propose actionable recommendations for improving trauma care outcomes and supporting the resilience and well-being of trauma nurses in their demanding roles.

1. Historical Context and Evolution of Trauma Nursing

Trauma nursing has evolved significantly over the past century, shaped by advancements in medical science, responses to global conflicts, and the growing recognition of trauma as a critical public health issue. Understanding its historical context provides insight into how this specialty emerged and adapted to meet the challenges of increasingly complex emergency care.In the late 19th and early 20th centuries, trauma care was rudimentary, with

little formal training for nurses in emergency medicine. Early trauma nursing practices were heavily influenced by wartime experiences (Van Way, 2016). During World War I (1914–1918) and World War II (1939–1945), nurses played a pivotal role in caring for injured soldiers. These experiences underscored the need for rapid assessment, triage, and treatment of life-threatening injuries in high-pressure environments. Field hospitals became laboratories for trauma care innovation. Nurses working on battlefields developed techniques to stabilize patients under resource-constrained conditions, such as controlling hemorrhage and preventing infection. These practices laid the groundwork for modern trauma care principles (Brown, 2015; Harris, 2014).

The mid-20th century saw the formalization of emergency and trauma nursing. The post-war period brought advancements in medicine, including the development of antibiotics and improvements in surgical techniques, which enhanced survival rates for trauma patients (Moparthi et al., 2024). In the 1960s, the establishment of civilian emergency medical services (EMS) systems revolutionized trauma care. This period marked the transition from informal on-the-job training to structured education for nurses in emergency medicine (Teman et al., 2014). The introduction of Advanced Trauma Life Support (ATLS) protocols in the 1970s further standardized trauma care, emphasizing the critical role of nurses in delivering timely interventions (Radvinsky et al., 2012).

The evolution of trauma nursing has been shaped by several key milestones, including the establishment of trauma centers in the 1960s and 1970s, the development of triage systems to prioritize patient care effectively, and the formation of emergency nursing associations, which have advanced professional standards and collaboration in the field. The establishment of designated trauma centers in hospitals recognized the need for specialized facilities and trained personnel to manage trauma patients. Nurses became integral members of multidisciplinary teams (Maull et al., 1986). Triage, the process of prioritizing patients based on injury severity, was formalized during wartime but adapted for civilian emergency departments (EDs) (Lampi, 2017). Trauma nurses became proficient in assessing and categorizing patients to ensure optimal use of resources.The Emergency Nurses Association (ENA), established in 1970, has been a pivotal organization advocating for emergency nursing excellence through education, advocacy, and certification. One of ENA's most influential contributions is the development of the Trauma Nursing Core Course (TNCC), a certification program designed to provide nurses with a systematic approach to trauma care. The TNCC emphasizes evidence-based practices and is often a mandatory requirement for emergency nurses in trauma centers. Moreover, it is recognized and recommended by institutions like the American College of Surgeons for hospitals aspiring to trauma center verification (Silva & Whalen, 2014). ENA has also led initiatives to elevate the professionalism of emergency nursing. Its advocacy encompasses policy issues, trauma prevention, and the development of safety protocols, significantly influencing care standards in emergency settings. These efforts have contributed to broader access to emergency care training, such as pediatric and critical care specialties, enriching the skillsets of nurses globally (Milbrath & Snyder, 2021). Through these initiatives, ENA has advanced emergency nursing by ensuring nurses are prepared for the complex demands of high-pressure environments, reinforcing the importance of continuous education and professional growth.

Recently, global health crises such as natural disasters, terrorist attacks, and pandemics have underscored the critical role of trauma nursing in disaster response and mass casualty incidents (MCIs). These events have accelerated advancements in trauma education and readiness, shaping the development of specialized training, interdisciplinary collaboration, and policy frameworks (Loke et al., 2021). Nurses serve as vital first responders during MCIs, where the demand for care exceeds available resources. Evidence-based guidelines tailored for nurses in prehospital and hospital environments emphasize the importance of disaster preparedness and efficient triage systems (Olchin& Krutz, 2012). Disaster simulations have emerged as a valuable tool in nursing education, providing opportunities for students to develop triage, communication, and teamwork skills. For instance, participation in MCI simulations significantly improved students' self-perceived knowledge and preparedness (Zinan et al., 2015). Advanced training curricula, such as the Advanced Disaster Medical Response (ADMR) Course, equip nurses with the core competencies needed for effective disaster management (Chang & Briggs, 2011). Events like the COVID-19 pandemic demonstrated the necessity of incorporating public health principles into trauma nursing. Nurses' roles expanded to include community education, infection control, and psychosocial support for both patients and healthcare teams (Rodriguez-Arrastia et al., 2021).

2. High-Pressure Emergency Settings in Trauma Nursing

Emergency settings are among the most demanding environments in healthcare, characterized by critical time sensitivity, unpredictable conditions, high patient volumes, and resource limitations. These environments require trauma nurses to perform under immense pressure while ensuring the highest standard of care. Understanding the dynamics of these settings is crucial to appreciating the skills and resilience required of trauma nurses (Abu-Alhaija& Gillespie, 2022). The urgency of care in emergency settings often requires trauma nurses to make rapid, life-saving decisions. Effective time management is critical to ensure patients receive timely interventions. Studies demonstrate that shorter response times and quicker preparation in emergency operations significantly enhance

rescue success rates and patient outcomes (Hu et al., 2023). Trauma nurses routinely prioritize procedures based on the severity of injuries using advanced triage systems to manage limited time effectively (Lee et al., 2013). Emergency settings are marked by unpredictable scenarios that demand flexibility and rapid adaptation. Nurses often face chaotic situations, such as mass casualty incidents or complex trauma cases, where the care environment can feel like a "war zone." These conditions contribute to chronic stress and secondary traumatic stress among nurses (Wolf et al., 2020). Unpredictability necessitates constant readiness for a wide range of medical emergencies, from trauma to mental health crises (Molloy et al., 2020). Overcrowding in emergency departments places significant stress on nurses and can compromise patient care. The simultaneous need to address multiple critical cases requires exceptional organizational and multitasking skills (Ferreira &Cividini, 2023). In high-volume settings, trauma nurses often experience a greater workload, contributing to physical exhaustion and burnout, which can affect their well-being and the quality of care provided (Adriaenssens et al., 2012).Limited access to critical resources, including personnel, equipment, and space, further complicates trauma nursing in emergencies. In rural or remote settings, these constraints are amplified, impacting patient outcomes significantly (Morgan & Calleja, 2020). Strategic allocation of scarce resources is essential to maximize patient survival during mass casualty incidents and other high-demand scenarios (Hirshberg et al., 2001).

3. The Role of Trauma Nurses in High-Pressure Environments

Trauma nurses operate in dynamic and high-pressure environments where their expertise is essential for patient stabilization, comprehensive care, and recovery support. Their roles are multifaceted, encompassing direct patient care, emotional support, and collaborative efforts in interdisciplinary teams (Ashworth et al., 2023).

Patient assessment and intervention: Trauma nurses are responsible for rapid, accurate assessments of patients, which include evaluating the mechanism of injury, vital signs, and potential complications. This initial care is critical to identifying life-threatening conditions and prioritizing treatments. They perform key interventions such as administering medications, managing airways, and initiating resuscitation. This ensures immediate response to the unique demands of trauma care (Baik et al., 2024).

Wound Care and Management: Trauma nurses provide specialized wound care, including cleaning, dressing, and monitoring for signs of infection. In cases of burns or severe injuries, they coordinate with plastic surgeons or wound care specialists for advanced treatment. They ensure thorough irrigation and debridement to remove contaminants, which is essential for effective wound management (Bonham, 2016). In cases involving severe injuries or burns, trauma nurses collaborate closely with plastic surgeons and wound care specialists to coordinate advanced treatments. This multidisciplinary approach ensures comprehensive care tailored to complex injuries (Steele, 2006). Specific interventions, such as the use of negative pressure wound therapy (NPWT), are often employed in trauma cases to promote granulation tissue formation and wound closure (Clevenger & Tepas, 1997). Trauma nurses monitor wounds for signs of infection, such as redness, swelling, or purulent discharge, to intervene promptly. Perianesthesia nurses in orthopedic trauma cases, for instance, focus on infection control practices due to the heightened risk of complications like osteomyelitis (Mamaril et al., 2007).Also, adherence to evidence-based protocols for dressing changes and use of sterile techniques is integral to minimizing infection risks and improving recovery times (Jennings & Mitchell, 2017).

Care coordination and continuity: Trauma nurses play a vital role in coordinating the care provided by various medical teams, ensuring seamless transitions from emergency departments to surgical units or intensive care facilities (Liu et al., 2019). For pediatric trauma patients, nurses specialize in adapting care protocols to meet developmental needs while mitigating emotional and physical stress (Wurster et al., 2009).

Management of complications:Trauma nurses are instrumental in preventing and managing complications, such as hypothermia, by using interventions to maintain body temperature and stabilize patients during resuscitation and surgery (van Veelen& Brodmann Maeder, 2021). They actively monitor patients for secondary trauma or psychological distress, intervening as needed to improve long-term outcomes (Adriaenssens et al., 2012).

Emotional and psychological support: Beyond physical care, trauma nurses provide critical emotional support to both patients and their families, addressing fears, uncertainties, and grief associated with traumatic events (Leng et al., 2022). Their role often includes mitigating secondary trauma effects, such as compassion fatigue and burnout, by employing resilience and coping strategies (Wentzel &Brysiewicz, 2014).

Collaboration: Trauma nurses act as central communication hubs, ensuring that all team members are informed about patient status, interventions, and critical care decisions. Structured communication, such as pre-trauma arrival briefings, has been shown to improve team concordance and task efficiency (Steinemann et al., 2016). Regular rounds involving trauma nurses and other specialists have improved patient outcomes and collaboration. For instance, structured trauma nursing rounds led to over 450 care improvements across a series of patients (Jennings & Mitchell, 2017).

Championing patient-centered decisions: Nurses frequently advocate for appropriate care decisions, such as prioritizing pain management or timely interventions. Advocacy is especially crucial during end-of-life decisions, where nurses facilitate discussions between families and the care team, balancing medical possibilities with patient preferences (Barndt, 2018).

4. Challenges and Stressors in Trauma Nursing

Trauma nursing in high-pressure emergency settings presents a unique set of challenges due to the high-stakes and emotionally charged nature of the work. Emergency nurses frequently encounter life-threatening situations and critical events, which can lead to emotional exhaustion and post-traumatic stress. These high-pressure scenarios, especially those involving aggression or patient mortality, are significant predictors of post-traumatic stress symptoms (de Wijn& van der Doef, 2020).

Secondary traumatic stress occurs when nurses experience emotional distress from exposure to patients' trauma. A significant proportion of emergency nurses report symptoms of STS, characterized by intrusive thoughts, emotional numbness, and hyperarousal (Alshammari et al., 2024). Factors contributing to STS include prolonged exposure to traumatic patient cases, lack of coping mechanisms or institutional support, and feelings of helplessness during severe patient deterioration or death (Duffy et al., 2015). The quality of trauma care can be compromised by inadequate resources, leading to dissatisfaction among nurses. Resource-related challenges include insufficient staffing or equipment during high-demand periods and poorly designed environments that disrupt efficient workflows and communication (Peng et al., 2023).

Nurses often experience psychological strain due to witnessing severe patient injuries, unexpected deaths, or adverse outcomes. The unpredictability of trauma cases adds an additional layer of stress, as even well-trained nurses sometimes feel unprepared to address every scenario effectively (Alzghoul, 2014). While trauma nurses are trained to prioritize patient care, the emotional toll of suppressing personal feelings to maintain professionalism is significant. This duality can lead to long-term emotional fatigue and reduced job satisfaction (Missouridou, 2017).

5. Recommendations for Improvement

Improving trauma nursing in high-pressure emergency settings is vital to enhance patient outcomes and reduce provider burnout. Trauma nurses often face dynamic and high-stakes situations requiring both technical expertise and emotional resilience. Recommendations for improvement span education, inter-professional collaboration, system efficiency, and mental health support (Haugland et al., 2023).

Continuing education programs focused on trauma-specific skills, such as advanced life support techniques, triage protocols, and rapid decision-making, are essential for maintaining nursing competence in emergency settings. Simulation-based training has been shown to improve clinical skills and teamwork under pressure (Castner et al., 2024). Trauma education programs, like the Advanced Trauma Life Support (ATLS) or Advanced Trauma Care for Nurses, have been shown to improve nurses' knowledge and skills in trauma triage and management. Participants reported increased confidence and preparedness to handle complex trauma scenarios (Awwad et al., 2021; Ireland et al., 2017). Innovative methods like in-situ simulation allow nurses to practice trauma resuscitation in realistic settings, enhancing teamwork and decision-making skills. This approach not only fosters confidence but also improves readiness to manage real-life trauma cases (Bellino et al., 2020). Multidisciplinary training aligns diverse roles, fostering understanding and synergy among healthcare professionals. Better coordination between emergency and surgical teams has been linked to improved trauma management outcomes (Alexandrino et al., 2023). Multidisciplinary training programs have been effective in reducing emergency department overcrowding and improving patient management outcomes, such as faster triage and reduced emergency department stay times (Heydari et al., 2019). Training emergency nurses in trauma care improves not only technical skills but also reduces risks of re-traumatization in vulnerable patients. Programs integrating psychological and social trauma aspects can enhance holistic care delivery (Akhtar et al., 2022). Training not only increases team efficiency but also boosts morale and job satisfaction, creating a more supportive work environment and reducing burnout among healthcare professionals (Roberts et al., 2014).

Effective communication and collaboration among healthcare professionals improve patient care during emergencies. Trauma teams should implement structured communication protocols like SBAR (Situation, Background, Assessment, and Recommendation) to ensure clarity and reduce errors. Structured communication tools help ensure clarity and reduce errors during handovers and critical moments. Implementation improves information transfer and reduces misunderstandings in high-pressure environments (Miller et al., 2012). Programs like TeamSTEPPS improve collaboration and communication within emergency teams, boosting perceptions of teamwork and patient safety. Pre- and post-training evaluations have demonstrated measurable improvements in team dynamics (Matzke et al., 2021). Trauma simulations in realistic settings train teams to communicate effectively under pressure. These exercises have been shown to improve communication, leadership, and task management skills, with positive impacts observed during real emergencies (Caldwell et al., 2020).

Nurses frequently experience burnout and PTSD due to exposure to traumatic events and workplace stressors. Studies report significant rates of emotional exhaustion and PTSD symptoms among trauma and critical care nurses (Li et al., 2021). Addressing the mental health needs of trauma nurses is critical given their exposure to high stress and traumatic events. Institutions should provide access to counseling services, peer support programs, and regular debriefings. Evidence indicates that such interventions lower the risk of burnout and post-traumatic stress disorder among emergency nurses (Flaubert et al., 2021). Access to mental health counseling and structured peer support programs significantly helps nurses cope with traumatic experiences. Such interventions improve psychological well-being and reduce the incidence of burnout and compassion fatigue (Fu et al., 2018). Mindfulness-based interventions (MBIs) effectively reduce stress, depression, and PTSD symptoms among nurses. Training in mindfulness helps enhance emotional resilience and coping mechanisms, fostering better mental health outcomes (Ramachandran et al., 2022). Trauma nurses also benefit from regular debriefings that provide opportunities to process experiences and receive emotional support. This reduces feelings of isolation and aids in emotional recovery after critical incidents (Darnell et al., 2018). Higher levels of social support, both personal and professional, are associated with lower levels of burnout and PTSD in nurses. Encouraging supportive work environments and relationships can buffer against stress (Lee & Kim, 2016). Training trauma nurses in delivering brief mental health interventions for themselves and peers has shown promise in improving their coping skills and reducing mental health symptoms (Darnell et al., 2018). Finally, institutions that implement structured programs addressing workplace stress and trauma exposure, such as resilience training and improved staffing models, enhance the well-being of their nursing staff (Jenkins & Elliott, 2004).

In conclusion, trauma nursing in high-pressure emergency settings is a cornerstone of modern healthcare, requiring a blend of clinical precision, rapid decision-making, and emotional resilience to address complex and life-threatening injuries. Trauma nurses play a vital role in patient stabilization, holistic recovery, and care coordination, often under significant stress and resource limitations. Despite these challenges, advancements in training programs, evidence-based practices, and multidisciplinary collaboration have improved both patient outcomes and professional standards in the field. Addressing the psychological and emotional toll on trauma nurses is critical for sustaining their well-being and effectiveness. Strategies such as continuing education, institutional support, and mental health interventions are essential for reducing burnout and secondary traumatic stress. By fostering resilience and providing adequate resources, healthcare systems can empower trauma nurses to deliver high-quality care while enhancing their professional satisfaction. As the demands on emergency healthcare grow, the role of trauma nurses remains indispensable, underscoring the need for ongoing investment in their development and support to ensure optimal outcomes for patients and caregivers alike.

References

Abu-Alhaija, D.M. and Gillespie, G.L., 2022. Critical clinical events and resilience among emergency nurses in 3 trauma hospital-based emergency departments: A cross-sectional study. Journal of emergency nursing, 48(5), pp.525-537.

Adriaenssens, J., De Gucht, V. and Maes, S., 2012. The impact of traumatic events on emergency room nurses: Findings from a questionnaire survey. International journal of nursing studies, 49(11), pp.1411-1422.

Akhtar, Z., Khan, M.H. and Noureen, S., Jamila, Asifa Shahzadi (2022). Trauma nurses' Role and Emergency Management skills at Tertiary Care Hospital, Lahore. Saudi J Nurs Health Care, 5(2), pp.36-38.

Alexandrino, H., Martinho, B., Ferreira, L. and Baptista, S., 2023. Non-technical skills and teamwork in trauma: from the emergency department to the operating room. Frontiers in medicine, 10, p.1319990.

Alshammari, B., Alanazi, N.F., Kreedi, F., Alshammari, F., Alkubati, S.A., Alrasheeday, A., Madkhali, N., Alshara, A., Bakthavatchaalam, V., Al-Masaeed, M. and Alshammari, S.K., 2024. Exposure to secondary traumatic stress and its related factors among emergency nurses in Saudi Arabia: a mixed method study. BMC nursing, 23(1), p.337.

Alzghoul, M.M., 2014. The experience of nurses working with trauma patients in critical care and emergency settings: A qualitative study from Scottish nurses' perspective. International journal of orthopaedic and trauma nursing, 18(1), pp.13-22.

Ashworth, H., Lewis-O'Connor, A., Grossman, S., Brown, T., Elisseou, S. and Stoklosa, H., 2023. Trauma-informed care (TIC) best practices for improving patient care in the emergency department. International Journal of Emergency Medicine, 16(1), p.38.

Awwad, K., Ng, Y.G., Lee, K., Lim, P.Y. and Rawajbeh, B., 2021. Advanced Trauma Life Support/Advanced Trauma Care for Nurses: A systematic review concerning the knowledge and skills of emergency nurse related to trauma triage in a community. International Emergency Nursing, 56, p.100994.

Baik, D., Yi, N., Han, O. and Kim, Y., 2024. Trauma nursing competency in the emergency department: a concept analysis. BMJ Open, 14(6), p.e079259.

Barndt, S.N., 2018. Death in trauma: the role of the ACNP in patient advocacy and familial support in end-of-life care decision-making. Journal of Trauma Nursing| JTN, 25(3), pp.171-176.
Bellino, A., Gordon, A.J. and Schertzer, K., 2020. Owning the Trauma Bay: Teaching Trauma Resuscitation to Emergency Medicine Residents and Nurses through In-situ Simulation. Journal of Education & Teaching in Emergency Medicine, 5(4), p.S108.
Bonham, J., 2016. Assessment and management of patients with minor traumatic wounds. Nursing Standard (2014+), 31(8), p.60.
Brown, W.J., 2015. Nursing in the 8th Evacuation Hospital, 1942-1945. US Army Medical Department Journal.
Caldwell, K.E., Lulla, A., Murray, C.T., Handa, R.R., Romo, E.J., Wagner, J.W., Wise, P.E., Leonard, J.M. and Awad, M.M., 2021. Multi-Disciplinary Trauma Evaluation and Management Simulation (MD-TEAMS) training for emergency medicine and general surgery residents. The American Journal of Surgery, 221(2), pp.285-290.
Castner, J., Zazzera, E.A. and Burchill, C.N., 2024. Continuing Education Certificate in Trauma Skills Among Emergency Nurses: A National Sample Survey Analysis. Journal of Trauma Nursing| JTN, 31(6), pp.290-300.
Chang, L. and Briggs, S.M., 2011. (A105) A Core Curriculum for Nurses in Disaster Preparedness and Response. Prehospital and Disaster Medicine, 26(S1), pp.s29-s29.
Clevenger, F.W. and Tepas III, J.J., 1997. Preoperative management of patients with major trauma injuries. AORN journal, 65(3), pp.583-594.
Darnell, D.A., Parker, L.E., Wagner, A.W., Dunn, C.W., Atkins, D.C., Dorsey, S. and Zatzick, D.F., 2019. Task-shifting to improve the reach of mental health interventions for trauma patients: findings from a pilot study of trauma nurse training in patient-centered activity scheduling for PTSD and depression. Cognitive behaviour therapy, 48(6), pp.482-496.
De Wijn, A.N. and Van der Doef, M.P., 2020. Patient-related stressful situations and stress-related outcomes in emergency nurses: A cross-sectional study on the role of work factors and recovery during leisure time. International Journal of Nursing Studies, 107, p.103579.
Duffy, E., Avalos, G. and Dowling, M., 2015. Secondary traumatic stress among emergency nurses: a cross-sectional study. International emergency nursing, 23(2), pp.53-58.
Ferreira, D.D.A.B. and Cividini, F.R., 2023. Challenges faced by nursing in the emergency room for the care of trauma victims: An integrative literature review. Research, Society and Development, 12(5), pp.e8512541504-e8512541504.
Flaubert, J.L., Le Menestrel, S., Williams, D.R., Wakefield, M.K. and National Academies of Sciences, Engineering, and Medicine, 2021. Supporting the Health and Professional Well-Being of Nurses. In The Future of Nursing 2020-2030: Charting a Path to Achieve Health Equity. National Academies Press (US).
Fu, C.Y., Yang, M.S., Leung, W., Liu, Y.Y., Huang, H.W. and Wang, R.H., 2018. Associations of professional quality of life and social support with health in clinical nurses. Journal of nursing management, 26(2), pp.172-179.
Hamblin, M., Huang, Y.Y., Wu, Q., Xuan, W., Ando, T., Xu, T., Sharma, S. and Kharkwal, G., 2011. Low-level light therapy aids traumatic brain injury. SPIE Newroonm.
Han, P., Duan, X., Zhao, S., Zhu, X. and Jiang, J., 2022. Nurse's psychological experiences of caring for severe COVID-19 patients in intensive care units: a qualitative Meta-synthesis. Frontiers in Public Health, 10, p.841770.
Harris, K., 2014. New horizons: Australian nurses at work in World War I. Endeavour, 38(2), pp.111-121.
Haugland, W.A., Crenshaw, J.T. and Gilder, R.E., 2023. Implementing a resilience bundle for emergency nurses: An evidence-based practice project. Journal of Emergency Nursing, 49(1), pp.40-49.
Henshall, C., Davey, Z. and Jackson, D., 2020. Nursing resilience interventions–A way forward in challenging healthcare territories. Journal of Clinical Nursing, 29(19-20), p.3597.
Heydari, F., Mohamadirizi, S. and Nasr-Esfahani, M., 2019. The effects of multidisciplinary education for nurses and physicians on the management of patients with multiple trauma. Nursing and Midwifery Studies, 8(4), pp.189-194.
Hirshberg, A., Holcomb, J.B. and Mattox, K.L., 2001. Hospital trauma care in multiple-casualty incidents: a critical view. Annals of emergency medicine, 37(6), pp.647-652.
Hu, P., Wang, H. and Zheng, Y., 2023. High-quality nursing benefits patients with emergency trauma undergoing surgical debridement and suturing and helps relieve pain. American Journal of Translational Research, 15(11), p.6495.
Iattoni, M., Ormazabal, M., Luvini, G. and Uccella, L., 2022. Effect of structured briefing prior to patient arrival on interprofessional communication and collaboration in the trauma team. Open Access Emergency Medicine, pp.385-393.

Ireland, S., Cross, R., Decker, K. and Mitra, B., 2017. Perceptions of an educational programme for registered nurses who work at non-major trauma services in Victoria, Australia: The Nursing Emergency eXternal Trauma Programme. Australasian emergency nursing journal, 20(3), pp.131-138.
Jenkins, R. and Elliott, P., 2004. Stressors, burnout and social support: nurses in acute mental health settings. Journal of advanced nursing, 48(6), pp.622-631.
Jennings, F.L. and Mitchell, M., 2017. Intensive care nurses' perceptions of Inter Specialty Trauma Nursing Rounds to improve trauma patient care—A quality improvement project. Intensive and Critical Care Nursing, 40, pp.35-43.
Lampi, M., 2017. TRIAGE: Management of the trauma patient (Doctoral dissertation, Linköping University Electronic Press).
Lee, G.A., Murray, A., Bushnell, R. and Niggemeyer, L.E., 2013. Challenges developing evidence-based algorithms for the trauma reception and resuscitation project. International emergency nursing, 21(2), pp.129-135.
Lee, H.J. and Kim, J.Y., 2016. Effects of psychiatric nurses' secondary traumatic stress and compassion satisfaction on burnout: the moderating effect of social support. Journal of Korean Academy of Psychiatric and Mental Health Nursing, 25(4), pp.399-408.
Leng, Y., Wu, Y., Wang, Z., Zhou, X. and Liao, J., 2022. A qualitative study exploring barriers and facilitators to establishing nurse-led, multidisciplinary psychological care for trauma patients: experiences from doctors and nurses. BMC nursing, 21(1), p.191.
Li, P., Kuang, H. and Tan, H., 2021. The occurrence of post-traumatic stress disorder (PTSD), job burnout and its influencing factors among ICU nurses. American journal of translational research, 13(7), p.8302.
Lima, R., Monteiro, A., Salgado, A.J., Monteiro, S. and Silva, N.A., 2022. Pathophysiology and therapeutic approaches for spinal cord injury. *International Journal of Molecular Sciences*, *23*(22), p.13833.
Liu, S., Zhang, Y., Liu, Y., Han, P., Zhuang, Y. and Jiang, J., 2023. The resilience of emergency and critical care nurses: a qualitative systematic review and meta-synthesis. Frontiers in Psychology, 14, p.1226703.
Liu, S.I., Curren, J., Leahy, N.E., Sobocinski, K., Zambardino, D., Shikar, M.M., Vasquez, C., Miluszusky, B. and Winchell, R.J., 2019. Trauma response nurse: Bringing critical care experience and continuity to early trauma care. Journal of Trauma Nursing| JTN, 26(4), pp.215-220.
Loke, A.Y., Guo, C. and Molassiotis, A., 2021. Development of disaster nursing education and training programs in the past 20 years (2000–2019): A systematic review. Nurse education today, 99, p.104809.
Lucena-Amaro, S. and Zolfaghari, P., 2022. Trauma nursing 1: an overview of major trauma and the care pathway. Nursing Times, 118(11), pp.1-15.
Magruder, K.M., McLaughlin, K.A. and Elmore Borbon, D.L., 2017. Trauma is a public health issue. European journal of psychotraumatology, 8(1), p.1375338.
Mamaril, M.E., Childs, S.G. and Sortman, S., 2007. Care of the orthopaedic trauma patient. Journal of PeriAnesthesia Nursing, 22(3), pp.184-194.
Matzke, C.M., DeGennaro, R. and Howie-Esquivel, J., 2021. Incorporating TeamSTEPPS training to improve staff collaboration in an academic level I emergency and trauma center. International emergency nursing, 55, p.100959.
Maull KI, Schwab CW, McHenry SD, Leavy P, Carl L, Woo P, Overholt S, Sinclair T, Aprahamian C. 1986. Trauma center verification. Journal of Trauma and Acute Care Surgery, 26(6), pp.521-524.
Mckee, A.C. and Daneshvar, D.H., 2015. The neuropathology of traumatic brain injury. Handbook of clinical neurology, 127, pp.45-66.
Milbrath, G. and Snyder, A., 2021. The emergency nurses association: 50 years of advocacy and advancement. Journal of emergency nursing, 47(2), pp.224-238.
Miller, D., Crandall, C., Washington III, C. and McLaughlin, S., 2012. Improving teamwork and communication in trauma care through in situ simulations. Academic Emergency Medicine, 19(5), pp.608-612.
Missouridou, E., 2017. Secondary posttraumatic stress and nurses' emotional responses to patient's trauma. Journal of trauma nursing| JTN, 24(2), pp.110-115.
Molloy, L., Fields, L., Trostian, B. and Kinghorn, G., 2020. Trauma-informed care for people presenting to the emergency department with mental health issues. Emergency Nurse, 28(2).
Moparthi, K.P., Javed, H., Kumari, M., Pavani, P., Paladini, A., Saleem, A., Ram, R. and Varrassi, G., 2024. Acute Care Surgery: Navigating Recent Developments, Protocols, and Challenges in the Comprehensive Management of Surgical Emergencies. Cureus, 16(1).
Morgan, J.M. and Calleja, P., 2020. Emergency trauma care in rural and remote settings: Challenges and patient outcomes. International emergency nursing, 51, p.100880.
Olchin, L. and Krutz, A., 2012. Nurses as first responders in a mass casualty: are you prepared?. Journal of Trauma Nursing| JTN, 19(2), pp.122-129.

Peng, X., Ye, Y., Ding, X. and Chandrasekaran, A., 2023. The impact of nurse staffing on turnover and quality: An empirical examination of nursing care within hospital units. Journal of Operations Management, 69(7), pp.1124-1152.
Radvinsky, D.S., Yoon, R.S., Schmitt, P.J., Prestigiacomo, C.J., Swan, K.G. and Liporace, F.A., 2012. Evolution and development of the Advanced Trauma Life Support (ATLS) protocol: a historical perspective. Orthopedics, 35(4), pp.305-311.
Ramachandran, H.J., Bin Mahmud, M.S., Rajendran, P., Jiang, Y., Cheng, L. and Wang, W., 2023. Effectiveness of mindfulness-based interventions on psychological well-being, burnout and post-traumatic stress disorder among nurses: A systematic review and meta-analysis. Journal of Clinical Nursing, 32(11-12), pp.2323-2338.
Roberts, N.K., Williams, R.G., Schwind, C.J., Sutyak, J.A., McDowell, C., Griffen, D., Wall, J., Sanfey, H., Chestnut, A., Meier, A.H. and Wohltmann, C., 2014. The impact of brief team communication, leadership and team behavior training on ad hoc team performance in trauma care settings. The American Journal of Surgery, 207(2), pp.170-178.
Rodriguez-Arrastia, M., García-Martín, M., Villegas-Aguilar, E., Ropero-Padilla, C., Martin-Ibañez, L. and Roman, P., 2022. Emotional and psychological implications for healthcare professionals in disasters or mass casualties: a systematic review. Journal of Nursing Management, 30(1), pp.298-309.
Rupp, R., 2021. Spinal Cord Injury. In Neuroprosthetics and Brain-Computer Interfaces in Spinal Cord Injury: A Guide for Clinicians and End Users (pp. 3-35). Cham: Springer International Publishing.
Silva, S. and Whalen, K.C., 2014. Trauma nursing core course for non-emergency nurses. Journal of Emergency Nursing, 40(5), pp.486-487.
Steele, S.E., 2006. When trauma means a stoma. Journal of Wound Ostomy & Continence Nursing, 33(5), pp.491-500.
Steinemann, S., Bhatt, A., Suares, G., Wei, A., Ho, N., Kurosawa, G., Lim, E. and Berg, B., 2016. Trauma team discord and the role of briefing. Journal of Trauma and Acute Care Surgery, 81(1), pp.184-189.
Teman, N.R., Gauger, P.G., Mullan, P.B., Tarpley, J.L. and Minter, R.M., 2014. Entrustment of general surgery residents in the operating room: factors contributing to provision of resident autonomy. Journal of the American College of Surgeons, 219(4), pp.778-787.
Van Veelen, M.J. and Brodmann Maeder, M., 2021. Hypothermia in trauma. International journal of environmental research and public health, 18(16), p.8719.
Van Way 3rd, C., 2016. War and Trauma: A History of Military Medicine-Part II. Missouri Medicine, 113(5), pp.336-340.
Wentzel, D. and Brysiewicz, P., 2014. The consequence of caring too much: compassion fatigue and the trauma nurse. Journal of emergency nursing, 40(1), pp.95-97.
Wolf, L.A., Delao, A.M., Perhats, C., Clark, P.R., Edwards, C. and Frankenberger, W.D., 2020. Traumatic stress in emergency nurses: does your work environment feel like a war zone?. International emergency nursing, 52, p.100895.
Woo, B.F.Y., Lee, J.X.Y. and Tam, W.W.S., 2017. The impact of the advanced practice nursing role on quality of care, clinical outcomes, patient satisfaction, and cost in the emergency and critical care settings: a systematic review. Human resources for health, 15, pp.1-22.
World Health Organization (2021) Injuries and violence. who.int, 19 March (accessed 18 October 2022).
Wurster, L.A., Coffey, C., Haley, K. and Covert, J., 2009. The role of the trauma nurse leader in a pediatric trauma center. Journal of Trauma Nursing| JTN, 16(3), pp.160-165.
Zinan, N., Puia, D. and Kinsley, T., 2015. Results of a mass casualty incident simulation in an undergraduate nursing program. Journal of Nursing Education and Practice, 5(12), p.71.