

December 2017

# Parameter Estimation In Weighted Rayleigh Distribution

M. Ajami

*Vali-e-Asr University of Rafsanjan, Rafsanjan, Iran, m.ajami@vru.ac.ir*

S. M. A. Jahanshahi

*University of Sistan and Baluchestan, Zahedan, Iran, mehdi.jahanshahi@gmail.com*

 Part of the [Applied Statistics Commons](#), [Social and Behavioral Sciences Commons](#), and the [Statistical Theory Commons](#)

## Recommended Citation

Ajami, M., & Jahanshahi, S. M. A. (2017). Parameter Estimation In Weighted Rayleigh Distribution. *Journal of Modern Applied Statistical Methods*, 16(2), 256-276. doi: 10.22237/jmasm/1509495240

# Parameter Estimation In Weighted Rayleigh Distribution

**M. Ajami**

Vali-e-Asr University of Rafsanjan  
Rafsanjan, Iran

**S. M. A. Jahanshahi**

University of Sistan and Baluchestan  
Zahedan, Iran

---

A weighted model based on the Rayleigh distribution is proposed and the statistical and reliability properties of this model are presented. Some non-Bayesian and Bayesian methods are used to estimate the  $\beta$  parameter of proposed model. The Bayes estimators are obtained under the symmetric (squared error) and the asymmetric (linear exponential) loss functions using non-informative and reciprocal gamma priors. The performance of the estimators is assessed on the basis of their biases and relative risks under the two above-mentioned loss functions. A simulation study is constructed to evaluate the ability of considered estimation methods. The suitability of the proposed model for a real data is shown by using the Kolmogorov-Smirnov goodness-of-fit test.

*Keywords:* Bayesian estimators, estimation methods, goodness-of-fit, loss function, reliability, weighted model

---

## Introduction

The Rayleigh distribution has been used in many areas of research, such as reliability, life-testing and survival analysis. Modeling the lifetime of random phenomena has been another area of study for which the Rayleigh distribution has been significantly used. Being first introduced by Rayleigh (1880), this statistical model was originally derived in connection with a problem in acoustics. More details on the Rayleigh distribution can be found in Johnson et al. (1994) and references therein.

The Rayleigh distribution has the following probability density function (pdf) and the cumulative distribution function (cdf), respectively,

---

*M. Ajami is an Assistant Professor in the Department of Statistics. Email him at [m.ajami@vru.ac.ir](mailto:m.ajami@vru.ac.ir). S. M. A. Jahanshahi is a Professor in the Department of Statistics. Email him at [mehdi.jahanshahi@gmail.com](mailto:mehdi.jahanshahi@gmail.com).*

$$f(x) = \frac{x}{\beta} \exp\left(-\frac{x^2}{2\beta}\right), \quad x > 0, \beta > 0,$$

$$F(x) = 1 - \exp\left(-\frac{x^2}{2\beta}\right), \quad x > 0, \beta > 0.$$

Weighted distributions are employed mainly in research associated with reliability, bio-medicine, meta-analysis, econometrics, survival analysis, renewal processes, physics, ecology and branching processes which are found in Patil and Rao (1978), Gupta and Kirmani (1990), Gupta and Keating (1985), Oluyede (1999), Patil and Ord (1976) and Zelen and Feinleib (1969). A weighted form of Rayleigh distribution has been published by Reshi et al. (2014). They introduced a new class of Size-biased Generalized Rayleigh distribution and also investigated the various structural and characterizing properties of that model. In addition, they studied the Bayes estimator of the parameter of the Rayleigh distribution under the Jeffrey's and the extended Jeffrey's priors assuming two different loss functions. They compared four estimation methods by using mean square error through simulation study with varying sample sizes. In fact, weighted distributions arise in practice when observations from a sample are recorded with unequal probabilities

Suppose  $X$  is a non-negative random variable with its unbiased pdf  $f(x, \beta)$ ,  $\beta$  is a parameter, then  $g$  distribution is weighted version of  $f$  and is defined as

$$g(x, \alpha, \beta) = \frac{w(x, \alpha) f(x, \beta)}{E(w(X, \alpha))},$$

where the weight function  $w(x, \alpha)$  is a non-negative function and  $0 < E(w(X, \alpha))$  is a normalizing constant which is  $E(w(X, \alpha)) = \int w(x, \alpha) f(x, \beta) dx$ . Furthermore,  $\alpha$  is a parameter which may or may not depend on  $\beta$  and  $E(w(X, \alpha)) = 1/E_g(1/w(X, \alpha))$  is the harmonic mean of  $w(x, \alpha)$  with the pdf  $g(\cdot)$ .

When  $w(x, \alpha) = x^\alpha$ ,  $\alpha = 0$ , the distribution is referred to as weighted distributions of order  $\alpha$ .

$$g(x, \alpha, \beta) = \frac{x^\alpha f(x, \beta)}{E(X^\alpha)}. \quad (1)$$

For  $\alpha = 1$  or  $2$ , the pdf (1) are referred to as length-biased (size-biased) and area-biased distributions, respectively.

A weighted Rayleigh (WR) distribution is proposed based on (1) and all calculations are done based upon this model, but in the sections of numerical simulations and application to real data a length-biased Rayleigh (LBR) distribution is used without loss of generality. Because determining the value of  $\alpha$  depends on the sampling method so it is not necessary to estimate  $\alpha$  in practice, therefore the focus on estimating the  $\beta$  parameter.

### Weighted Rayleigh distribution

In the following, the  $WR(\alpha, \beta)$  distribution is introduced and then, some properties including the  $r^{\text{th}}$  moment, the corresponding CDF and hazard rate function are calculated.

**Definition 1.** A nonnegative random variable  $X$  is said to have the  $WR(\alpha, \beta)$  distribution provided that the variable's density function is given by

$$g(x, \alpha, \beta) = \frac{1}{2^{\alpha/2} \beta^{\alpha/2+1} \Gamma(\alpha/2+1)} x^{\alpha+1} e^{-x^2/2\beta}, \quad x > 0, \alpha, \beta > 0. \quad (2)$$

**Remark 1.** Suppose that  $X$  follows  $WR(\alpha, \beta)$  and let  $U = X^2/2\beta$ , then  $U$  follows  $\Gamma(\alpha/2+1, 1)$  distribution.

**Remark 2.** The  $WR(\alpha, \beta)$  distribution belongs to the exponential family. Therefore,  $T = \sum_{i=1}^n X_i^2$  is a sufficient complete statistic.

The  $r^{\text{th}}$  moments are useful for inference and model fitting. A result that allows us to compute the moments of the  $WR(\alpha, \beta)$  distribution is given in the following lemma.

**Lemma 1.** If  $X$  be a random variable with density function (2), then the  $r^{\text{th}}$  moment is given by

$$E(X^r) = 2^{r/2} \beta^{r/2} \frac{\Gamma\left(\frac{\alpha+r}{2}+1\right)}{\Gamma\left(\frac{\alpha}{2}+1\right)},$$

where  $r$  is a positive integer.

**Proof.** According to (2)

$$E(X^r) = \int_0^\infty \frac{x^{\alpha+r+1}}{2^{\alpha/2} \beta^{\alpha/2+1} \Gamma(\alpha/2+1)} e^{-x^2/2\beta} dx,$$

let  $x^2/2\beta = u^2$ , then we have

$$E(X^r) = \frac{2\beta(\sqrt{2\beta})^{\alpha+r}}{2^{\alpha/2} \beta^{\alpha/2+1} \Gamma(\alpha/2+1)} \int_0^\infty u^{\alpha+r+1} e^{-u^2} du = 2^{r/2} \beta^{r/2} \frac{\Gamma\left(\frac{\alpha+r}{2}+1\right)}{\Gamma\left(\frac{\alpha}{2}+1\right)}.$$

■

Lemma 1 concludes

$$E(X) = 2^{1/2} \beta^{1/2} \frac{\Gamma\left(\frac{\alpha+1}{2}+1\right)}{\Gamma\left(\frac{\alpha}{2}+1\right)},$$

$$V(X) = 2\beta \left[ \frac{\Gamma\left(\frac{\alpha+2}{2}+1\right)}{\Gamma\left(\frac{\alpha}{2}+1\right)} - \left( \frac{\Gamma\left(\frac{\alpha+1}{2}+1\right)}{\Gamma\left(\frac{\alpha}{2}+1\right)} \right)^2 \right],$$

$$CV(X) = \frac{\sqrt{\Gamma\left(\frac{\alpha}{2}+1\right)\Gamma\left(\frac{\alpha+2}{2}+1\right) - \Gamma^2\left(\frac{\alpha+1}{2}+1\right)}}{\Gamma\left(\frac{\alpha+1}{2}+1\right)}.$$

## PARAMETER ESTIMATION IN WEIGHTED RAYLEIGH DISTRIBUTION

The corresponding CDF of the  $WR(\alpha, \beta)$  distribution is as follows:

$$G(x) = \frac{1}{\Gamma(\alpha/2 + 1)} \int_0^{x^2/2\beta} t^{\alpha/2} e^{-t} dt = \frac{1}{\Gamma(\alpha/2 + 1)} \gamma(\alpha/2 + 1, x^2/2\beta),$$

where  $\gamma(a, z) = \int_0^z t^{a-1} e^{-t} dt$  denotes the lower incomplete gamma function.

In addition, the survival and the hazard rate functions of the  $WR(\alpha, \beta)$  distribution are

$$\bar{G}(x) = \frac{1}{\Gamma(\alpha/2 + 1)} \int_{x^2/2\beta}^{\infty} t^{\alpha/2} e^{-t} dt = \frac{1}{\Gamma(\alpha/2 + 1)} \gamma_1(\alpha/2 + 1, x^2/2\beta),$$

and

$$h(x) = \frac{x^{\alpha+1} e^{-x^2/2\beta}}{2^{\alpha/2} \beta^{\alpha/2+1} \gamma_1(\alpha/2 + 1, x^2/2\beta)},$$

respectively, where  $\gamma_1(a, z) = \int_z^{\infty} t^{a-1} e^{-t} dt$  denotes upper incomplete gamma function.


In special cases, if  $\alpha = 1$ , corresponding length-biased distribution is

$$g(x) = \sqrt{\frac{2}{\pi}} \frac{x^2}{\beta^{3/2}} e^{-x^2/2\beta}, \quad x > 0, \beta > 0,$$


and if  $\alpha = 2$  corresponding area-biased distribution is

$$g(x) = \frac{x^3}{2\beta^2} e^{-x^2/2\beta}, \quad x > 0, \beta > 0.$$

Plots of length-biased and area-biased (ABR) distributions for some parameter values are displayed in [Figure 1](#). Some possible shapes of the LBR and ABR hazard rate functions are displayed in [Figure 2](#)



**Figure 1.** The LBR( $\beta$ ) (left panel) and ABR( $\beta$ ) (right panel) density functions for some parameter values.



**Figure 2.** The LBR( $\beta$ ) (left panel) and ABR( $\beta$ ) (right panel) hazard rate functions for some parameter values.

## Parameter estimation

In this section, the method of moments, the maximum likelihood method, uniformly minimum variance unbiased method, maximum goodness-of-fit method and some Bayesian methods are used to estimate the  $\beta$  parameter of the model.

**Method of moments estimator**

Hereafter, let  $X_1, \dots, X_n$  be a random sample from the  $WR(\alpha, \beta)$  distribution. The method of moments estimator (MME) is

$$\hat{\beta}_{MME} = \frac{\bar{X}^2}{2} \left( \frac{\Gamma\left(\frac{\alpha}{2} + 1\right)}{\Gamma\left(\frac{\alpha + 1}{2} + 1\right)} \right)^2.$$

**Maximum likelihood estimator**

The likelihood function can be written as

$$L(\beta; x_1, K, x_n) = \left( \frac{1}{2^{\alpha/2} \beta^{\alpha/2+1} \Gamma(\alpha/2+1)} \right)^n \prod_{i=1}^n X_i^{\alpha+1} e^{-\frac{\sum_{i=1}^n X_i^2}{2\beta}}, \quad x > 0, \alpha, \beta > 0.$$

One can easily calculate maximum likelihood estimator (MLE) of  $\beta$  by taking natural logarithm and derivative relative to  $\beta$  as

$$\hat{\beta}_{MLE} = \frac{T}{n(\alpha + 2)},$$

where  $T = \sum_{i=1}^n X_i^2$ .

To study asymptotic normality of  $\hat{\beta}_{MLE}$ , calculate the Fisher information  $I(\beta)$  as

$$I(\beta) = \frac{n}{\beta^2} \left( \frac{2\Gamma\left(\frac{\alpha}{2} + 2\right)}{\Gamma\left(\frac{\alpha}{2} + 1\right)} - \left(\frac{\alpha}{2} + 1\right) \right), \tag{3}$$

So according to theorem 18 of Ferguson (1996)

$$\sqrt{I(\beta)}(\hat{\beta}_{MLE} - \beta) \xrightarrow{D} N(0,1) \text{ as } n \rightarrow \infty.$$

Therefore, an  $100(1 - \alpha)\%$  approximate confidence interval of  $\beta$  can be obtained as

$$\hat{\beta}_{MLE} \pm Z_{\alpha/2} \sqrt{I^{-1}(\beta)},$$

where  $Z_{\alpha/2}$  is the  $\alpha/2^{\text{th}}$  percentile point of the standard normal distribution.

### Uniformly minimum variance unbiased estimator

Based upon Lemma 1,

$$E(X^2) = 2\beta \frac{\Gamma\left(\frac{\alpha+2}{2} + 1\right)}{\Gamma\left(\frac{\alpha}{2} + 1\right)}.$$

Therefore,

$$E(T) = 2n\beta \frac{\Gamma\left(\frac{\alpha+2}{2} + 1\right)}{\Gamma\left(\frac{\alpha}{2} + 1\right)},$$

and

$$E\left(\frac{T}{2n} \frac{\Gamma\left(\frac{\alpha}{2} + 1\right)}{\Gamma\left(\frac{\alpha+2}{2} + 1\right)}\right) = \beta,$$

which is a function of the sufficient and complete statistic  $T$  that is unbiased for  $\beta$ . Thus based on Lehman-Scheffe theorem we have

$$\hat{\beta}_{UMVUE} = \frac{T}{2n} \frac{\Gamma\left(\frac{\alpha}{2} + 1\right)}{\Gamma\left(\frac{\alpha + 2}{2} + 1\right)}.$$

### Maximum goodness-of-fit estimators

Maximum goodness-of-fit estimators (otherwise known as minimum distance estimators) of the parameters of the CDF can be calculated by minimizing any distance of the empirical distribution function (EDF) statistics regarding to the unknown parameters. As other research has shown there is no unique EDF statistic which can be considered the most efficient for all situations (Alizadeh and Arghami, 2011). Kolmogorov-Smirnov, Cramer-von Mises and Anderson-Darling statistics seem to be momentous in situations are

$$D_n = \sqrt{n} \sup_{1 \leq i \leq n} |G_n(x_{(i)}) - G(x_{(i)})|,$$

$$W_n^2 = n \sum_{i=1}^n [G_n(x_{(i)}) - G(x_{(i)})]^2 p(x_{(i)}),$$

$$A_n^2 = n \sum_{i=1}^n \frac{[G_n(x_{(i)}) - G(x_{(i)})]^2}{G(x_{(i)})(1 - G(x_{(i)}))} p(x_{(i)}),$$

where  $p(x_{(i)}) = G(x_{(i)}) - G(x_{(i)} - 1)$  is the probability under  $H_0$  and considering that  $G_n(\cdot)$  is EDF for  $G(\cdot)$ .

### Bayes estimators of $\beta$

Considering  $\beta$  as a random variable, two different priors, namely Jeffreys and reciprocal gamma are considered for  $\beta$ . Taking into account the priors, two different loss functions are used for the  $WR(\alpha, \beta)$  model, the first one is the squared error loss (SEL) function and the second one is linear exponential (LINEX) loss function.

### Bayes estimator based on Jeffreys' prior

Based on (3) the Jeffreys' prior is

$$\pi(\beta) \propto \sqrt{I(\beta)} = 1/\beta, \beta > 0,$$

and then, the posterior density will be

$$\pi(\beta | \underline{x}) \propto \frac{1}{\beta^{n(\alpha/2+1)+1}} e^{-\sum_{i=1}^n X_i^2 / 2\beta}, \tag{4}$$

which follows reciprocal gamma distribution as

$$\beta | \underline{x} : \text{rgamma} \left( n \left( \frac{\alpha}{2} + 1 \right), \frac{\sum_{i=1}^n X_i^2}{2} \right).$$

The Bayesian estimator of  $\beta$  under the SEL function is

$$\hat{\beta}_{SEL} = \frac{T}{n(\alpha + 2) - 2},$$

where the SEL function is

$$L(\beta, \hat{\beta}) = (\hat{\beta} - \beta)^2,$$

and  $T = \sum_{i=1}^n X_i^2$ .

In the following, Bayesian estimator is calculated under the LINEX loss function. This loss function was proposed by Varian (1975) and Zellner (1986). The LINEX loss function for scale parameter  $\beta$  is given by

$$L(\Delta) = e^{a\Delta} - a\Delta - 1, a \neq 0, \tag{5}$$

where  $\Delta = \frac{\hat{\beta}}{\beta} - 1$  and  $\hat{\beta}$  is an estimator of  $\beta$ . The sign and magnitude of “a” represent the direction and degree of asymmetry respectively (see Soliman, 2000, and Sanku, 2012). Under LINEX loss function (5) and using the posterior (4), the posterior mean of loss function,  $L(\Delta)$ , is

## PARAMETER ESTIMATION IN WEIGHTED RAYLEIGH DISTRIBUTION

$$E[L(\Delta)] = e^{-a} E\left[e^{a\left(\frac{\hat{\beta}}{\beta}\right)}\right] - aE\left[\left(\frac{\hat{\beta}}{\beta}\right) - 1\right] - 1,$$

one can easily obtain  $\hat{\beta}$  which minimizes the posterior expectation of the loss function (5), denoted by  $\hat{\beta}_{LJ}$  as

$$\hat{\beta}_{LJ} = \frac{T}{2a} \left( 1 - \exp\left(-\frac{a}{n(\alpha/2 + 1) + 1}\right) \right).$$

### Bayes estimator based on reciprocal gamma prior

Suppose  $\beta$  follows reciprocal gamma distribution as prior distribution which is

$$\pi(\beta) = \frac{b^\sigma}{\Gamma(\sigma)} \frac{e^{-b/\beta}}{\beta^{\sigma+1}}, \quad \beta, \sigma > 0.$$

Then, the posterior density satisfies

$$\pi(\beta | \underline{x}) \propto \frac{1}{\beta^{n(\alpha/2+1)+\sigma+1}} e^{-(b+\sum_{i=1}^n X_i^2/2)/\beta},$$

so the Bayesian estimator of  $\beta$  under the SEL function is

$$\hat{\beta}_{SERG} = \frac{T + 2b}{c},$$

where  $c = n(\alpha + 2) + 2\sigma - 2$ .

In special case, if we suppose  $\sigma = 1$ ,  $b = 0$  then Bayesian estimator of  $\beta$  is

$$\hat{\beta}_{SERG} = \frac{T}{n(\alpha + 2)},$$

which is equal to MLE.

In addition, Bayesian estimator of  $\beta$  under the LINEX loss function is

$$\hat{\beta}_{LRG} = d(T + 2b),$$

where  $d = \frac{1 - \exp\left(-\frac{a}{n(\alpha/2+1) + \sigma + 1}\right)}{2a}$ .

**The risk efficiency of  $\hat{\beta}_{SEJ}$  regarding to  $\hat{\beta}_{LJ}$  under LINEX and squared errors loss function based on Jeffreys' prior**

If random variable  $X$  follows the distribution function (2), so  $X^2$  obeys  $\Gamma((\alpha/2+1), 2\beta)$  then  $T : \Gamma(n(\alpha/2+1), 2\beta)$  as

$$h_T(t) = \frac{1}{(2\beta)^{n(\alpha/2+1)} \Gamma(n(\alpha/2+1))} t^{n(\alpha/2+1)-1} e^{-\frac{t}{2\beta}} \quad \alpha, \beta > 0.$$

Because the risk functions of estimators  $\hat{\beta}_{SEJ}$  and  $\hat{\beta}_{LJ}$  are important, calculate these risk functions which are denoted by  $R_L(\hat{\beta}_{LJ})$ ,  $R_L(\hat{\beta}_{SEJ})$ ,  $R_S(\hat{\beta}_{LJ})$ , and  $R_S(\hat{\beta}_{SEJ})$  where the subject  $L$  denotes risk relative LINEX loss function and the subject  $S$  denotes risk relative to SEL.

**Lemma 2.** Let  $X : WR(\alpha, \beta)$ , then risk function of  $\hat{\beta}_{SEJ}$  under LINEX loss function with respect to the Jeffreys' prior is

$$R_L(\hat{\beta}_{SEJ}) = e^{-a} \left(1 - \frac{a}{n(\alpha/2+1) - 1}\right)^{-n(\alpha/2+1)} - \frac{an(\alpha/2+1)}{n(\alpha/2+1) - 1} + a - 1.$$

**Proof.** By definition,

$$\begin{aligned}
 R_L(\hat{\beta}_{SEJ}) &= E[L(\Delta)] = \int_0^\infty \left\{ e^{a\left[\left(\frac{\hat{\beta}_{SEJ}}{\beta}\right)^{-1}\right]} - a\left[\left(\frac{\hat{\beta}_{SEJ}}{\beta}\right)^{-1}\right] - 1 \right\} h(t) dt \\
 &= \int_0^\infty e^{a\left[\left(\frac{\hat{\beta}_{SEJ}}{\beta}\right)^{-1}\right]} h(t) dt - a \int_0^\infty \left(\frac{\hat{\beta}_{SEJ}}{\beta}\right) h(t) dt + a - 1.
 \end{aligned} \tag{6}$$

It is easy to verify

$$\text{(I). } \int_0^\infty e^{a\left[\left(\frac{\hat{\beta}_{SEJ}}{\beta}\right)^{-1}\right]} h(t) dt = e^{-a} \left( 1 - \frac{a}{n(\alpha/2+1)-1} \right)^{-n(\alpha/2+1)},$$

$$\text{(II). } \int_0^\infty \left(\frac{\hat{\beta}_{SEJ}}{\beta}\right) h(t) dt = \frac{n(\alpha/2+1)}{n(\alpha/2+1)-1}.$$

Substituting (I)-(II) into (6), the result desired follows. ■

**Corollary 1.** Based on Lemma 2, one can conclude that

$$R_L(\hat{\beta}_{LJ}) = e^{-\frac{a}{n(\alpha/2+1)+1}} - n(\alpha/2+1) \left( 1 - e^{-\frac{a}{n(\alpha/2+1)+1}} \right) + a - 1.$$

**Lemma 3.** Let  $X : WR(\alpha, \beta)$ , then the risk function of  $\hat{\beta}_{LJ}$  under SEL function with respect to the Jeffreys' prior is

$$R_S(\hat{\beta}_{LJ}) = \frac{\beta^2 \left( 1 - e^{-\frac{a}{n\left(\frac{\alpha}{2}+1\right)}} \right)}{a} \left( \frac{\left( n\left(\frac{\alpha}{2}+1\right) \right) \left( n\left(\frac{\alpha}{2}+1\right) + 1 \right) \left( 1 - e^{-\frac{a}{n\left(\frac{\alpha}{2}+1\right)}} \right)}{a} \right) \left( -n(\alpha+2) + a \left( 1 - e^{-\frac{a}{n\left(\frac{\alpha}{2}+1\right)}} \right)^{-1} \right).$$

**Proof.** By definition,

$$R_S(\hat{\beta}_{LJ}) = \int_0^\infty (\hat{\beta}_{LJ} - \beta)^2 h(t) dt = E(\hat{\beta}_{LJ}^2) - 2\beta E(\hat{\beta}_{LJ}) + \beta^2. \tag{7}$$

However,

$$(I). E(\hat{\beta}_{LJ}^2) = \frac{\beta^2 (n(\alpha/2+1))(n(\alpha/2+1)+1) \left(1 - e^{-\frac{a}{n(\alpha/2+1)}}\right)^2}{a^2}$$

$$(II). 2\beta E(\hat{\beta}_{LJ}) = \frac{\beta^2 n(\alpha+2) \left(1 - e^{-\frac{a}{n(\alpha/2+1)}}\right)}{a}.$$

Substituting (I)-(II) into (7), the proof is completed. ■

**Corollary 2.** In the same procedure of Lemma 3 the  $R_S(\hat{\beta}_{SEJ})$  under the SEL is

$$R_S(\hat{\beta}_{SEJ}) = \beta^2 \left[ \frac{(n(\alpha/2+1))(n(\alpha/2+1)+1)}{(n(\alpha/2+1)-1)^2} - \frac{2n(\alpha/2+1)}{(n(\alpha/2+1)-1)} + 1 \right].$$

**Definition 2.** The risk efficiency of  $\hat{\beta}_2$  regarding to  $\hat{\beta}_1$  under  $L$  loss function is defined as

$$RE_L(\hat{\beta}_1, \hat{\beta}_2) = \frac{R_L(\hat{\beta}_2)}{R_L(\hat{\beta}_1)}.$$

**The risk efficiency of  $\hat{\beta}_{SERG}$  regarding to  $\hat{\beta}_{LRG}$  under LINEX and SEL functions based on reciprocal gamma's prior**

In the following, the risk functions of estimators  $\hat{\beta}_{SERG}$  and  $\hat{\beta}_{LRG}$  are calculated. Therefore, they are denoted by  $R_L(\hat{\beta}_{LRG})$ ,  $R_L(\hat{\beta}_{SERG})$ ,  $R_S(\hat{\beta}_{LRG})$ , and  $R_S(\hat{\beta}_{SERG})$ .

**Corollary 3.** Let  $X: WR(\alpha, \beta)$ , then the risk function of  $\hat{\beta}_{SERG}$  under the LINEX and the SEL functions and reciprocal gamma prior are

$$R_L(\hat{\beta}_{SERG}) = \frac{e^{-a\left(1-\frac{2b}{\beta c}\right)}}{(1-2c)^{n(\alpha/2+1)}} - \frac{a}{\beta c}(2b+n\beta(\alpha+2))+a-1,$$

$$R_L(\hat{\beta}_{SERG}) = \frac{2n\beta^2(\alpha+2)+n^2\beta^2(\alpha+2)^2+4b^2+4nb\beta(\alpha+2)}{c^2} - \frac{2\beta(2b+n\beta(\alpha+2))}{c} + \beta^2.$$

**Corollary 4.** Similar to Corollary 3 under the LINEX and the SEL functions and reciprocal gamma prior we have

$$R_L(\hat{\beta}_{LRG}) = \frac{e^{\frac{b}{\beta}(1-k)-a}}{(2k-1)^{n(\alpha/2+1)}},$$

and

$$R_S(\hat{\beta}_{LRG}) = d^2 \left[ 2n\beta^2(\alpha+2)+n^2\beta^2(\alpha+2)^2+4b^2+4nb\beta(\alpha+2) \right] - 2\beta^2 dn(\alpha+2) - 4bd\beta + \beta^2,$$

where  $k = \exp\left(\frac{-a}{n(\alpha/2+1)+\sigma+1}\right)$ .

### Numerical simulations

In the following, some experimental results are presented to investigate the effectiveness of the different estimation methods which have been so far performed. Bias and MSE for non-Bayesian estimators are mostly compared for different estimation methods. In this study, different sample sizes of  $n = 10, 20$  (small), 30, 40 (moderate), 50 (large) and 100 (very large) are considered. In Table 1, the average estimates of  $\beta$  based on 10,000 replications are presented for different estimation methods in which the MSEs are noted in the parentheses.

As can be seen in Table 1, among simple estimators the MLE and UMVUE have the smallest values of bias and MSE for various values of sample size so MLE and UMVUE are the best estimation methods in terms of bias and MSE. In addition, the other two good methods of estimation in priority of order are MME and CVM.

**Table 1.** Bias and MSE values of simple estimators for  $\beta$  parameter

$n$	MLE	MME	UMVUE	KS	CVM	AD
10	<b>-0.002550</b>	0.014100	<b>-0.002550</b>	0.024430	0.023930	0.030840
	<b>(0.000007)</b>	(0.000995)	<b>(0.000007)</b>	(0.000601)	(0.000572)	(0.000952)
20	<b>-0.002770</b>	0.006340	<b>-0.002770</b>	0.012050	0.011170	0.014600
	<b>(0.000006)</b>	(0.000040)	<b>(0.000006)</b>	(0.000145)	(0.000125)	(0.000213)
30	<b>-0.000260</b>	0.005690	<b>-0.000260</b>	0.009430	0.009170	0.011480
	<b>(0.000003)</b>	(0.000032)	<b>(0.000003)</b>	(0.000088)	(0.000084)	(0.000131)
40	<b>-0.001070</b>	0.003430	<b>-0.001070</b>	0.006660	0.005620	0.007670
	<b>(0.000002)</b>	(0.000012)	<b>(0.000002)</b>	(0.000044)	(0.000031)	(0.000058)
50	<b>-0.003730</b>	-0.000380	<b>-0.003730</b>	0.001920	0.001050	0.002810
	<b>(0.000001)</b>	(0.000000)	<b>(0.000001)</b>	(0.000000)	(0.000000)	(0.000000)
100	<b>0.000550</b>	0.002840	<b>0.000550</b>	0.003610	0.003110	0.004190
	<b>(0.000000)</b>	(0.000000)	<b>(0.000000)</b>	(0.000001)	(0.000001)	(0.000001)

Bias values and risk functions are computed to compare considered Bayesian estimators. A comparison of this type is needed to check whether an estimator is inadmissible under some loss function. Therefore, if it is so, the estimator would not be used for the losses specified by that loss function. For this purpose, the risks of the estimators and the efficiency of them are computed. In each case,  $a = 1, a = -1, b = 2$  and  $\sigma = 2$  are taken without loss of generality.

Because comparing different loss functions is not reasonable, compare the results in similar loss function, but in different priors. According to results compiled in Tables 2, 3, 5 and 6, all the four considered Bayesian estimators

## PARAMETER ESTIMATION IN WEIGHTED RAYLEIGH DISTRIBUTION

based on reciprocal gamma prior have small values of bias. Further, the  $\hat{\beta}_{SERG}$  estimator has smaller bias than  $\hat{\beta}_{LRG}$  estimator for  $a = 1$  while  $\hat{\beta}_{LRG}$  estimator has smaller bias than  $\hat{\beta}_{SERG}$  estimator for  $a = -1$ .

According to Tables 2, 3, 5 and 6, among the four considered Bayesian risks based SEL the  $R_S(\hat{\beta}_{LRG})$  has the smallest values of risk for various values of sample size.

Also among the four considered Bayesian risks based LINEX, the  $R_L(\hat{\beta}_{LJ})$  has the smallest values of risk for various values of sample size.

**Table 2.** Bias and risk values of Bayesian estimators for  $\beta$  parameter and  $a = 1$

$n$	$bias(\hat{\beta}_{SEJ})$	$bias(\hat{\beta}_{LJ})$	$R_S(\hat{\beta}_{SEJ})$	$R_S(\hat{\beta}_{LJ})$	$R_L(\hat{\beta}_{SEJ})$	$R_L(\hat{\beta}_{LJ})$
10	<b>0.072</b>	-0.090	0.100	0.056	0.047	<b>0.031</b>
20	<b>0.035</b>	-0.047	0.041	0.031	0.020	<b>0.016</b>
30	<b>0.024</b>	-0.031	0.025	0.021	0.012	<b>0.011</b>
40	<b>0.016</b>	-0.025	0.018	0.016	0.009	<b>0.008</b>
50	<b>0.011</b>	-0.022	0.014	0.013	0.007	<b>0.007</b>
100	<b>0.005</b>	-0.011	0.007	0.007	0.003	<b>0.003</b>

**Table 3.** Bias and risk values of Bayesian estimators for  $\beta$  parameter and  $a = -1$

$n$	$bias(\hat{\beta}_{SEJ})$	$bias(\hat{\beta}_{LJ})$	$R_S(\hat{\beta}_{SEJ})$	$R_S(\hat{\beta}_{LJ})$	$R_L(\hat{\beta}_{SEJ})$	$R_L(\hat{\beta}_{LJ})$
10	0.065	<b>-0.039</b>	0.099	0.071	0.037	<b>0.032</b>
20	0.034	<b>-0.017</b>	0.041	0.035	0.018	<b>0.016</b>
30	0.022	<b>-0.011</b>	0.025	0.023	0.012	<b>0.011</b>
40	0.018	<b>-0.007</b>	0.018	0.017	0.009	<b>0.008</b>
50	0.013	<b>-0.007</b>	0.014	0.014	0.007	<b>0.007</b>
100	0.007	<b>-0.003</b>	0.007	0.007	0.003	<b>0.003</b>

**Table 4.** Relative risk values of Bayesian estimators for  $\beta$  parameter

$n$	10	20	30	40	50	100
$RE_{S(a=1)}(\hat{\beta}_{LJ}, \hat{\beta}_{SEJ})$	1.524	1.233	1.149	1.110	1.087	1.043
$RE_{L(a=1)}(\hat{\beta}_{LJ}, \hat{\beta}_{SEJ})$	1.788	1.338	1.214	1.157	1.124	1.060
$RE_{S(a=-1)}(\hat{\beta}_{LJ}, \hat{\beta}_{SEJ})$	1.164	1.078	1.051	1.038	1.031	1.015
$RE_{L(a=-1)}(\hat{\beta}_{LJ}, \hat{\beta}_{SEJ})$	1.382	1.174	1.112	1.083	1.066	1.032

**Table 5.** Bias and risk values of Bayesian estimators for  $\beta$  parameter and  $a =$

$n$	$bias(\hat{\beta}_{SERG})$	$bias(\hat{\beta}_{LRG})$	$R_s(\hat{\beta}_{SERG})$	$R_s(\hat{\beta}_{LRG})$	$R_L(\hat{\beta}_{SERG})$	$R_L(\hat{\beta}_{LRG})$
10	<b>0.061</b>	-0.083	0.073	<b>0.045</b>	0.504	2.317
20	<b>0.031</b>	-0.046	0.035	<b>0.027</b>	0.501	2.487
30	<b>0.022</b>	-0.031	0.023	<b>0.019</b>	0.500	2.556
40	<b>0.015</b>	-0.025	0.017	<b>0.015</b>	0.500	2.593
50	<b>0.011</b>	-0.021	0.014	<b>0.012</b>	0.500	2.617
100	<b>0.007</b>	-0.009	0.007	<b>0.006</b>	0.500	2.666

**Table 6.** Bias and risk values of Bayesian estimators for  $\beta$  parameter and  $a = -1$

$n$	$bias(\hat{\beta}_{SERG})$	$bias(\hat{\beta}_{LRG})$	$R_s(\hat{\beta}_{SERG})$	$R_s(\hat{\beta}_{LRG})$	$R_L(\hat{\beta}_{SERG})$	$R_L(\hat{\beta}_{LRG})$
10	0.060	<b>-0.031</b>	0.073	<b>0.050</b>	4.916	0.474
20	0.034	<b>-0.014</b>	0.035	<b>0.029</b>	5.365	0.425
30	0.022	<b>-0.011</b>	0.023	<b>0.020</b>	5.531	0.407
40	0.015	<b>-0.009</b>	0.017	<b>0.015</b>	5.616	0.397
50	0.014	<b>-0.006</b>	0.014	<b>0.013</b>	5.670	0.392
100	0.007	<b>-0.003</b>	0.007	<b>0.006</b>	5.778	0.380

**Table 7.** Relative risk values of Bayesian estimators for  $\beta$  parameter

$n$	10	20	30	40	50	100
$RE_{S(a=1)}(\hat{\beta}_{LRG}, \hat{\beta}_{SERG})$	1.743	1.331	1.211	1.156	1.123	1.060
$RE_{L(a=1)}(\hat{\beta}_{LRG}, \hat{\beta}_{SERG})$	0.218	0.201	0.196	0.193	0.192	0.188
$RE_{S(a=-1)}(\hat{\beta}_{LRG}, \hat{\beta}_{SERG})$	1.511	1.241	1.157	1.117	1.093	1.046
$RE_{L(a=-1)}(\hat{\beta}_{LRG}, \hat{\beta}_{SERG})$	10.372	12.625	13.593	14.131	14.474	15.205

### Application to real data

Here, in order to display the usage of proposed model in real data, it is needed to analyze two sets of the seven from the afore presented data in paper by Bennett and Filliben (2000). Reportedly, they have notified minority electron mobility for p-type  $Ga_{1-x}Al_xAs$  with seven different values of mole fraction. To do so, two data sets are employed relating to the mole fractions of 0.25 and 0.30. The data values are as followed:

## PARAMETER ESTIMATION IN WEIGHTED RAYLEIGH DISTRIBUTION

Data Set 1 (belongs to mole fraction 0.25): 3.051, 2.779, 2.604, 2.371, 2.214, 2.045, 1.715, 1.525, 1.296, 1.154, 1.016, 0.7948, 0.7007, 0.6292, 0.6175, 0.6449, 0.8881, 1.115, 1.397, 1.506, 1.528.

Data Set 2 (belongs to mole fraction 0.30): 2.658, 2.434, 2.288, 2.092, 1.959, 1.814, 1.530, 1.366, 1.165, 1.041, 0.9198, 0.7241, 0.6403, 0.576, 0.5647, 0.5873, 0.8013, 1.002, 1.250, 1.347, 1.368.

To evaluate the fitting quality of the Rayleigh and LBR distributions, the Kolmogorov-Smirnov (K-S) tests and AIC and BIC's criterions are used. The information about comparing both models are given in Table 8. Since probability values of the LBR model are greater than corresponding values of the Rayleigh model and the AIC and BIC criterions of the LBR model are less than corresponding values of the Rayleigh model. Although the values of considered statistics are not significantly different but we it can be infered that the LBR distribution fits better than the Rayleigh distribution in both considered data.

The MLEs of  $\beta$  are 0.9322 and 0.7309 and the 95 percent confidence intervals of  $\beta$  based on MLEs as suggested above under heading [Parameter Estimation](#), can be obtained as (0.6067,1.2577) and (0.4757,0.9861) respectively.

**Table 8.** Comparing related statistics for Rayleigh and LBR

Data	Model	D	p.value	AIC	BIC
1	Rayleigh	0.1411	0.7458	46.0090	47.0540
1	LBR	<b>0.1275</b>	<b>0.8427</b>	<b>45.9160</b>	<b>46.9610</b>
2	Rayleigh	0.1354	0.7883	40.3870	41.4320
2	LBR	<b>0.1311</b>	<b>0.8180</b>	<b>39.7820</b>	<b>40.8260</b>

## Conclusion

Different estimation procedures were studied for estimating the unknown scale parameter of the  $WR(\alpha,\beta)$  distribution being the maximum likelihood estimator, the method of moment estimator, uniformly minimum variance unbiased estimator, maximum goodness-of-fit estimators and the Bayes estimators. Since it is not possible to compare different methods theoretically, some simulations were used for comparison of different estimators with respect to biases, mean squared errors and risks.

All the four considered Bayesian estimators based on reciprocal gamma prior have small values of bias. In addition, the  $\hat{\beta}_{SERG}$  estimator has smaller bias

than  $\hat{\beta}_{LRG}$  estimator for  $a = 1$  but  $\hat{\beta}_{LRG}$  estimator has smaller bias than  $\hat{\beta}_{SERG}$  estimator for  $a = -1$ .

Among the four considered Bayesian risks based SEL the  $R_S(\hat{\beta}_{LRG})$  has the smallest values of risk and based LINEX, the  $R_L(\hat{\beta}_{LJ})$  has the smallest values of risk for various values of sample size. Thus from a Bayesian perspective we suggest using  $\hat{\beta}_{LRG}$  estimator based on SEL and using  $\hat{\beta}_{LJ}$  based on LINEX loss function.

The performance of the MLE and UMVUE is also quite satisfactory and in overall non-Bayesian estimators are better than Bayesian estimators, thereby employing of the MLE and UMVUE estimators can be recommend for all practical purposes.

## Acknowledgements

Portions of this paper are developed from the authors' earlier conference presentation (Ajami & Jahanshahi, 2016).

## References

- Ajami, M. & Jahanshahi, S. M. A. (2016, August 24-26). Comparison of various estimation methods for size-biased Rayleigh distribution. Paper presented at the 13<sup>th</sup> Iranian Statistical Conference, Shahid Bahonar University of Kerman, Iran.
- Alizadeh, N. H. & Arghami, N. R. (2011). Monte Carlo comparison of five exponentiality tests using different entropy estimates. *Journal of Statistical Computation and Simulation*, 81(11), 1579–1592. doi: 10.1080/00949655.2010.496368
- Bennett, H. S. & Filliben, J. J. A. (2000). A systematic approach for multidimensional, closed form analytic modeling: minority electron mobilities in  $Ga_{1-x}Al_xAs$  heterostructures, *Journal of Research of the National Institute of Standards and Technology*, 105(3), 441–452. doi: 10.6028/jres.105.037
- Ferguson, T. S. (1996). *A Course In Large Sample Theory*. New York: Chapman and Hall.
- Gupta, R. C. & Keating, J. P. (1985). Relations for reliability measures under length biased sampling, *Scandinavian Journal of Statistics*, 13, 49–56.

## PARAMETER ESTIMATION IN WEIGHTED RAYLEIGH DISTRIBUTION

- Gupta, R. C. & Kirmani, S. N. U. A. (1990). The role of weighted distributions in stochastic modeling, *Communications in Statistics - Theory and Methods*, 19(9), 3147–3162. doi: 10.1080/03610929008830371
- Johnson N. L., Kotz, S., & Balakrishnan, N. (1994). *Continuous univariate distributions, Vol 1* (2<sup>nd</sup> Ed.). New York: Wiley.
- Oluyede, B. O. (1999). On inequalities and selection of experiments for length-Biased Distributions. *Probability in the Engineering and Informational Sciences*, 13(2), 169–185. doi: 10.1017/s0269964899132030
- Patil, G. P. & Rao, C. R. (1978). Weighted distributions and size-biased sampling with applications to wildlife populations and human families. *Biometrics*, 34(2), 179–184. doi: 10.2307/2530008
- Patil, G. P. & Ord, J. K. (1976). On size-biased sampling and related form-invariant weighted distribution. *Sankhya: The Indian Journal Of Statistics B*, 38, 48–61.
- Rayleigh, J. W. S. (1880). On the resultant of a large number of vibrations of the some pitch and of arbitrary phase, *Philosophical Magazine, 5th Series*, 10(60), 73–78. doi: 10.1080/14786448008626893
- Reshi, J. A., Ahmed, A. & Mir, K. A. (2014). Characterizations and estimation in the length-biased generalized Rayleigh distribution. *Mathematical Theory and Modeling*, 4(6), 87-98.
- Sanku, D. (2012). Bayesian estimation of the parameter and reliability function of an inverse Rayleigh distribution, *Malaysian Journal of Mathematical Sciences*, 6(1), 113–124.
- Soliman, A. A. (2000). Comparison of LINEX and quadratic Bayes estimators for the Rayleigh distribution. *Communications in Statistics - Theory and Methods*, 29(1), 95–107. doi: 10.1080/03610920008832471
- Varian, M. M. (1975). A Bayesian approach to real estate assessment. In S. E. Fienberg and A. Zellner, Ed. *Studies in Bayesian Econometrics and Statistics (In Honor of Leonard J. Savage)*, pp. 195–208. Amsterdam: North Holland/Elsevier.
- Zelen, M. & Feinleib, M. (1969). On the theory of chronic diseases, *Biometrika*, 56(3), 601–614. doi: 10.2307/2334668
- Zellner, A. (1986). Bayesian estimation and prediction using loss functions, *Journal of the American Statistical Association*. 81(394), 446–451. doi: 10.2307/2289234