

ASYMPTOTIC SPREADING OF PREDATOR-PREY POPULATIONS IN A SHIFTING ENVIRONMENT

KING-YEUNG LAM AND RAY LEE

ABSTRACT. Inspired by a recent study associating shifting temperature conditions with changes in the efficiency with which predators convert prey to offspring, we propose a predator-prey model of reaction-diffusion type to analyze the consequence of such effects on the population dynamics and spread of the predator species. In the model, the predator conversion efficiency is represented by a spatially heterogeneous function depending on the variable $\xi = x - c_1 t$ for some given $c_1 > 0$. Using the Hamilton-Jacobi approach, we provide explicit formulas for the spreading speed of the predator species. When the conversion function is monotone increasing, the spreading speed is determined in all cases and non-local pulling is possible. When the function is monotone decreasing, we provide formulas for the spreading speed when the rate of shift of the conversion function is sufficiently fast or slow.

1. INTRODUCTION

A fundamental challenge in ecology is to understand the effect of climate change on the population abundances and distributions of species. Climate-induced changes in species abundance and distribution have significant implications for biodiversity [61, 48, 59], the functioning of ecosystems [59, 60, 66], the spread of disease [54, 69] and human welfare [54]. While many species are vulnerable to a changing climate, for many others climate change may facilitate expansion to new areas and population growth [62, 4, 11, 34]. These outcomes arise not only from isolated effects of climate on individual species, but also from changes to interactions between species [30].

Mathematical modeling can be used to determine why certain species decline while others prosper under the changing climate. The study of species persistence and spread often depends on the spatial context, and much analysis in the classical literature has been based on reaction-diffusion models. A prominent example is the Fisher-KPP equation [27, 39], which describes the spreading of a single population

$$u_t = du_{xx} + f(u) \quad \text{for } x \in \mathbb{R}, t > 0, \quad (1.1)$$

where d corresponds to the dispersal rate, $f(u) = ru(1 - u)$, and r is the intrinsic growth rate of the species u .

The problem of spreading speed for (1.1) was first investigated by Kolmogorov et al. [39] for heaviside initial condition, who showed under monostable assumption on $f(u)$ (i.e. $f(s)/s$ is decreasing in s) that

$$c^* = 2\sqrt{df'(0)}.$$

Received by the editors 18 March 2024; accepted 19 June 2024; published online 22 July 2024.

2010 *Mathematics Subject Classification*. Primary 35K58, 35B40; Secondary 35D40.

Key words and phrases. Hamilton-Jacobi equations, viscosity solution, spreading speed, non-cooperative system, predator-prey, reaction-diffusion equations.

K.-Y. Lam and R. Lee were supported in part by NSF Grant DMS-2325195.

Here, c^* refers to the key notion of spreading speed introduced by Aronson and Weinberger [2, 3], which refers to the number $c^* > 0$ such that

$$\lim_{t \rightarrow \infty} \sup_{|x| > ct} u(t, x) = 0 \quad \text{for } c \in (c^*, \infty), \quad \text{and} \quad \lim_{t \rightarrow \infty} \inf_{|x| > ct} u(t, x) > 0 \quad \text{for } c \in (0, c^*).$$

The above result was also generalized in [3] to any compactly supported initial data and in higher spatial dimensions. This theoretical spreading speed has yielded good estimates for range expansion observed in nature [57].

Various studies have since revisited Fisher's model with an interest in the impact of a shifting environment. To consider climate change, it is often assumed that the behavior of the species depends on the variable $\xi = x - ct$, where the constant c corresponds to the velocity of a shifting climate [55, 8, 45, 9]. See also [26], in which this formulation is applied in the context of an SIS model, and [64] for a survey on reaction-diffusion models in shifting environments. Many models have proposed the case where the growth rate $ru(1 - u)$ in (1.1) is replaced by a shifting logistic form $u(r(x - ct) - u)$, where $r(x - ct)$ denotes the species' intrinsic growth rate [55, 8]. These works assumed the growth rate $r(\xi)$ to be positive on a bounded patch of suitable habitat and negative elsewhere, and were broadly interested in the effects of a shifting climate on the persistence of the species.

A shifting environment also leads to new spreading phenomena. In [41], the spreading speed for solutions of Fisher's equation with growth rate $f(u, x - ct) = (r(x - ct) - u)u$ was determined using the Hamilton-Jacobi method, in the case that the intrinsic growth rate of the species is positive and monotone. They showed that, for a certain range of velocities of climate shift, the species spreads with speed distinct from either of the limiting KPP invasion speeds in a phenomenon called non-local pulling [35, 31]. When the growth rate is non-monotone, the existence of forced waves and their attractivity is studied in [9].

In addition to single species equations, the spreading dynamics for systems of equations has received considerable attention. Building on the earlier works on order-preserving systems (such as cooperative systems and competitive systems of two species) [44, 65, 46, 47], the spreading of two competing species in a shifting habitat is studied in [71, 18].

By contrast, for predator-prey systems a comparison principle is not immediately available and many studies regarding propagation phenomena in these systems have focused on the dynamics of traveling wave solutions. The existence of traveling wave solutions for two-species predator prey equations was established in [24, 29], and studied further in [36, 51], while some results on the stability of traveling wave solutions were established in [28].

Until recently, few works have treated the spreading dynamics of predator-prey systems with general initial data. In [53], Pan determined the spreading speed of the predator for a predator-prey system with initially constant prey density and compactly-supported predator. Shortly thereafter, Ducrot, Giletti, and Matano [22] used methods from uniform persistence theory to characterize the spreading dynamics when both predator and prey are initially compactly supported. They showed that the behavior can be classified based on the speeds of the prey in the absence of predator, and of the predator when prey is abundant (see also [23], for the case of a predator-prey equation with non-local dispersal, and [19, 52]). Since these works, the spreading speeds regarding the Cauchy problem for predator-prey systems with three species was studied in [21] (see also [67]). There, it was shown that the nonlocal pulling phenomenon can occur in a system with two predators and one prey.

For other types of non-cooperative systems and their spreading speeds, we refer to [63], which characterizes the spreading speed for a general class of non-cooperative reaction diffusion systems as the minimal traveling wave speed. We also note [20], which determined the spreading speed of infectious disease in an epidemic model, and [50], which considered the spreading dynamics for competition between

three species. Spreading dynamics are also studied for nonlocal diffusion problems, here we mention [70, 68] for such results in predator-prey models in the absence of shifting environment.

1.1. The predator-prey model in a shifting environment. We are interested in the effect of the heterogeneous shifting profile of the conversion efficiency of prey to predator, represented below by the function $\tilde{a}(x - c_1t)$, on the spreading dynamics. Though the temperature-dependence of the conversion efficiency is not well-understood, there is some evidence that the conversion efficiency may be impacted by climate. Using an experimental system of predator and prey, Daugaard et al. [16] found that the conversion efficiency of the predator increased with warming. Moreover, while conversion efficiency was not directly considered, a meta-analysis by Lang et. al [42] identified a trend toward increasing efficiency of *energy assimilation* by consumers with increasing temperature. On the other hand, many biological processes depend unimodally on temperature, such that measures of species performance and fitness decline once temperature increases sufficiently beyond a “thermal optimum” [37, 17, 13]. It is thus plausible that predators currently experiencing climates at or near their thermal optima may experience declines in conversion efficiency with additional warming.

To this end, we propose the following predator-prey model of reaction-diffusion type to analyze the consequence of such effects on the population dynamics and spread of species:

$$\begin{cases} \tilde{u}_t = d_1\tilde{u}_{xx} + \tilde{u}(-\kappa - \alpha_1\tilde{u} + \tilde{a}(x - c_1t)\tilde{v}) & \text{in } (0, \infty) \times \mathbb{R}, \\ \tilde{v}_t = d_2\tilde{v}_{xx} + \tilde{v}(\tilde{r} - \alpha_2\tilde{v} - \tilde{b}\tilde{u}) & \text{in } (0, \infty) \times \mathbb{R}, \\ \tilde{u}(0, x) = \tilde{u}_0(x), \quad \tilde{v}(0, x) = \tilde{v}_0(x) & \text{in } \mathbb{R}. \end{cases} \quad (1.2)$$

Here the predator and prey densities are represented by $\tilde{u}(t, x)$ and $\tilde{v}(t, x)$. It is assumed that the predator cannot persist in the absence of prey, and competes with other predators, while in the absence of predation the prey exhibits logistic growth and is described by the standard Fisher-KPP equation. The interaction rates between predator and prey are mediated by the consumption rate $\tilde{b} > 0$ of prey by predator, and by the predator’s conversion efficiency function $\tilde{a}(x - c_1t)$, which describes the degree to which consumed prey can be successfully converted to additional predators. For simplicity, we assume the conversion efficiency has a fixed profile in the moving coordinate $y = x - c_1t$ with constant velocity c_1 . Finally, $d_i, \alpha_i, \kappa, \tilde{r}$ are positive parameters, where d_i are the random dispersal rates, α_i are the intraspecific competition rates, κ is the natural death rate of the predator species and \tilde{r} is the natural birth rate of the prey species.

Without loss of generality, we may non-dimensionalize the problem (1.2) and obtain the following model:

$$\begin{cases} u_t = u_{xx} + (-1 - u + a(x - c_1t)v)u & \text{in } (0, \infty) \times \mathbb{R} \\ v_t = dv_{xx} + r(1 - v - bu)v & \text{in } (0, \infty) \times \mathbb{R} \\ u(0, x) = u_0(x), \quad v(0, x) = v_0(x) & \text{in } \mathbb{R}. \end{cases} \quad (1.3)$$

We assume the following throughout our study of the reaction-diffusion system (1.3):

(H1) The function $a : \mathbb{R} \rightarrow \mathbb{R}$ is monotone, and satisfies

$$\beta := 1 - b(\|a\|_\infty - 1) > 0, \quad \inf_{s \in \mathbb{R}} a(s) > \frac{1}{\beta}, \quad \text{and} \quad \|a\|_\infty > 1.$$

By observing (via **(IC)** below and maximum principle) that the density of the predator is bounded from above by $\|a\|_\infty - 1$, it follows that the quantity $\beta := 1 - b(\|a\|_\infty - 1)$ corresponds to the minimum carrying capacity for the prey.

Remark 1.1. It is documented in a microbial predator-prey system [16] that the quantity of predators produced for each prey consumed increases when temperature is increased. This corresponds to the

case when $c_1 > 0$ and $a(\cdot)$ is decreasing. We also study the case when increasing temperature decreases the predator efficiency, i.e. $a(\cdot)$ is increasing.

We are interested in the situation when the initial data of the predator is compactly supported, while that of the prey has a positive upper and lower bound. For simplicity, we will assume throughout the discussion that $(u_0, v_0) \in C^2(\mathbb{R})$ satisfies

(IC) $0 \leq u_0 \leq \|a\|_\infty - 1$, $\beta \leq v_0 \leq 1$, and u_0 has compact support.

Finally, we define the following limiting growth rates (at $\pm\infty$), to simplify the statements and proofs of the main results.

$$\begin{cases} r_1 = a(-\infty) - 1, & r_2 = a(+\infty) - 1, \\ \underline{r}_1 = \beta a(-\infty) - 1, & \underline{r}_2 = \beta a(+\infty) - 1. \end{cases} \tag{1.4}$$

Here, r_1 and r_2 correspond to the limiting growth rates of the predator behind and ahead of the environmental shift, respectively, when the prey density is at its minimum value $v = \beta$, while \underline{r}_1 and \underline{r}_2 are the limiting growth rates of the predator behind and ahead of the shift, respectively, when the prey density is at its maximum value $v = 1$.

1.2. Main Results. In this paper, we are interested in the asymptotic speed of spread (or spreading speed) as the predator species u expands its territory. Up to a change of coordinates $x \mapsto -x$, it is enough to focus our discussion on the rightward spreading speed, while allowing the spatial heterogeneity $a(\cdot)$ to be monotonically increasing or decreasing.

In the remainder of this paper, we will refer to the rightward spreading speed c^* simply as the spreading speed, which is defined as follows:

Definition 1.1. Suppose a species has population density $u(t, x)$. We say that the species u has spreading speed given by $c^* > 0$ if

$$\lim_{t \rightarrow \infty} \sup_{x > ct} u(t, x) = 0 \quad \text{for each } c \in (c^*, \infty), \tag{1.5}$$

$$\lim_{t \rightarrow \infty} \inf_{0 < x < ct} u(t, x) > 0 \quad \text{for each } c \in (0, c^*). \tag{1.6}$$

Following [33, Definition 1.2], we introduce the notion of maximal and minimal speed:

Definition 1.2. Suppose a species has population density $u(t, x)$, then its maximal speed \bar{c}_* and minimal speed \underline{c}_* are given by

$$\begin{cases} \bar{c}_* := \inf \left\{ c > 0 \mid \limsup_{t \rightarrow \infty} \sup_{x > ct} u(t, x) = 0 \right\}, \\ \underline{c}_* := \sup \left\{ c > 0 \mid \liminf_{t \rightarrow \infty} \inf_{0 \leq x < ct} u(t, x) > 0 \right\}. \end{cases} \tag{1.7}$$

Remark 1.2. It follows that the species u has a spreading speed if and only if $\bar{c}_* = \underline{c}_*$. In such a case, the spreading speed c^* is given by the common value $\bar{c}_* = \underline{c}_*$.

The following two main theorems characterize the spreading speed of u for the cases (i) $a(\cdot)$ is monotonically increasing and (ii) $a(\cdot)$ is monotonically decreasing, respectively. Assuming the positive axis points poleward and temperature is rising, they correspond to the cases when the the conversion efficiency of the predator is suppressed or enhanced by the warming climate.

Theorem 1.1. *Let $c_1 > 0$ be given, $a : \mathbb{R} \rightarrow \mathbb{R}$ be increasing, and suppose **(H1)** holds. If $(u(t, x), v(t, x))$ is the solution of (1.3) with initial data satisfying **(IC)**, then the spreading speed of u exists, and is given by*

$$c^* := \begin{cases} 2\sqrt{r_2} & \text{if } c_1 \leq 2\sqrt{r_2} \\ \frac{c_1}{2} - \sqrt{r_2 - r_1} + \frac{r_1}{\frac{c_1}{2} - \sqrt{r_2 - r_1}} & \text{if } 2\sqrt{r_2} < c_1 < 2\sqrt{r_1} + 2\sqrt{r_2 - r_1} \\ 2\sqrt{r_1} & \text{if } c_1 \geq 2\sqrt{r_1} + 2\sqrt{r_2 - r_1}. \end{cases} \quad (1.8)$$

Theorem 1.2. *Let $c_1 \in \mathbb{R} \setminus (2\sqrt{r_2}, 2\sqrt{r_1}]$ be given, $a : \mathbb{R} \rightarrow \mathbb{R}$ be decreasing, and suppose **(H1)** holds. If $(u(t, x), v(t, x))$ is the solution of (1.3) with initial data satisfying **(IC)**, then the spreading speed of u exists, and is given by*

$$c^* = \begin{cases} 2\sqrt{r_2} & \text{if } c_1 \leq 2\sqrt{r_2}, \\ 2\sqrt{r_1} & \text{if } c_1 > 2\sqrt{r_1}. \end{cases}$$

For the proof of Theorems 1.1 and 1.2, see Subsection 5.2

We also show the convergence of (u, v) to the homogeneous state in the moving frames with speed different from c_1 or c^* .

Theorem 1.3. *Suppose the spreading speed of u exists and is denoted by $c^* > 0$. Define, for each $c \in (0, \infty) \setminus \{c_1, c^*\}$, the translation of (u, v)*

$$(u_c(t, x), v_c(t, x)) = (u(t, x + ct), v(t, x + ct)).$$

(a) *If $c \in (c^*, \infty)$, then $\lim_{t \rightarrow \infty} (u_c(\cdot, t), v_c(\cdot, t)) \rightarrow (0, 1)$ in $C_{loc}(\mathbb{R})$.*

(b) *If $c \in (0, c^*) \setminus \{c_1\}$, then*

$$\lim_{t \rightarrow \infty} (u_c(\cdot, t), v_c(\cdot, t)) \rightarrow (\tilde{u}, \tilde{v}) \quad \text{in } C_{loc}(\mathbb{R}),$$

where the constant vector $(\tilde{u}, \tilde{v}) \in \mathbb{R}^2$ is the unique positive equilibrium of the kinetic system

$$\frac{d}{dt} \tilde{u} = \tilde{u}(-1 - \tilde{u} + A\tilde{v}), \quad \frac{d}{dt} \tilde{v} = r\tilde{v}(1 - \tilde{v} - b\tilde{u}),$$

such that $A = a(-\infty)$ if $c < c_1$ and $A = a(+\infty)$ if $c > c_1$.

For the proof of Theorem 1.3, see Section 6.

Remark 1.3. Note that the case $c_1 \in (2\sqrt{r_2}, 2\sqrt{r_1}]$ is not covered by Theorem 1.2. In that case the Hamilton-Jacobi approach that we adopt in this paper does not directly apply. We conjecture that $c^* = c_1$ in that case and the predator advances in locked step with the environment. See [9, 26] for results regarding a single species in a shifting habitat. A possible approach is to use the persistence theory as in [21].

Remark 1.4. In the case of $a(\cdot) \equiv a_0$ being a constant and $v_0 \equiv 1$, the spreading speed of the predator u was determined by Pan in [53, Theorem 2.1].

To consolidate the formulas for the spreading speeds in Theorems 1.1 and 1.2, we will denote $\lambda^* = \frac{c_1}{2} - \sqrt{|r_2 - r_1|}$. Then the spreading speed for all cases can be given by

$$\sigma(c_1; r_1, r_2) = \begin{cases} 2\sqrt{r_2} & \text{if } r_1 < r_2 \quad \text{and} \quad c_1 \leq 2\sqrt{r_2} \\ \lambda^* + \frac{r_1}{\lambda^*} & \text{if } r_1 < r_2 \quad \text{and} \quad 2\sqrt{r_2} < c_1 < 2\sqrt{r_1} + 2\sqrt{r_2 - r_1} \\ 2\sqrt{r_1} & \text{if } r_1 < r_2 \quad \text{and} \quad c_1 \geq 2\sqrt{r_1} + 2\sqrt{r_2 - r_1}, \\ 2\sqrt{r_2} & \text{if } r_1 > r_2 \quad \text{and} \quad c_1 \leq 2\sqrt{r_2} \\ 2\sqrt{r_1} & \text{if } r_1 > r_2 \quad \text{and} \quad c_1 > 2\sqrt{r_1}, \end{cases} \quad (1.9)$$

or, equivalently,

$$\sigma(c_1; r_1, r_2) = \begin{cases} 2\sqrt{r_2} & \text{if } c_1 \leq 2\sqrt{r_2} \\ \lambda^* + \frac{r_1}{\lambda^*} & \text{if } r_1 < r_2 \text{ and } 2\sqrt{r_2} < c_1 \leq 2\sqrt{r_1} + 2\sqrt{r_2 - r_1} \\ 2\sqrt{r_1} & \text{if } c_1 > 2\sqrt{r_1} + 2\sqrt{\max\{0, r_2 - r_1\}}, \end{cases} \quad (1.10)$$

1.3. Related mathematical results. We also mention a closely related work of Choi, Giletti, and Guo [15], where they considered a two-species predator-prey system similar to (1.3), with the intrinsic growth rate $r = r(x - c_1 t)$ for the prey subject to the climate shift instead of the coefficient a . They considered the case when both initial data u_0 and v_0 are compactly supported and a non-decreasing profile for the growth rate with r changing sign, $r(-\infty) < 0 < r(\infty)$. In the case of local dispersal, they showed that the prey persists by spreading if and only if the maximal speed of the prey exceeds the environmental speed (i.e., $2\sqrt{dr(+\infty)} > c_1$), while the predator persists by spreading at the speed given by the smaller of the prey and maximal predator spreading speeds. In their setting both species tends to zero in $\{(t, x) : x < c_1 t\}$, while in the zone ahead of the environmental shift, the density of the prey is strictly decreasing so there is no nonlocal pulling phenomenon. We also mention [32] for the case of two weak-competing predators and one prey, and [1], for the case of one predator and two preys. For compactly supported initial data, the invasion wave of the prey resembles the effect of a shifting environment studied in our paper. However, the exact spreading speed of the predator(s) is not completely determined.

Finally, we mention the work of Bramson [12], in which established via probabilistic techniques a correction term of $\frac{3}{2} \log t$ which separates the the location of the spreading front for solutions to the Fisher-KPP equation (1.1) and the asymptotic location of the minimal traveling wave solution. This result was later generalized using maximum principle arguments by Lau [43] to KPP-like nonlinearities $f(s)$ satisfying $f'(s) \leq f'(0)$ on $[0, 1]$. For systems of equations of predator-prey type, the existence and characterization of such a delay between the spreading front and the asymptotic rate of spread is a challenging open question.

1.4. Organization of the paper. The rest of the paper will be organized as follows. In Section 2, we give a quick proof of the upper estimate of the spreading speed (namely, $\bar{c}_* \leq \sigma(c_1; r_1, r_2)$) by invoking the recent results on the diffusive logistic equation in shifting environment due to [41]. In Section 3, we derive some rough estimates for the prey density $v(t, x)$, and state five separate cases for the key parameters c_1, r_1, r_2 where the spreading speed has to be treated separately. In Section 4, we outline, in several lemmas, the conceptual steps to estimate the spreading speed from below via explicit solution of some Hamilton-Jacobi equation (4.4) obtained as the limiting problem of the first equation of (1.3). These lemmas will be proved in Subsections 4.1, 4.2 and 4.3. In Section 5, we determine the explicit formulas of the unique solution $\hat{\rho}$ to the limiting problem in each case, and prove that the upper bound of c^* obtained in Section 3 is also the lower bound. Thus the spreading speed is determined and Theorems 1.1 and 1.2 are proved. In Section 6, we prove Theorem 6.1 regarding the convergence to homogeneous state. Finally, in the Appendix, we collect some useful comparison results regarding the limiting Hamilton-Jacobi equations in [41], which are rephrased in a format suitable for our purpose here.

2. UPPER BOUND ON SPREADING SPEED

In this section, we give a quick proof of $\bar{c}_* \leq \sigma(c_1; r_1, r_2)$, where $\sigma(c_1; r_1, r_2)$ is given by (1.10) (which is equivalent to (1.9)). i.e., the spreading speed c^* is bounded above by $\sigma(c_1; r_1, r_2)$.

First, we establish some preliminary estimates on the solutions u and v of (1.3).

Lemma 2.1. *Assume $0 \leq u_0 \leq \|a\|_\infty - 1$ and $\beta \leq v_0 \leq 1$. Then the corresponding solutions $u(t, x)$ and $v(t, x)$ of (1.3) satisfy $0 \leq u(t, x) \leq \|a\|_\infty - 1$ and $\beta \leq v(t, x) \leq 1$ for all $(t, x) \in (0, \infty) \times \mathbb{R}$.*

Proof. By the classical theory of reaction-diffusion equations, there exists a unique solution (u, v) satisfying (1.3) for all $(t, x) \in (0, \infty) \times \mathbb{R}$; see, e.g. [58]. Moreover, since $0 \leq u_0 \leq \|a\|_\infty - 1$ and $\beta \leq v_0 \leq 1$, the maximum principle (see [56, Chapter 3, Section 6, Theorem 10] or [40, Theorem 6.2.1]) implies $0 \leq u \leq \|a\|_\infty - 1$ and $\beta \leq v \leq 1$ on $(0, \infty) \times \mathbb{R}$. \square

The global upper bound for v established in Lemma 2.1, combining with existing results for the diffusive logistic equation with heterogeneous shifting coefficients [41], can be used to determine an upper bound for the spreading speed of u .

Proposition 2.2. *Let $(u(t, x), v(t, x))$ be the solution of (1.3), with the associated maximal spreading speed \bar{c}_* as given in (1.7). Then*

$$\lim_{t \rightarrow \infty} \sup_{x \geq (\sigma + \eta)t} u(t, x) = 0 \quad \text{for each } \eta > 0. \tag{2.1}$$

In particular, $\bar{c}_ \leq \sigma$, where $\sigma = \sigma(c_1; r_1, r_2)$ is defined in (1.10).*

Proof. By Lemma 2.1, $v(t, x) \leq 1$ for all $(t, x) \in (0, \infty) \times \mathbb{R}$, hence we may regard $u(t, x)$ as a subsolution of the following scalar problem

$$\begin{cases} \bar{u}_t = \bar{u}_{xx} + \bar{u}(-1 - \bar{u} + a(x - c_1 t)) & \text{in } (0, \infty) \times \mathbb{R} \\ \bar{u}(0, x) = u_0(x) & \text{in } \mathbb{R}. \end{cases} \tag{2.2}$$

Let \bar{u} be the classical solution of (2.2) with initial data $u_0(x)$. By the parabolic maximum principle we have

$$u(t, x) \leq \bar{u}(t, x) \quad \text{for all } (t, x) \in (0, \infty) \times \mathbb{R}. \tag{2.3}$$

In the case $r_2 > r_1$, we may invoke [41, Theorem 6] to deduce that \bar{u} satisfies

$$\lim_{t \rightarrow \infty} \sup_{x \geq (\sigma + \eta)t} \bar{u}(t, x) = 0 \quad \text{for each } \eta > 0, \tag{2.4}$$

where $\sigma = \sigma(c_1; r_1, r_2)$ is given in (1.10).

In case $r_1 > r_2$, we define

$$U(t, x) = \begin{cases} \exp(-\sqrt{r_1}x + 2r_1 t) & \text{if } c_1 \geq 2\sqrt{r_1}, \\ \min\{r_1, \exp(-\lambda(x - c_1 t))\} & \text{if } 2\sqrt{r_2} < c_1 < 2\sqrt{r_1}, \lambda = \frac{1}{2} (c_1 - \sqrt{c_1^2 - 4r_2}) \\ \min\{r_1, \exp(-\sqrt{r_2}x + 2r_2 t)\} & \text{if } 0 < c_1 \leq 2\sqrt{r_2}. \end{cases}$$

Then it can be verified that U is a generalized supersolution of (2.2) (see [40, Definition 1.1.1] or [10, Definition 4.2] for the definition). Hence, we again deduce that (2.4) holds where

$$\sigma = \begin{cases} 2\sqrt{r_1} & \text{if } c_1 \geq 2\sqrt{r_1}, \\ 2\sqrt{r_2} & \text{if } c_1 \leq 2\sqrt{r_2}, \\ c_1 & \text{otherwise.} \end{cases}$$

Combining with (2.3), we conclude that

$$\lim_{t \rightarrow \infty} \sup_{x \geq (\sigma + \eta)t} u(t, x) = 0 \quad \text{for each } \eta > 0,$$

where σ is given in (1.10) (and $\sigma = c_1$, in case $2\sqrt{r_2} < c_1 < 2\sqrt{r_1}$). This completes the proof. \square

3. ROUGH ESTIMATE FOR $v(t, x)$

Having established that the spreading speed is bounded above by $\sigma = \sigma(c_1; r_1, r_2)$, we may also deduce in the following lemma that $v(t, x)$ converges to its carrying capacity as $t \rightarrow \infty$ in the region $\{(x, t) : x > \sigma t\}$.

Lemma 3.1. *Let $(u(t, x), v(t, x))$ be the solution of (1.3). Then*

$$\lim_{t \rightarrow \infty} \sup_{x \geq (\sigma + \eta)t} |v(t, x) - 1| = 0 \quad \text{for each } \eta > 0, \quad (3.1)$$

where σ is given by (1.10).

Proof. Since $v(t, x) \leq 1$ (thanks to Lemma 2.1), it suffices to show the lower bound. We shall follow the proof of Theorem 5.1 in [21]. Fix $c > \sigma(c_1; r_1, r_2)$. We may suppose for contradiction that there exists a sequence $\{(t_n, x_n)\}$ with $t_n \rightarrow \infty$ and $x_n \geq ct_n$ such that $\limsup_{n \rightarrow \infty} v(t_n, x_n) < 1$. Denote $(u_n, v_n)(t, x) = (u, v)(t + t_n, x + x_n)$. By standard parabolic estimates, we may pass to a further subsequence so that (u_n, v_n) converges to an entire in time solution (u_∞, v_∞) of (1.3) in $C_{loc}(\mathbb{R}^2)$. Since $c > \sigma(c_1; r_1, r_2)$, by Proposition 2.2, we have $u_\infty \equiv 0$. Thus, v_∞ is an entire solution satisfying the equation

$$(v_\infty)_t = d(v_\infty)_{xx} + rv_\infty(1 - v_\infty) \quad \text{for } (t, x) \in \mathbb{R}^2.$$

Since $v \geq \beta$ for all $(t, x) \in (0, \infty) \times \mathbb{R}$, we deduce that $v_\infty \geq \beta$ for all $(t, x) \in \mathbb{R}^2$. By the classification of entire solutions of the diffusive logistic equation (see, e.g. [49, Lemma 2.3(d)]) we have $v_\infty \equiv 1$. This is in contradiction with the statement $\limsup_{n \rightarrow \infty} v(t_n, x_n) < 1$. \square

Having established the upper bound of the spreading speed, the outstanding task is to estimate the spreading speed from below. We will do so by adopting the Hamilton-Jacobi approach [25]. To this end, define

$$F^\epsilon(t, x) = -1 + v\left(\frac{t}{\epsilon}, \frac{x}{\epsilon}\right)a\left(\frac{x}{\epsilon} - \frac{c_1 t}{\epsilon}\right) \quad (3.2)$$

and its (lower) half-relaxed limit [6]

$$F_*(t, x) = \liminf_{\substack{\epsilon \rightarrow 0 \\ (t', x') \rightarrow (t, x)}} F^\epsilon(t', x'). \quad (3.3)$$

Thanks to (H1), the function $a(s)$ is monotone.

We will divide the proof of the spreading speed into the following cases, depending on the speed of environmental shift c_1 and the profile of the conversion efficiency $a(x - c_1 t)$.

- Case 1(a):** $r_1 < r_2$ and $c_1 \leq 2\sqrt{r_2}$
- Case 1(b):** $r_1 < r_2$ and $2\sqrt{r_2} < c_1 < 2(\sqrt{r_1} + \sqrt{r_2 - r_1})$
- Case 1(c):** $r_1 < r_2$ and $c_1 \geq 2(\sqrt{r_1} + \sqrt{r_2 - r_1})$
- Case 2(a):** $r_1 > r_2$ and $c_1 < 2\sqrt{r_2}$
- Case 2(b):** $r_1 > r_2$ and $c_1 = 2\sqrt{r_2}$
- Case 2(c):** $r_1 > r_2$ and $c_1 > 2\sqrt{r_1}$

In Case 1(a) - (c), we have $r_1 < r_2$, and we let

$$R_1(s) = \begin{cases} R_{1a}(s) = \begin{cases} r_2 & \text{for } s > 2\sqrt{r_2} \\ r_2 & \text{for } c_1 < s \leq 2\sqrt{r_2} \\ r_1 & \text{for } s \leq c_1 \end{cases} & \text{if } c_1 \leq 2\sqrt{r_2}, \\ R_{1b}(s) = \begin{cases} r_2 & \text{for } s > c_1, \\ r_1 & \text{for } \lambda^* + \frac{r_1}{\lambda^*} < s \leq c_1 \\ r_1 & \text{for } s \leq \lambda^* + \frac{r_1}{\lambda^*} \end{cases} & \text{if } 2\sqrt{r_2} < c_1 < 2(\sqrt{r_1} + \sqrt{r_2 - r_1}), \\ R_{1c}(s) = \begin{cases} r_2 & \text{for } s > c_1, \\ r_1 & \text{for } 2\sqrt{r_1} < s \leq c_1 \\ r_1 & \text{for } s \leq 2\sqrt{r_1} \end{cases} & \text{if } c_1 \geq 2(\sqrt{r_1} + \sqrt{r_2 - r_1}) \end{cases} \quad (3.4)$$

In Cases 2(a)-(c), we have $r_1 > r_2$, and we let

$$R_2(s) = \begin{cases} R_{2a}(s) = \begin{cases} r_2 & \text{for } s > 2\sqrt{r_2}, \\ r_2 & \text{for } c_1 \leq s \leq 2\sqrt{r_2} \\ r_1 & \text{for } s < c_1. \end{cases} & \text{if } c_1 < 2\sqrt{r_2}, \\ R_{2b}(s) = \begin{cases} r_2 & \text{for } s > 2\sqrt{r_2}, \\ \min\{r_2, r_1\} & \text{for } s = 2\sqrt{r_2} \\ r_1 & \text{for } s < 2\sqrt{r_2}. \end{cases} & \text{if } c_1 = 2\sqrt{r_2}, \\ R_{2c}(s) = \begin{cases} r_2 & \text{for } s \geq c_1, \\ r_1 & \text{for } 2\sqrt{r_1} < s < c_1 \\ r_1 & \text{for } s \leq 2\sqrt{r_1} \end{cases} & \text{if } c_1 > 2\sqrt{r_1}. \end{cases} \quad (3.5)$$

Lemma 3.2. $F_*(t, x) \geq R_i(x/t)$ in cases 1(a)-(c) and 2(a)-(c).

Proof. The lemma follows from the definition of F_* , in (3.3), and the global bounds $\beta \leq v(t, x) \leq 1$ (Lemma 2.1). \square

4. LOWER BOUND ON THE SPREADING SPEED

We will use the Hamilton-Jacobi method to prove a lower bound for the spreading speed. To this end, define the WKB-Ansatz [25]

$$w^\epsilon(t, x) = -\epsilon \log u^\epsilon(t, x) \quad \text{where} \quad u^\epsilon(t, x) = u(t/\epsilon, x/\epsilon), \quad (4.1)$$

and consider the half-relaxed limits [7]

$$w^*(t, x) = \limsup_{\substack{\epsilon \rightarrow 0 \\ (t', x') \rightarrow (t, x)}} w^\epsilon(t', x') \quad \text{and} \quad w_*(t, x) = \liminf_{\substack{\epsilon \rightarrow 0 \\ (t', x') \rightarrow (t, x)}} w^\epsilon(t', x'). \quad (4.2)$$

In the following lemma, we show that $w^*(t, x)$ and $w_*(t, x)$ can be related to one-dimensional functions $\rho^*(s)$ and $\rho_*(s)$, respectively.

Lemma 4.1. *Let w^* and w_* be defined as in (4.2). Then $w^*(t, x) = t\rho^*(x/t)$ and $w_*(t, x) = t\rho_*(x/t)$ for some functions ρ^* and ρ_* .*

Proof. For the existence of ρ^* , we may compute

$$w^*(t, x) = \limsup_{\substack{\epsilon \rightarrow 0 \\ (t', x') \rightarrow (t, x)}} -\epsilon \log u\left(\frac{t'}{\epsilon}, \frac{x'}{\epsilon}\right) = t \limsup_{\substack{\epsilon \rightarrow 0 \\ (t'', x'') \rightarrow (1, x/t)}} -(\epsilon/t) \log u\left(\frac{t''}{\epsilon/t}, \frac{x''}{\epsilon/t}\right).$$

Thus $w^*(t, x) = tw^*(1, x/t)$, and the first part of the result is proved if we take $\rho^*(s) = w^*(1, s)$. The proof of the second part is analogous. \square

Next, we describe a bird’s-eye view of the Hamilton-Jacobi approach in order to achieve our final goal of bounding the spreading speed from below by the optimal constant $\sigma > 0$. For clarity, we will state the necessary lemmas and provide their proofs later on.

We start with the following lemma which is due to [25] for the KPP equation. The proof is presented in Subsection 4.1.

Lemma 4.2. *Suppose that there is $s_0 > 0$ such that $\rho^*(s) = 0$ for all $s \in [0, s_0]$. Then there exists $\delta_0 > 0$ such that*

$$\liminf_{t \rightarrow \infty} \inf_{\eta t < x < (s_0 - \eta)t} u(t, x) \geq \delta_0 \quad \text{for each } \eta > 0 \text{ sufficiently small.}$$

Hence, a lower bound of the spreading speed (and hence the complete proofs of our main theorems) can be obtained by determining the set $\{s : \rho^*(s) = 0\}$. Precisely, it is sufficient to show that $\rho^*(s) = 0$ for $s \in [0, \sigma]$, where $\sigma = \sigma(c_1; r_1, r_2)$ as in (1.10).

To this end, we derive a limiting Hamilton-Jacobi equation for w^* and then for ρ^* . Observe that w^ϵ satisfies

$$w_t^\epsilon - \epsilon w_{xx}^\epsilon + |w_x^\epsilon|^2 + (F^\epsilon(t, x) - u^\epsilon) = 0 \quad \text{for } (t, x) \in (0, \infty) \times \mathbb{R}, \tag{4.3}$$

where F^ϵ is given in (3.2). By the fact that $F_*(t, x) \geq R(x/t)$, it is standard [25, 41] to deduce the following.

Lemma 4.3. *Suppose $F_*(t, x) \geq R(x/t)$, then ρ^* is a viscosity subsolution of*

$$\min\{\rho - s\rho' + |\rho'|^2 + R(s), \rho\} = 0 \quad \text{for } s \in (0, \infty). \tag{4.4}$$

Moreover,

$$\rho^*(0) = 0 \quad \text{and} \quad \rho^*(s) < \infty \quad \text{for all } s \in [0, \infty). \tag{4.5}$$

Proof. We postpone the proof to Subsection 4.2. \square

By the comparison principle, discussed in the Appendix, the Hamilton-Jacobi equation (4.4) has a unique viscosity solution.

Lemma 4.4. *For any given case $(i, j) \in \{(1, a), (1, b), (1, c), (2, a), (2, b), (2, c)\}$ as stated in Section 3, let R be given by $R = R_{ij}$. The Hamilton-Jacobi equation (4.4) has a unique viscosity solution, $\hat{\rho}$, satisfying*

$$\hat{\rho}(0) = 0 \quad \text{and} \quad \lim_{s \rightarrow \infty} \frac{\hat{\rho}(s)}{s} = \infty. \tag{4.6}$$

Moreover, $\hat{\rho}$ is nondecreasing in s , i.e.

$$\hat{\rho}(s) \equiv 0 \quad \text{for } 0 \leq s \leq \sup\{s' \geq 0 : \hat{\rho}(s') = 0\}. \tag{4.7}$$

Furthermore, the following Lemma holds:

Lemma 4.5. *For any given case $(i, j) \in \{(1, a), (1, b), (1, c), (2, a), (2, b), (2, c)\}$ as stated in Section 3, let R be given by $R = R_{ij}$ and let $\hat{\rho}$ be the unique solution of (4.4) as specified in Lemma 4.4. Then*

$$0 \leq \rho^*(s) \leq \hat{\rho}(s) \quad \text{for } s \geq 0.$$

The statement (4.7) and Lemma 4.5 imply that

$$\rho^*(s) = 0 \quad \text{for } 0 \leq s \leq \sup\{s' \geq 0 : \hat{\rho}(s') = 0\}. \quad (4.8)$$

Together with Lemma 4.2, this enables us to establish

$$c_* \geq \sup\{s \geq 0 : \hat{\rho}(s) = 0\}.$$

Next, we will establish the explicit formula of the unique solution $\hat{\rho}$ satisfying (4.4) in viscosity sense and (4.6) in classical sense in Lemma 5.1.

For example, in Case 1(a) ($r_1 < r_2$ and $c_1 \leq 2\sqrt{r_2}$), we find that

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > 2\sqrt{r_2} \\ 0 & \text{for } 0 \leq s \leq 2\sqrt{r_2}. \end{cases} \quad (4.9)$$

Thanks to Lemma 4.5, we have $0 \leq \rho^*(s) \leq \hat{\rho}(s)$. Hence, we deduce that

$$\rho^*(s) = 0 \quad \text{for } 0 \leq s \leq 2\sqrt{r_2}.$$

By Lemma 4.2, we conclude that $c_* \geq 2\sqrt{r_2}$. This establishes the lower bound of the spreading speed in Case 1(a). The spreading speed in Case 1(a) is thus determined, since $2\sqrt{r_2}$ is also the upper bound of spreading speed (thanks to Proposition 2.2).

In the remainder of this section, we present the proofs of the above lemmas.

4.1. Proof of Lemma 4.2.

Proof of Lemma 4.2. Our proof is adapted from Theorem 1.1 of [25]. Fix a small $0 < \eta \ll 1$. It is sufficient to show that there exists $\delta_0 = \delta_0(\eta) > 0$ such that

$$\liminf_{\epsilon \rightarrow 0} \inf_K u^\epsilon(t, x) \geq \delta_0 \quad (4.10)$$

for any compact set given by $K = \{(1, x) : \eta \leq x \leq s_0 - \eta\} \subset \subset \{(t, x) : 0 < x/t < s_0\}$. Indeed,

$$\liminf_{t \rightarrow \infty} \inf_{\eta t < x < (s_0 - \eta)t} u(t, x) = \liminf_{\epsilon \rightarrow 0} \inf_K u^\epsilon(t, x) \geq \delta_0.$$

To show (4.10), we first observe that $w^*(t, x) = t\rho^*(x/t) = 0$ in some compact subset \tilde{K} such that

$$K \subset \text{Int } \tilde{K} \subset \{(t, x) : 0 < x/t < s_0\},$$

which implies $w^\epsilon(t, x) \rightarrow 0$ uniformly in a neighborhood of K . Now for $(t_0, x_0) \in K$, let $\psi(t, x) = (t - t_0)^2 + (x - x_0)^2$. Then $w^* - \psi$ has a strict local maximum at (t_0, x_0) . Since $w^\epsilon \rightarrow 0$ uniformly in a neighborhood of K , for each $\epsilon > 0$ sufficiently small, the function $w^\epsilon - \psi$ has a local maximum at $(t_\epsilon, x_\epsilon) \in K$, where $(t_\epsilon, x_\epsilon) \rightarrow (t_0, x_0)$ as $\epsilon \rightarrow 0$. Thus,

$$o(1) = \partial_t \psi - \epsilon \partial_{xx} \psi + |\partial_x \psi|^2 \leq \partial_t w^\epsilon - \epsilon \partial_{xx} w^\epsilon + |\partial_x w^\epsilon|^2 = u^\epsilon - F^\epsilon \leq u^\epsilon - \delta_0 \quad (4.11)$$

at $(t, x) = (t_\epsilon, x_\epsilon)$, where $\delta_0 = \beta \inf_{s \in \mathbb{R}} a(s) - 1 > 0$.

Using the fact that $w^\epsilon - \psi$ has a local maximum at (t_ϵ, x_ϵ) , we deduce that

$$w^\epsilon(t_\epsilon, x_\epsilon) \geq (w^\epsilon - \psi)(t_\epsilon, x_\epsilon) \geq (w^\epsilon - \psi)(t_0, x_0) = w^\epsilon(t_0, x_0)$$

which implies that $u^\epsilon(t_0, x_0) \geq u^\epsilon(t_\epsilon, x_\epsilon)$. Combining with (4.11), we have

$$u^\epsilon(t_0, x_0) \geq u^\epsilon(t_\epsilon, x_\epsilon) \geq \delta_0 + o(1).$$

Since the above argument is uniform for arbitrary $(t_0, x_0) \in K$, this implies (4.10). \square

4.2. **Proof of Lemma 4.3.** We recall the definition of viscosity solutions of (4.4), following [6, 38] (see also [25, 41] for the definition involving variational inequalities).

Definition 4.1. In the following let R^* and R_* be the upper and lower envelope of R , which is given by

$$R^*(s) = \limsup_{s' \rightarrow s} R(s') \quad \text{and} \quad R_*(s) = \liminf_{s' \rightarrow s} R(s').$$

- A lower semicontinuous function $\hat{\rho}$ is called a viscosity supersolution of (4.4) if $\hat{\rho} \geq 0$, and for any test function $\phi \in C^1$, if s_0 is a strict local minimum of $\hat{\rho} - \phi$, then

$$\hat{\rho}(s_0) - s_0\phi'(s_0) + |\phi'(s_0)|^2 + R^*(s_0) \geq 0.$$

- An upper semicontinuous function $\hat{\rho}$ is called a viscosity subsolution of (4.4) if for any test function $\phi \in C^1$, if s_0 is a strict local maximum of $\hat{\rho} - \phi$ and $\hat{\rho}(s_0) > 0$, then

$$\hat{\rho}(s_0) - s_0\phi'(s_0) + |\phi'(s_0)|^2 + R_*(s_0) \leq 0.$$

- We say $\hat{\rho}$ is a viscosity solution of (4.4) if $\hat{\rho}$ is a viscosity super- and subsolution.

We will first show that ρ^* is nonnegative and that $\rho^*(0) = 0$.

Lemma 4.6. *Let ρ^* be defined as in Lemma 4.1. Then*

$$\rho^*(0) = 0 \quad \text{and} \quad \rho^*(s) \geq 0 \quad \text{for } s \geq 0. \quad (4.12)$$

Proof. We first show $w^*(t, x) \geq 0$ for $(t, x) \in (0, \infty) \times \mathbb{R}$. Indeed, since $u(t, x) \leq \max\{r_2, r_1\}$ for all $(t, x) \in [0, \infty) \times \mathbb{R}$, by the definition of w^ϵ we have $w^\epsilon \geq -\epsilon \log(\max\{r_2, r_1\})$ for each $\epsilon > 0$ and $(t, x) \in (0, \infty) \times \mathbb{R}$, and we may compute

$$w^*(t, x) = \limsup_{\substack{\epsilon \rightarrow 0 \\ (t', x') \rightarrow (t, x)}} w^\epsilon(t', x') \geq 0 \quad \text{for } (t, x) \in (0, \infty) \times \mathbb{R}. \quad (4.13)$$

In particular, $w^*(t, 0) \geq 0$ for $t > 0$.

The proof will be complete once we show $w^*(t, 0) \leq 0$ for $t > 0$. Denote $\underline{r} = r_1 \wedge r_2$. Then using the lower bound $v \geq \beta$, we see that u is a supersolution of

$$\underline{u}_t - \underline{u}_{xx} = \underline{u}(\underline{r} - \underline{u}) \quad \text{in } (0, \infty) \times \mathbb{R}. \quad (4.14)$$

Let $\underline{u}(t, x)$ be the solution of (4.14) with identical (compactly supported) initial condition as $u(t, x)$, then the classical spreading result for the Fisher-KPP equation [3] says that \underline{u} has spreading speed $2\sqrt{\underline{r}}$. In particular,

$$\liminf_{t \rightarrow \infty} \inf_{|x| < \sqrt{\underline{r}t}} \underline{u}(t, x) \geq 2\delta_1 \quad \text{for some } \delta_1 > 0.$$

By the comparison principle, $u \geq \underline{u}$, i.e. there exists $t_1 > 0$ such that

$$\inf_{|x| < \sqrt{\underline{r}t}} u(t, x) \geq \delta_1 \quad \text{for } t \geq t_1,$$

which implies

$$\sup_{\substack{|x| < \sqrt{\underline{r}t} \\ t \geq \epsilon t_1}} w^\epsilon(t, x) \leq -\epsilon \log \delta_1. \quad (4.15)$$

Now, fix an arbitrary $t_0 > 0$. Let $(t, x) \rightarrow (t_0, 0)$ and $\epsilon \rightarrow 0$, we deduce

$$w^*(t_0, 0) = \limsup_{\substack{\epsilon \rightarrow 0 \\ (t, x) \rightarrow (t_0, 0)}} w^\epsilon(t, x) \leq 0. \quad (4.16)$$

Combining (4.13) and (4.16), we have

$$w^*(t, 0) = 0 \quad \text{for all } t > 0. \quad (4.17)$$

We recall $w^*(t, x) = t\rho^*(x/t)$ (thanks to Lemma 4.1), so that (4.12) directly follows from (4.13) and (4.17). This completes the proof. \square

The following lemma implies that $\rho^*(s) < \infty$ for $s \in [0, \infty)$.

Lemma 4.7. *Let w^ϵ be a solution of (4.3). Then for each compact subset Q of $(0, \infty) \times \mathbb{R}$, there is a constant $C(Q)$ independent of ϵ such that*

$$w^\epsilon(t, x) \leq C(Q) \quad \text{for } (t, x) \in Q \text{ and } \epsilon \in (0, 1/C(Q)).$$

In particular,

$$w^*(t, x) < +\infty \quad \text{for each } (t, x) \in (0, \infty) \times \mathbb{R} \quad \text{and} \quad \rho^*(s) < +\infty \quad \text{for each } s \in [0, \infty). \quad (4.18)$$

Proof. We only prove the bound for $Q \subset (0, \infty) \times [0, \infty)$. The case for $Q \subset (0, \infty) \times (-\infty, 0]$ is similar and is omitted. Our proof follows the ideas in [25]. Fix $\delta \in (0, 1)$ such that $Q \subset [\delta, 1/\delta] \times [0, 1/\delta]$. Define

$$z^\epsilon(t, x) = \frac{|x + 2\delta|^2}{4t} + \frac{\epsilon}{2} \log t + C_\delta(1 + t).$$

By taking $C_\delta > 0$ to be a large constant depending on δ , z^ϵ is a (classical) supersolution of (4.3) in $(0, \infty) \times (0, \infty)$.

By (4.17) in the proof of Lemma 4.6 and the definition of w^* , there is a constant $C_\delta > 0$ such that,

$$\sup_{0 < \epsilon \leq 1/2} w^\epsilon(t + \delta/2, 0) \leq C_\delta \quad \text{for } t \in [0, 1/\delta].$$

Observe that for ϵ sufficiently small, we have

$$\begin{cases} w^\epsilon(\delta/2, x) < \infty = z^\epsilon(0, x) & \text{for } x \geq 0 \\ w^\epsilon(t + \delta/2, 0) \leq C_\delta \leq z^\epsilon(t, 0) & \text{for } t \in [0, 1/\delta]. \end{cases}$$

It follows from the maximum principle that

$$w^\epsilon(t + \delta/2, x) \leq z^\epsilon(t, x) \quad \text{for } (t, x) \in [0, 1/\delta] \times [0, \infty).$$

Taking supremum over $[\delta/2, 1/\delta] \times [0, 1/\delta]$, we have

$$\sup_{[\delta/2, 1/\delta] \times [0, 1/\delta]} w^\epsilon(t + \delta/2, x) \leq C'_\delta := \sup_{[\delta/2, 1/\delta] \times [0, 1/\delta]} \left[\frac{|x + 2\delta|^2}{4t} + \log t + C_\delta(1 + t) \right]. \quad (4.19)$$

This completes the proof. \square

Now, we are in position to prove Lemma 4.3.

Proof of Lemma 4.3. Since (4.5) is a consequence of Lemma 4.6 and (4.18), it remains to show that ρ^* is a viscosity subsolution of (4.4).

Let $\phi \in C^1$ be a test function and suppose that $\rho^* - \phi$ has a strict local maximum at $s = s_0$, and that $\rho^*(s_0) > 0$. Without loss of generality, we may assume that $\rho^* - \phi \leq 0$ for all s near s_0 , with equality holding only at $s = s_0$. We will show that

$$\rho^*(s_0) - s_0\phi'(s_0) + |\phi'(s_0)|^2 + R_*(s_0) \leq 0.$$

(Note that $R_*(s_0) = R(s_0)$ by our definition of R in (3.4) and (3.5).) First, we note that $w^*(t, x) = t\rho^*(x/t)$ and that $w^*(t, x) - t\phi(x/t) - (t-1)^2 \leq 0$ for all (t, x) near $(1, s_0)$, with equality holding only at $(t, x) = (1, s_0)$. Define, in terms of ϕ , a two variable test function

$$\varphi(t, x) = t\phi(x/t) - (t-1)^2.$$

Then by the definition of w^* , there exists a sequence $\epsilon_n \rightarrow 0$ and sequence of points $(t_n, x_n) \rightarrow (1, s_0)$ as $n \rightarrow \infty$ such that: $w^{\epsilon_n} - \varphi$ has a local maximum at (t_n, x_n) , and $w^{\epsilon_n}(t_n, x_n) \rightarrow w^*(t_0, x_0) > 0$. Thus, for $(t, x) = (t_n, x_n)$, we have

$$\begin{aligned} \partial_t \varphi &= \partial_t w^{\epsilon_n} = \epsilon_n \partial_{xx} w^{\epsilon_n} - |\partial_x w^{\epsilon_n}|^2 - (F^{\epsilon_n} - u^{\epsilon_n}) \\ &\leq \epsilon_n \partial_{xx} \varphi - |\partial_x \varphi|^2 - (F^{\epsilon_n} - u^{\epsilon_n}). \end{aligned}$$

Thus,

$$\partial_t \varphi - \epsilon_n \partial_{xx} \varphi + |\partial_x \varphi|^2 + (F^{\epsilon_n} - u^{\epsilon_n}) \leq 0 \quad (4.20)$$

for $(t, x) = (t_n, x_n)$. Letting $n \rightarrow \infty$, we obtain

$$\partial_t \varphi(t_0, x_0) + |\partial_x \varphi(t_0, x_0)|^2 + F_*(t_0, x_0) \leq 0, \quad (4.21)$$

where we have used the fact that $w^{\epsilon_n}(t_n, x_n) \rightarrow w^*(t_0, x_0) > 0$ implies $u^{\epsilon_n}(t_n, x_n) \rightarrow 0$. Since $F_*(t_0, x_0) \geq R(x_0/t_0)$, it follows from (4.21) that

$$\phi(s_0) - s_0 \phi'(s_0) + |\phi'(s_0)|^2 + R_*(s_0) \leq 0.$$

Thus, ρ^* is a viscosity subsolution of (4.4). □

4.3. Proofs of Lemmas 4.4 and 4.5.

Proof of Lemma 4.4. Observe, in each case $i = 1, 2$, that our choice of R satisfies (B1)-(B2) of the Appendix. By Corollary A.2 of the Appendix, there exists a unique $\hat{\rho}$ satisfying (4.4) in the viscosity sense, and the boundary condition (4.6) in the classical sense. Moreover, $s \mapsto \hat{\rho}$ is nondecreasing. This proves Lemma 4.4. □

Proof of Lemma 4.5. We observe that ρ^* is a viscosity subsolution (by Lemma 4.3) and that $\hat{\rho}$ is a viscosity supersolution (by Lemma 4.4). Moreover, by (4.5) and (4.6), we have

$$\rho^*(0) = \hat{\rho}(0) = 0, \quad \text{and} \quad \lim_{s \rightarrow \infty} \frac{\rho^*(s)}{s} \leq +\infty = \lim_{s \rightarrow \infty} \frac{\hat{\rho}(s)}{s}.$$

We can therefore apply the comparison principle (see Lemma A.1 of the Appendix) to derive

$$\rho^*(s) \leq \hat{\rho}(s) \quad \text{for } s \geq 0.$$

Finally, $\rho^*(s) \geq 0$ is proved in Lemma 4.6. □

5. SOLVING FOR THE SPREADING SPEED VIA EXPLICIT FORMULAS FOR $\hat{\rho}$

For each of the cases 1(a)-(c), 2(a)-(c), we will propose an explicit formula for $\hat{\rho}$ in Subsection 5.1. Thanks to the uniqueness result in Lemma 4.4, it is enough to verify (separately for each of the cases) that the given expression defines a viscosity solution of (4.4). This will be done in Subsection 5.2.

5.1. **Explicit formulas for $\hat{\rho}$.** Below, we state the explicit formula for $\hat{\rho}$ in each case. Subsequently, we will verify in Lemma 5.1 that $\hat{\rho}$ solves (4.4) by invoking the definition of the viscosity solution [6].

- **Case 1(a):** $r_1 < r_2$ and $c_1 \leq 2\sqrt{r_2}$.

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > 2\sqrt{r_2} \\ 0 & \text{for } 0 \leq s \leq 2\sqrt{r_2}, \end{cases} \quad (5.1)$$

- **Case 1(b):** $r_1 < r_2$ and $2\sqrt{r_2} < c_1 < 2(\sqrt{r_1} + \sqrt{r_2 - r_1})$.

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > c_1 \\ \lambda^* s - ((\lambda^*)^2 + r_1) & \text{for } (\lambda^* + \frac{r_1}{\lambda^*}) < s \leq c_1 \\ 0 & \text{for } 0 \leq s \leq \lambda^* + \frac{r_1}{\lambda^*}, \end{cases} \quad (5.2)$$

where $\lambda^* = c_1/2 - \sqrt{r_2 - r_1}$.

- **Case 1(c):** $r_1 < r_2$ and $c_1 \geq 2(\sqrt{r_1} + \sqrt{r_2 - r_1})$.

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > c_1 \\ \lambda^* s - ((\lambda^*)^2 + r_1) & \text{for } 2\lambda^* < s \leq c_1 \\ \frac{s^2}{4} - r_1 & \text{for } 2\sqrt{r_1} < s \leq 2\lambda^* \\ 0 & \text{for } 0 \leq s \leq 2\sqrt{r_1}. \end{cases} \quad (5.3)$$

- **Case 2(a):** $r_1 > r_2$ and $c_1 < 2\sqrt{r_2}$.

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > 2\sqrt{r_2} \\ 0 & \text{for } 0 \leq s \leq 2\sqrt{r_2}, \end{cases} \quad (5.4)$$

- **Case 2(b):** $r_1 > r_2$ and $c_1 = 2\sqrt{r_2}$.

In this case, $\hat{\rho}(s)$ is defined the same as in (5.4).

- **Case 2(c):** $r_1 > r_2$ and $c_1 > 2\sqrt{r_1}$.

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > 2\tilde{\lambda} \\ \tilde{\lambda}s - (\tilde{\lambda}^2 + r_2) & \text{for } c_1 < s \leq 2\tilde{\lambda} \\ \frac{s^2}{4} - r_1 & \text{for } 2\sqrt{r_1} < s < c_1 \\ 0 & \text{for } 0 \leq s \leq 2\sqrt{r_1}, \end{cases} \quad (5.5)$$

where $\tilde{\lambda} = c_1/2 + \sqrt{r_1 - r_2}$.

5.2. $\hat{\rho}$ solves the HJE (4.4) in the viscosity sense.

Lemma 5.1. *There exists a unique viscosity solution $\hat{\rho}$ of (4.4) satisfying (in the classical sense)*

$$\hat{\rho}(0) = 0 \quad \text{and} \quad \lim_{s \rightarrow \infty} \frac{\hat{\rho}(s)}{s} = \infty. \quad (5.6)$$

Moreover, it is given by the formulas of Subsection 5.1.

Proof. By Lemma 4.4, $\hat{\rho}$ is unique whenever it exists. To prove the lemma, it is enough to show that $\hat{\rho}$, as given in each case of Subsection 5.1, satisfies (i) the Hamilton-Jacobi equation (4.4) in the viscosity sense, as well as (ii) the boundary condition (5.6) in the classical sense. In view of the explicit formulas, (ii) is obvious. Thus, it remains to verify that $\hat{\rho}$ is a viscosity solution of (4.4) in each case.

Since the proof only differs slightly in each case, we consider only two representative cases 1(b) and 2(c) here, and omit the verification of the rest.

Let us proceed with Case 1(b), where we fix r_1, r_2, c_1 satisfying

$$r_1 < r_2, \quad \text{and} \quad 2\sqrt{r_2} < c_1 < 2(\sqrt{r_1} + \sqrt{r_2 - r_1}). \quad (5.7)$$

Next, we set

$$R(s) = R_{1b}(s) = \begin{cases} r_2 & \text{for } s > c_1, \\ r_1 & \text{for } \lambda^* + \frac{r_1}{\lambda^*} < s \leq c_1 \\ \underline{r}_1 & \text{for } s \leq \lambda^* + \frac{r_1}{\lambda^*}. \end{cases}$$

and

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > c_1 \\ \lambda^* s - ((\lambda^*)^2 + r_1) & \text{for } (\lambda^* + \frac{r_1}{\lambda^*}) < s \leq c_1 \\ 0 & \text{for } 0 \leq s \leq \lambda^* + \frac{r_1}{\lambda^*}, \end{cases} \quad (5.8)$$

where $\lambda^* = c_1/2 - \sqrt{r_2 - r_1}$.

First, observe that $\hat{\rho}$ is continuous, thanks to our choice of λ^* .

Next, we show that $\hat{\rho}$ is a viscosity subsolution of (4.4). To this end, observe that $\hat{\rho}$ satisfies the equation (4.4) in the classical sense almost everywhere in $[0, \infty)$. (In fact, it satisfies the equation classically for $s \in \mathbb{R} \setminus \{c_1, \lambda^* + r_1/\lambda^*\}$.) By the convexity of the Hamiltonian, we can apply [5, Proposition 5.1] to conclude that it is in fact a viscosity subsolution of (4.4).

Next, we show that $\hat{\rho}$ is a viscosity supersolution of (4.4). Suppose $\hat{\rho} - \phi$ obtains a strict local minimum at $s_0 \in [0, \infty)$ for some $\phi \in C^1$. Now, $\hat{\rho}$ is a classical solution of (4.4) for all $s \notin \{\lambda^* + r_1/\lambda^*, c_1\}$, so it automatically satisfies (4.4) in the viscosity sense. We need only consider $s_0 \in \{\lambda^* + r_1/\lambda^*, c_1\}$. Suppose $s_0 = \lambda^* + r_1/\lambda^*$. Then $0 \leq \phi'(s_0) \leq \lambda^*$, and $R^*(s_0) = \max\{\underline{r}_1, r_1\} = r_1$. Therefore, at the point $s = s_0$, it holds that

$$\rho - s_0\phi' + |\phi'|^2 + R^* = -(\lambda^* + \frac{r_1}{\lambda^*})\phi' + |\phi'|^2 + r_1 = (\phi' - \lambda^*)(\phi' - \frac{r_1}{\lambda^*}) \geq 0,$$

where the last inequality is a consequence of $\phi'(s_0) \leq \lambda^* < \frac{r_1}{\lambda^*}$ (which in turn follows from the choice of λ^* and the condition (5.7)).

If $s_0 = c_1$, then $R^*(s_0) = \max\{r_1, r_2\} = r_2$, and we have

$$\rho - s_0\phi' + |\phi'|^2 + R^* = \left(\frac{c_1^2}{4} - r_2\right) - c_1\phi' + |\phi'|^2 + r_2 = (\phi' - \frac{c_1}{2})^2 \geq 0 \quad \text{at } s = s_0.$$

This proves that $\hat{\rho}$ is a viscosity supersolution.

This completes the proof that the unique viscosity solution $\hat{\rho}$ as guaranteed by Lemma 4.4 is given by the explicit formula (5.8) for the first representative Case 1(b).

Let us proceed with Case 2(c), where we fix r_1, r_2, c_1 satisfying

$$r_1 > r_2, \quad \text{and} \quad c_1 > 2\sqrt{r_1}. \quad (5.9)$$

Next, we set

$$R(s) = R_{2c}(s) = \begin{cases} r_2 & \text{for } s \geq c_1, \\ r_1 & \text{for } 2\sqrt{r_1} < s < c_1 \\ \underline{r}_1 & \text{for } s \leq 2\sqrt{r_1} \end{cases}$$

and

$$\hat{\rho}(s) = \begin{cases} \frac{s^2}{4} - r_2 & \text{for } s > 2\tilde{\lambda} \\ \tilde{\lambda}s - (\tilde{\lambda}^2 + r_2) & \text{for } c_1 < s \leq 2\tilde{\lambda} \\ \frac{s^2}{4} - r_1 & \text{for } 2\sqrt{r_1} < s \leq c_1 \\ 0 & \text{for } 0 \leq s \leq 2\sqrt{r_1}, \end{cases} \quad (5.10)$$

where $\tilde{\lambda} = c_1/2 + \sqrt{r_1 - r_2}$.

Again, we first observe that $\hat{\rho}$, as given in (5.10), is continuous, and satisfies the equation (4.4) in the classical sense for $s \in [0, \infty) \setminus \{c_1, 2\sqrt{r_1}\}$ (note that it is in fact continuously differentiable in a neighborhood of $s = 2\tilde{\lambda}$). It then follows again from [5, Proposition 5.1] that $\hat{\rho}$ is a viscosity subsolution of (4.4).

Next, we verify that it is also a viscosity supersolution. Suppose $\hat{\rho} - \phi$ obtains a strict local minimum at $s_0 \in [0, \infty)$ for some $\phi \in C^1$. Since $\hat{\rho}$ is a classical solution of (4.4) for all $s \notin \{c_1, 2\sqrt{r_1}\}$, we need only consider $s_0 = c_1$ or $s_0 = 2\sqrt{r_1}$. Suppose $s_0 = c_1$, then

$$\rho - s_0\phi' + |\phi'|^2 + R^* = \left(\frac{c_1^2}{4} - r_1\right) - c_1\phi' + |\phi'|^2 + r_1 = \left(\phi' - \frac{c_1}{2}\right)^2 \geq 0 \quad \text{at } s = s_0.$$

Suppose $s_0 = 2\sqrt{r_1}$, then

$$\rho - s_0\phi' + |\phi'|^2 + R^* = 0 - (2\sqrt{r_1})\phi' + |\phi'|^2 + r_1 = (\phi' - \sqrt{r_1})^2 \geq 0 \quad \text{at } s = s_0.$$

This verifies that $\hat{\rho}$ is a viscosity supersolution of (4.4). This completes the proof that the unique viscosity solution $\hat{\rho}$ as guaranteed by Lemma 4.4 is given by the explicit formula (5.10) for the first representative Case 2(c).

We omit the verification of the other cases since they are analogous. □

Corollary 5.2. *Suppose either that (1) $r_1 < r_2$, or that (2) $r_1 > r_2$ and $c_1 \notin (2\sqrt{r_1}, 2\sqrt{r_2}]$. Then there exists $\delta_0 > 0$ such that for each $\eta > 0$,*

$$\liminf_{t \rightarrow \infty} \inf_{|x| < (\sigma - \eta)t} u(t, x) \geq \delta_0 \quad \text{for each } \eta > 0 \text{ small enough,}$$

where $\sigma = \sigma(c_1; r_1, r_2)$ is given in (1.10). In particular, we have

$$\underline{c}_* \geq \sigma(c_1; r_1, r_2). \tag{5.11}$$

Proof. Observe that

$$u_t \geq u_{xx} + u(\delta' - u) \quad \text{for } (t, x) \in (0, \infty) \times \mathbb{R}$$

where $\delta' = \beta \inf_{s \in \mathbb{R}} a(s) - 1 > 0$, thanks to (H1). Since u has compactly supported initial data, it follows from standard theory that the spreading speed of u is bounded from below by $2\sqrt{\delta'}$, i.e.

$$\liminf_{t \rightarrow \infty} \inf_{|x| < 3\sqrt{\delta'}t/2} u(t, x) \geq \delta_0. \tag{5.12}$$

For given $c_1, r_1, r_2 > 0$ satisfying any of the cases 1(a)-(c) and 2(a)-(c), Lemma 5.1 says that the unique solution $\hat{\rho}$ guaranteed in Lemma 4.4 is given as in Subsection 5.1. If we define

$$\hat{s} := \sup\{s \geq 0 : \hat{\rho}(s) = 0\},$$

then it is easy to see that $\hat{s} = \sigma(c_1; r_1, r_2)$, where the latter is given in (1.10). By Lemmas 4.5 and 4.6, $0 \leq \rho^*(s) \leq \hat{\rho}(s)$ for $s \geq 0$. Thus, $\rho^*(s) = 0$ for all $s \in [0, \sigma]$, where $\sigma = \sigma(c_1; r_1, r_2)$.

It follows from Lemma 4.2 that

$$\liminf_{t \rightarrow \infty} \inf_{\eta t < x < (\sigma - \eta)t} u(t, x) \geq \delta_0 \quad \text{for each } \eta > 0 \text{ small enough,} \tag{5.13}$$

and a similar statement holds for $x < 0$. The desired result follows by combining with (5.12). □

Proof of Theorems 1.1 and Theorem 1.2. We recall that $\underline{c}_* \leq \bar{c}_*$, by construction. On the other hand, by Proposition 2.2, $\bar{c}_* \leq \sigma_1(c_1; r_1, r_2)$, and by Corollary 5.2, $\underline{c}_* \geq \sigma_1(c_1; r_1, r_2)$. It follows that $\underline{c}_* = \bar{c}_* = \sigma_1(c_1; r_1, r_2)$, so that the spreading speed of u is given by $c^* = \sigma_1(c_1; r_1, r_2)$. □

6. CONVERGENCE TO HOMOGENEOUS STATE

In this section, we apply the previous spreading result to characterize the long-time behavior of solutions of (1.3) in the moving frame where the predator persists.

Having established the existence of the spreading speed c^* , and recalling Lemma 3.1, it follows that $(u, v) \rightarrow (0, 1)$ locally uniformly in any moving frames with speed above c^* , and that u persists in any moving frames with speed below c^* . In the following we discuss the asymptotic behavior of the solutions in the latter case. Define, for $i = 1, 2$,

$$u_i = \frac{a_i - 1}{1 + a_i b}, \quad v_i = \frac{1 + b}{1 + a_i b} \quad (6.1)$$

where $a_1 := a(-\infty)$ and $a_2 := a(+\infty)$. Then (u_i, v_i) is the unique positive root of the algebraic system

$$u(-1 - u + a_i v) = 0 = rv(1 - v - bu).$$

Then Theorem 1.3 can be restated as follows.

Theorem 6.1. *Let (u, v) be the solution of (1.3), where $(u_0, v_0) \in C^2(\mathbb{R})$ satisfies (IC).*

(a) *Suppose $c_1 \geq c^*$. For any $\eta \in (0, c^*/2)$,*

$$\lim_{t \rightarrow \infty} \sup_{|x| < (c^* - \eta)t} \|(u, v) - (u_1, v_1)\| = 0. \quad (6.2)$$

where $\|\cdot\|$ denotes the Euclidean norm in \mathbb{R}^2 .

(b) *Suppose $c_1 < c^*$. For any $\eta \in (0, c_1/2)$,*

$$\lim_{t \rightarrow \infty} \sup_{|x| < (c_1 - \eta)t} \|(u, v) - (u_1, v_1)\| = 0, \quad (6.3)$$

and for any $\eta \in (0, (c^* - c_1)/2)$,

$$\lim_{t \rightarrow \infty} \sup_{(c_1 + \eta)t < x < (c^* - \eta)t} \|(u, v) - (u_2, v_2)\| = 0, \quad (6.4)$$

where (u_i, v_i) is defined in (6.1).

Proof of Theorem 6.1. Denote $c^* = \sigma(c_1; r_1, r_2)$. Thanks to Corollary 5.2, for each $\eta > 0$, there exists $T = T(\eta) > 0$ and $\delta_0 > 0$ such that

$$u(t, x) \geq \delta_0 \quad \text{for } t \geq T, \quad -\eta t \leq x \leq (c^* - \eta/2)t. \quad (6.5)$$

Suppose for a contradiction that there exists $\epsilon_0, \eta > 0$ and a sequence (t_k, x_k) with $t_k \rightarrow \infty$ and $0 \leq x_k < (c^* - \eta)t_k$ (the case for $x_k \leq 0$ is similar), such that $\|(u, v)(t_k, x_k) - (\tilde{u}, \tilde{v})\| > \epsilon_0$ for all $k \geq 1$, where

$$(\tilde{u}, \tilde{v}) = \begin{cases} (u_2, v_2) & \text{if } c_1 < c^* \quad \text{and } x_k \in ((c_1 + \eta)t_k, (c^* - \eta)t_k), \\ (u_1, v_1) & \text{otherwise.} \end{cases} \quad (6.6)$$

Let

$$u_k(t, x) := u(t + t_k, x + x_k) \quad \text{and} \quad v_k(t, x) := v(t + t_k, x + x_k),$$

then

$$u_k(t, x) \geq \delta_0 \quad \text{in } \Omega_k,$$

where

$$\Omega_k = \{(t, x) : t + t_k \geq T, \quad -\eta(t + t_k) \leq x + x_k \leq (c^* - \eta/2)(t + t_k)\}.$$

Claim. $\Omega_k \rightarrow \mathbb{R}^2$, i.e., for any compact subset $K \subset \mathbb{R}^2$ there exists $k_1 > 1$ such that

$$K \subset \Omega_k \quad \text{for all } k \geq k_1.$$

Indeed, given K , choose $R > 0$ such that $K \subset [-R, R] \times [-R, R]$. Then for $k \gg 1$, we have

$$\begin{aligned} K &\subset [-R, R]^2 \\ &\subseteq \{(t, x) : |t| \leq R, \quad \eta(R - t_k) \leq x \leq -(c^* - \eta/2)R + \eta t_k/2\} \\ &\subseteq \{(t, x) : |t| \leq R, \quad -\eta(t + t_k) \leq x \leq (c^* - \eta/2)t + \eta t_k/2\} \\ &\subseteq \{(t, x) : t + t_k \geq T, \quad -\eta(t + t_k) - x_k \leq x \leq (c^* - \eta/2)t - x_k + (c^* - \eta/2)t_k\} \\ &= \Omega_k. \end{aligned}$$

This proves the claim.

It follows from the claim and Lemma 2.1 that there exists constants $0 < \delta_1 < 1$ independent of k such that

$$\delta_1 \leq u_k \leq \frac{1}{\delta_1} \quad \text{and} \quad \delta_1 \leq v_k \leq \frac{1}{\delta_1} \quad \text{in } \Omega_k \quad (6.7)$$

Using the above L^∞ bounds and parabolic L^p estimates we may deduce, re-labelling a subsequence if necessary, that (u_k, v_k) converges weakly in $W_{loc}^{2,1,p}(\mathbb{R}^2)$ (and strongly in $C_{loc}^{1+\alpha, (1+\alpha)/2}(\mathbb{R}^2)$ thanks to Sobolev embedding) to an entire solution (u_∞, v_∞) of the system

$$\begin{cases} u_t &= u_{xx} + u(-1 - u + \tilde{a}v) \\ v_t &= dv_{xx} + rv(1 - v - bu), \end{cases}$$

where $\tilde{a} = a_2$ if $c_1 < c^*$ and $x_k \in ((c_1 + \eta)t_k, (c^* - \eta)t_k)$, and $\tilde{a} = a_1$ otherwise. Moreover, (6.7) also implies that

$$\delta_1 \leq u_\infty \leq \frac{1}{\delta_1} \quad \text{and} \quad \delta_1 \leq v_\infty \leq \frac{1}{\delta_1} \quad \text{in } \mathbb{R}^2. \quad (6.8)$$

Having established the positive upper and lower bounds for (u_∞, v_∞) on \mathbb{R}^2 , one can then repeat a standard argument via Lyapunov functional (see the proof of Lemma 4.1 in [21]) that (u_∞, v_∞) is identically equal to the homogeneous steady state (\tilde{u}, \tilde{v}) given in (6.6), i.e., $(u_k, v_k) \rightarrow (\tilde{u}, \tilde{v})$ in $C_{loc}(\mathbb{R}^2)$. This in particular implies

$$(u, v)(t_k, x_k) = (u_k, v_k)(0, 0) \rightarrow (\tilde{u}, \tilde{v}) \quad \text{as } k \rightarrow \infty.$$

But this is a contradiction, which completes the proof. \square

APPENDIX A. COMPARISON PRINCIPLE

Recall the Hamilton-Jacobi equation

$$\min\{\rho - s\rho' + |\rho'|^2 + \tilde{R}(s), \rho\} = 0 \quad \text{for } s \in (0, \infty). \quad (\text{A.1})$$

We prove a comparison principle for (A.1) for discontinuous $\tilde{R} : \mathbb{R} \rightarrow \mathbb{R}$ that is locally monotone [14].

Definition A.1. A function $h : \mathbb{R} \rightarrow \mathbb{R}$ is locally monotone if for every $s_0 \in \mathbb{R}$, either

$$\lim_{\delta \rightarrow 0} \inf_{\substack{|s_i - s_0| < \delta \\ s_1 > s_2}} (h(s_1) - h(s_2)) \geq 0 \quad \text{or} \quad \lim_{\delta \rightarrow 0} \sup_{\substack{|s_i - s_0| < \delta \\ s_1 > s_2}} (h(s_1) - h(s_2)) \leq 0.$$

The assumptions on \tilde{R} are stated precisely as follows.

(B1) $\tilde{R}(s)$ is locally monotone;

(B2) $\tilde{R}^*(s) = \tilde{R}_*(s)$ almost everywhere, and $\inf_{s>0} R(s) > 0$, where \tilde{R}^* and \tilde{R}_* are defined by

$$\tilde{R}^*(s) = \limsup_{s' \rightarrow s} \tilde{R}(s') \quad \text{and} \quad \tilde{R}_*(s) = \liminf_{s' \rightarrow s} \tilde{R}(s').$$

Next, note that the specific form of the Hamilton-Jacobi equation (A.1) and hypotheses (B1)-(B2) imply the hypotheses (H1)-(H6) in [41]. The following two results are due to [41].

Lemma A.1. *Suppose $\tilde{R}(s)$ satisfies (B1)-(B2). Let $\bar{\rho}$ and $\underline{\rho}$ be non-negative viscosity super- and subsolutions, respectively, of (A.1) such that*

$$\underline{\rho}(0) \leq \bar{\rho}(0) \quad \text{and} \quad \lim_{s \rightarrow \infty} \frac{\underline{\rho}(s)}{s} \leq \lim_{s \rightarrow \infty} \frac{\bar{\rho}(s)}{s}. \quad (\text{A.2})$$

Then $\underline{\rho} \leq \bar{\rho}$ in $(0, \infty)$.

Proof. This is a consequence of [41, Proposition 2.11]. \square

Corollary A.2. *Let $\tilde{R} : \mathbb{R} \rightarrow \mathbb{R}$ satisfy the assumptions (B1)-(B2). Then there exists a unique viscosity solution $\hat{\rho}$ to (A.1) satisfying the boundary conditions*

$$\hat{\rho}(0) = 0 \quad \text{and} \quad \lim_{s \rightarrow \infty} \frac{\hat{\rho}(s)}{s} = \infty. \quad (\text{A.3})$$

Moreover, $\hat{\rho}$ is nondecreasing.

Proof. This is a consequence of [41, Proposition 1.7(b) and Lemma 2.9] \square

REFERENCES

- [1] I. Ahn, W. Choi, A. Ducrot, and J.-S. Guo, *Spreading dynamics for a three species predator-prey system with two preys in a shifting environment*, J. Dyn. Diff. Equat. (2022). <https://doi.org/10.1007/s10884-022-10237-z>.
- [2] D. G. Aronson and H. F. Weinberger, *Nonlinear diffusion in population genetics, combustion, and nerve pulse propagation*, in Partial differential equations and related topics (Program, Tulane Univ., New Orleans, LA., 1974), Lecture Notes in Math., Vol. 446, Springer, Berlin-New York, 1975, pp. 5–49.
- [3] D. G. Aronson and H. F. Weinberger, *Multidimensional nonlinear diffusion arising in population genetics*, Adv. Math. **30**(1978), 33–76.
- [4] R. B. Aronson, K. E. Smith, S. C. Vos, J. B. McClintock, M. O. Amsler, P.-O. Moksnes, D. S. Ellis, J. Kaeli, H. Singh, J. W. Bailey, J. C. Schiferl, R. van Woesik, M. A. Martin, B. V. Steffel, M. E. Deal, S. M. Lazarus, J. N. Havenhand, R. Swalethorp, S. Kjellerup, S. Thatje, *No barrier to emergence of bathyal king crabs on the antarctic shelf*, Proceedings of the National Academy of Sciences, **112**(2015), 12997–13002.
- [5] M. Bardi and I. Capuzzo-Dolcetta, *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*, Systems & Control: Foundations & Applications, Birkhäuser Boston, Inc., Boston, MA, 1997. With appendices by Maurizio Falcone and Pierpaolo Soravia.
- [6] G. Barles, *An introduction to the theory of viscosity solutions for first-order Hamilton-Jacobi equations and applications*, in Hamilton-Jacobi equations: approximations, numerical analysis and applications, vol. 2074 of Lecture Notes in Math., Springer, Heidelberg, 2013, pp. 49–109.
- [7] G. Barles and B. Perthame, *Discontinuous solutions of deterministic optimal stopping time problems*, RAIRO Modél. Math. Anal. Numér. **21**(1987), 557–579.
- [8] H. Berestycki, O. Diekmann, C. J. Nagelkerke, and P. A. Zegeling, *Can a species keep pace with a shifting climate?*, Bull. Math. Biol. **71**(2009), 399–429.
- [9] H. Berestycki and J. Fang, *Forced waves of the Fisher-KPP equation in a shifting environment*, J. Differential Equations, **264**(2018), 2157–2183.
- [10] H. Berestycki, J.-M. Roquejoffre, and L. Rossi, *The shape of expansion induced by a line with fast diffusion in Fisher-KPP equations*, Comm. Math. Phys. **343**(2016), 207–232.
- [11] B. A. Bradley, D. S. Wilcove, and M. Oppenheimer, *Climate change increases risk of plant invasion in the eastern united states*, Biological Invasions, **12**(2010), 1855–1872.
- [12] M. Bramson, *Convergence of solutions of the Kolmogorov equation to travelling waves*, Mem. Amer. Math. Soc. **44**(1983), pp. iv+190.
- [13] L. B. Buckley and J. G. Kingsolver, *Functional and phylogenetic approaches to forecasting species' responses to climate change*, Annual Review of Ecology, Evolution, and Systematics, **43**(2012), 205–226.
- [14] X. Chen and B. Hu, *Viscosity solutions of discontinuous Hamilton-Jacobi equations*, Interfaces Free Bound. **10**(2008), 339–359.
- [15] W. Choi, T. Giletti, and J.-S. Guo, *Persistence of species in a predator-prey system with climate change and either nonlocal or local dispersal*, J. Differential Equations **302**(2021), 807–853.
- [16] U. Daugaard, O. L. Petchev, and F. Pennkamp, *Warming can destabilize predator-prey interactions by shifting the functional response from type iii to type ii*, J. Anim. Ecol., **88**(2019), 1575–1586.

- [17] C. A. Deutsch, J. J. Tewksbury, R. B. Huey, K. S. Sheldon, C. K. Ghalambor, D. C. Haak, and P. R. Martin, *Impacts of climate warming on terrestrial ectotherms across latitude*, Proceedings of the National Academy of Sciences **105**(2008), 6668–6672.
- [18] F.-D. Dong, B. Li, and W.-T. Li, *Forced waves in a Lotka-Volterra competition-diffusion model with a shifting habitat*, J. Differential Equations **276**(2021), 433–459.
- [19] A. Ducrot, *Convergence to generalized transition waves for some Holling-Tanner prey-predator reaction-diffusion system*, J. Math. Pures Appl. **100**(2013), 1–15.
- [20] A. Ducrot, *Spatial propagation for a two component reaction-diffusion system arising in population dynamics*, J. Differential Equations **260**(2016), 8316–8357.
- [21] A. Ducrot, T. Giletti, J.-S. Guo, and M. Shimojo, *Asymptotic spreading speeds for a predator-prey system with two predators and one prey*, Nonlinearity **34**(2021), 669–704.
- [22] A. Ducrot, T. Giletti, and H. Matano, *Spreading speeds for multidimensional reaction-diffusion systems of the prey-predator type*, Calc. Var. Partial Differential Equations **58**(2019): 137.
- [23] A. Ducrot, J.-S. Guo, G. Lin, and S. Pan, *The spreading speed and the minimal wave speed of a predator-prey system with nonlocal dispersal*, Z. Angew. Math. Phys. **70**(2019): 146
- [24] S. R. Dunbar, *Travelling wave solutions of diffusive Lotka-Volterra equations*, J. Math. Biol. **17**(1983), 11–32.
- [25] L. C. Evans and P. E. Souganidis, *A PDE approach to geometric optics for certain semilinear parabolic equations*, Indiana Univ. Math. J. **38**(1989), 141–172.
- [26] J. Fang, Y. Lou, and J. Wu, *Can pathogen spread keep pace with its host invasion?*, SIAM J. Appl. Math. **76** (2016), 1633–1657.
- [27] R. A. FISHER, *The wave of advance of advantageous genes*, Annals of Eugenics, **7**(1937), 355–369.
- [28] R. Gardner and C. K. R. T. Jones, *Stability of travelling wave solutions of diffusive predator-prey systems*, Trans. Amer. Math. Soc. **327**(1991), 465–524.
- [29] R. A. Gardner, *Existence of travelling wave solutions of predator-prey systems via the connection index*, SIAM J. Appl. Math. **44**(1984), 56–79.
- [30] S. E. Gilman, M. C. Urban, J. Tewksbury, G. W. Gilchrist, and R. D. Holt, *A framework for community interactions under climate change*, Trends in Ecology & Evolution **25**(2010), 325–331.
- [31] L. Girardin and K.-Y. Lam, *Invasion of open space by two competitors: spreading properties of monostable two-species competition-diffusion systems*, Proc. Lond. Math. Soc. **119**(2019), 1279–1335.
- [32] J.-S. Guo, M. Shimojo, and C.-C. Wu, *Spreading dynamics for a predator-prey system with two predators and one prey in a shifting habitat*, Discrete Contin. Dyn. Syst. Ser. B **28**(2023), 6126–6141.
- [33] F. Hamel and G. Nadin, *Spreading properties and complex dynamics for monostable reaction-diffusion equations*, Comm. Partial Differential Equations, **37**(2012), 511–537.
- [34] R. J. Hobbs and H. A. Mooney, *Invasive species in a changing world*, Island Press, 2000.
- [35] M. Holzer and A. Scheel, *Accelerated fronts in a two-stage invasion process*, SIAM J. Math. Anal. **46** (2014), 397–427.
- [36] J. Huang, G. Lu, and S. Ruan, *Existence of traveling wave solutions in a diffusive predator-prey model*, J. Math. Biol. **46**(2003), 132–152.
- [37] R. B. Huey and R. Stevenson, *Integrating thermal physiology and ecology of ectotherms: a discussion of approaches*, American Zoologist **19**(1979), 357–366.
- [38] H. Ishii, *Hamilton-Jacobi equations with discontinuous Hamiltonians on arbitrary open sets*, Bull. Fac. Sci. Engrg. Chuo Univ. **28**(1985), 33–77.
- [39] A. Kolmogorov, I. Petrovskii, and N. Piscounov, *Étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique*, Moscow Univ. Bull. Ser. Boarding School Section. HAS, **1**(1937), p. 126.
- [40] K.-Y. Lam and Y. Lou, *Introduction to Reaction-Diffusion Equations: Theory and Applications to Spatial Ecology and Evolutionary Biology*, Lecture Notes on Mathematical Modelling in the Life Sciences, Springer, Cham, 2022.
- [41] K.-Y. Lam and X. Yu, *Asymptotic spreading of KPP reactive fronts in heterogeneous shifting environments*, J. Math. Pures Appl. **167**(2022), 1–47.
- [42] B. Lang, R. B. Ehnes, U. Brose, and B. C. Rall, *Temperature and consumer type dependencies of energy flows in natural communities*, Oikos, **126**(2017), 1717–1725.
- [43] K.-S. Lau, *On the nonlinear diffusion equation of Kolmogorov, Petrovsky, and Piscounov*, J. Differential Equations **59**(1985), 44–70.
- [44] M. A. Lewis, B. Li, and H. F. Weinberger, *Spreading speed and linear determinacy for two-species competition models*, J. Math. Biol. **45**(2002), 219–233.

- [45] B. Li, S. Bewick, J. Shang, and W. F. Fagan, *Persistence and spread of a species with a shifting habitat edge*, SIAM J. Appl. Math. **74**(2014), 1397–1417.
- [46] B. Li, H. F. Weinberger, and M. A. Lewis, *Spreading speeds as slowest wave speeds for cooperative systems*, Math. Biosci. **196**(2005), 82–98.
- [47] X. Liang and X.-Q. Zhao, *Asymptotic speeds of spread and traveling waves for monotone semiflows with applications*, Comm. Pure Appl. Math. **60**(2007), 1–40.
- [48] S. D. Ling, *Range expansion of a habitat-modifying species leads to loss of taxonomic diversity: a new and impoverished reef state*, Oecologia **156**(2008), 883–894.
- [49] Q. Liu, S. Liu, and K.-Y. Lam, *Asymptotic spreading of interacting species with multiple fronts I: a geometric optics approach*, Discrete Contin. Dyn. Syst. **40**(2020), 3683–3714.
- [50] Q. Liu, S. Liu, and K.-Y. Lam, *Stacked invasion waves in a competition-diffusion model with three species*, J. Differential Equations **271**(2021), 665–718.
- [51] K. Mischaikow and J. F. Reineck, *Travelling waves in predator-prey systems*, SIAM J. Math. Anal. **24**(1993), 1179–1214.
- [52] S. Pan, *Asymptotic spreading in a Lotka-Volterra predator-prey system*, J. Math. Anal. Appl. **407**(2013), 230–236.
- [53] S. Pan, *Invasion speed of a predator-prey system*, Appl. Math. Lett. **74**(2017), 46–51.
- [54] G. T. Pecl, M. B. Araújo, J. D. Bell, J. Blanchard, T. C. Bonebrake, I.-C. Chen, T. D. Clark, R. K. Colwell, F. Danielsen, B. Evengård, L. Falconi, S. Ferrier, S. Frusher, R. A. Garcia, R. B. Griffis, A. J. Hobday, C. Janion-Scheepers, M. A. Jarzyna, S. Jennings, J. Lenoir, H. I. Linnetved, V. Y. Martin, P. C. McCormack, J. McDonald, N. J. Mitchell, T. Mustonen, J. M. Pandolfi, N. Pettorelli, E. Popova, S. A. Robinson, B. R. Scheffers, J. D. Shaw, C. J. B. Sorte, J. M. Strugnell, J. M. Sunday, M.-N. Tuanmu, A. Vergés, C. Villanueva, T. Wernberg, E. Wapstra, and S. E. Williams, *Biodiversity redistribution under climate change: Impacts on ecosystems and human well-being*, Science **355**(6332)(2017):aa9214. <https://www.science.org/doi/10.1126/science.aa9214>.
- [55] A. B. Potapov and M. A. Lewis, *Climate and competition: the effect of moving range boundaries on habitat invasibility*, Bull. Math. Biol. **66**(2004), 975–1008.
- [56] M. H. Protter and H. F. Weinberger, *Maximum principles in differential equations*, Springer-Verlag, New York, 1984.
- [57] N. Shigesada and K. Kawasaki, *Biological Invasions: Theory and Practice*, Oxford Series in Ecology and Evolution, Oxford University Press Inc., New York, 1997.
- [58] J. Smoller, *Shock waves and reaction-diffusion equations*, vol. 258 of Grundlehren der Mathematischen Wissenschaften, Springer-Verlag, New York-Berlin, 1983.
- [59] C. J. B. Sorte, S. L. Williams, and J. T. Carlton, *Marine range shifts and species introductions: Comparative spread rates and community impacts*, Glob. Ecol. Biogeogr. **19**(2010), 303–316.
- [60] A. Vergés, C. Doropoulos, H. A. Malcolm, M. Skye, M. Garcia-Pizá, E. M. Marzinelli, A. H. Campbell, E. Ballesteros, A. S. Hoey, A. Vila-Concejo, Y.-M. Bozec, P. D. Steinberg, *Long-term empirical evidence of ocean warming leading to tropicalization of fish communities, increased herbivory, and loss of kelp*, Proceedings of the National Academy of Sciences **113**(2016), 13791–13796.
- [61] P. D. Wallingford, T. L. Morelli, J. M. Allen, E. M. Beaury, D. M. Blumenthal, B. A. Bradley, J. S. Dukes, R. Early, E. J. Fusco, D. E. Goldberg, I. Ibanez, B. B. Laginhas, M. Vila, and C. J. B. Sorte, *Adjusting the lens of invasion biology to focus on the impacts of climate-driven range shifts*, Nature Climate Change **10**(2020), 398–405.
- [62] G.-R. Walther, A. Roques, P. E. Hulme, M. T. Sykes, P. Pyšek, I. Kühn, M. Zobel, S. Bacher, Z. Botta-Dukát, H. Bugmann, B. Czucz, J. Dauber, T. Hickler, V. Jarošík, M. Kenis, S. Klotz, D. Minchin, M. Moora, W. Nentwig, J. Ott, V. E. Panov, B. Reineking, C. Robinet, V. Semchenko, W. Solarz, W. Thuiller, M. Vilà, K. Vohland, J. Settele, *Alien species in a warmer world: risks and opportunities*, Trends in Ecology & Evolution **24**(2009), 686–693.
- [63] H. Wang, *Spreading speeds and traveling waves for non-cooperative reaction-diffusion systems*, J. Nonlinear Sci. **21**(2011), 747–783.
- [64] J.-B. Wang, W.-T. Li, F.-D. Dong, and S.-X. Qiao, *Recent developments on spatial propagation for diffusion equations in shifting environments*, Discrete Contin. Dyn. Syst. Ser. B **27**(2022), 5101–5127.
- [65] H. F. Weinberger, M. A. Lewis, and B. Li, *Analysis of linear determinacy for spread in cooperative models*, J. Math. Biol. **45**(2002), 183–218.
- [66] S. R. Weiskopf, M. A. Rubenstein, L. G. Crozier, S. Gaichas, R. Griffis, J. E. Halofsky, K. J. Hyde, T. L. Morelli, J. T. Morissette, R. C. Muñoz, A. J. Pershing, D. L. Peterson, R. Poudel, M. D. Staudinger, A. E. Sutton-Grier, L. Thompson, J. Vose, J. F. Weltzin, K. P. Whyte, *Climate change effects on biodiversity, ecosystems, ecosystem services, and natural resource management in the United States*, Science of the Total Environment **733**(2020): 137782.

- [67] C.-C. Wu, *The spreading speed for a predator-prey model with one predator and two preys*, Appl. Math. Lett. **91**(2019), 9–14.
- [68] S.-L. Wu, L. Pang, and S. Ruan, *Propagation dynamics in periodic predator-prey systems with nonlocal dispersal*, J. Math. Pures Appl. **170**(2023), 57–95.
- [69] X. Wu, Y. Lu, S. Zhou, L. Chen, and B. Xu, *Impact of climate change on human infectious diseases: Empirical evidence and human adaptation*, Environment International **86**(2016), 14–23.
- [70] F. Yang, W. Li, and R. Wang, *Invasion waves for a nonlocal dispersal predator-prey model with two predators and one prey*, Commun. Pure Appl. Anal., 20 (2021), pp. 4083–4105.
- [71] Z. Zhang, W. Wang, and J. Yang, *Persistence versus extinction for two competing species under a climate change*, Nonlinear Anal. Model. Control **22**(2017), 285–302.

CORRESPONDING AUTHOR, DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OH 43220, USA
Email address: lam.184@osu.edu

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY, COLUMBUS, OHIO 43220, USA
Email address: lee.7355@buckeyemail.osu.edu