

Improving quality parameters of spinach by adjusting light spectra under moderate water deprivation conditions

Tahooria B. ZARGAR, Oqba BASAL*, Szilvia VERES

*University of Debrecen, Faculty of Agricultural and Food Sciences and Environmental Management,
Hungary; tahooria@mailbox.unideb.hu; oqba@agr.unideb.hu (*corresponding author); szveres@agr.unideb.hu*

Abstract

Spinach (*Spinacia oleracea* L.) is a highly valuable leafy vegetable, abundant in vitamins, minerals, and antioxidants, offering various health benefits such as enhancing cardiovascular health, reducing inflammation, and aiding digestion. Consequently, it is crucial to effectively cultivate and maintain spinach's quality. Light plays a pivotal role in the growth and development of plants, including spinach, where different light qualities can influence its morpho-physiological traits and overall quality. To explore this, an experiment was conducted, using three distinct LED lighting sets for spinach cultivation. Light 1 emphasized blue light, Light 2 served as the standard white light control, and Light 3 focused more on red light with occasional brief UV-C flashes. Additionally, mild water deprivation was induced using 2.5% polyethylene glycol (PEG). The results revealed that using a high red-to-blue light ratio with intermittent UV-C radiation significantly reduced various growth parameters of spinach, such as root length, shoot length, root volume, fresh and dry root and shoot weight, as well as total and relative chlorophyll contents, when compared to the control group. Furthermore, water deprivation had a negative impact on spinach's growth, affecting shoot and root length, and fresh and dry weight in all light qualities, proving to be fatal under Light 3 conditions. Therefore, it is essential to carefully select appropriate light qualities throughout the plant's life cycle to enhance the quality of spinach, especially when mild water deprivation is involved. Opting for a higher blue-to-red light ratio was found to be somewhat beneficial in improving the overall quality of spinach.

Keywords: abiotic stress; green leafy vegetables; light quality; pigments; water deprivation

Introduction

Spinach (*Spinacia oleracea* L. cv. 'Matador') is one of the economic vegetables with a minimum growth cycle and is grown annually (Ribera *et al.*, 2020). The increasing consumption of fresh (in salads) and processed spinach demonstrates its rising popularity worldwide. When fresh, steamed or barely cooked, spinach has a high biological value and is particularly abundant in antioxidants (Ribera *et al.*, 2020). It has high levels of fibers, minerals (Mn, Fe, K, Mg, Ca, Se), and vitamins (A, E, K, C, B2, B6, B9, folic acid) (Roberts and Moreau, 2016). Spinach is also known for its anti-inflammatory, antitumoral, anti-obesity and lipid-lowering effects. Consuming more antioxidant-rich fruits and vegetables, like spinach, may halt the deterioration of antioxidant defense mechanisms that occur with ageing (Hassan *et al.*, 2022). Spinach crops must be improved in quality

Received: 31 Jul 2023. Received in revised form: 20 Sep 2023. Accepted: 27 Sep 2023. Published online: 17 Nov 2023.

From Volume 49, Issue 1, 2021, Notulae Botanicae Horti Agrobotanici Cluj-Napoca journal uses article numbers in place of the traditional method of continuous pagination through the volume. The journal will continue to appear quarterly, as before, with four annual numbers.

and yield by finding new approaches. Moreover, efforts to reduce the potential environmental impact on the production and consumption of spinach are essential.

Severe abiotic stresses, such as drought stress, can negatively affect plant growth. However, mild stress can be intentionally induced to enhance antioxidant compounds without negatively impacting productivity in the edible part of the plant and promote plant adaptation to stressful conditions (Cogo *et al.*, 2011; Paim *et al.*, 2020). Additionally, there is evidence that plants subjected to minor abiotic stressors are more resilient to subsequent stresses such as those brought on by the postharvest storage procedure. The metabolic memory created by the original stimulation would be the reason for this effect (Paim *et al.*, 2020). Priming is the deliberate application of mild stresses to withstand stress in the future (Sorrentino *et al.*, 2020). The development of technical approaches to lower water resource use in food production has therefore been the subject of several initiatives (Akıncı and Lösel, 2012). So, the application of modest water deprivation may be worthwhile to stimulate the metabolic memory to withstand resilient conditions in the future. Spinach quality may be improved by mild water deprivation through a number of physiological and biochemical processes. Spinach plants adapt in ways that enhance their nutritional value, flavor, and general quality when they experience a minor water shortage. Here are several instances of how mild drought stress may affect spinach quality, mild water deprivation can lead to increased antioxidant content, enhanced phytochemical content, increased mineral uptake, improved leaf texture and taste, and changes in leaf morphology and composition. Antioxidants help plants cope with oxidative stress and can have positive effects on human health when consumed (Barros *et al.*, 2016).

Among various environmental factors, light is one of the most significant abiotic factors that regulate plant growth, photosynthesis, survival, and adaptation. Light quality affects plant development, photomorphogenesis, and tissue composition; in addition to light quantity, factors like intensity and length (photoperiod) also influence plant productivity (Ouzounis *et al.*, 2015). Red and blue light, for instance, are the colors most effectively used for photosynthesis, and red light promotes the development of the photosynthetic apparatus (Paradiso *et al.*, 2011). Far-red light encourages blooming in long-day plants, whereas blue light impacts stomatal opening, plant height, and chlorophyll biosynthesis. The red/far red ratio controls stem and branching length, leaf area, and reproduction (Zheng *et al.*, 2019). Seedlings grown under a combination of green, red, and blue lights were taller than those grown under monochromatic red or blue lights in *Arabidopsis thaliana* (Folta, 2004) and increased fresh and dry weight of tomato plant and fruit in combination with red and blue light is recorded (Kaiser *et al.*, 2019). By raising the expression of synthesis pathway genes, blue and UV wavelengths have been demonstrated to promote the accumulation of bioactive chemicals in plant tissues (Hasan and Bae, 2017). Bian *et al.* (2014) highlighted the positive effects of blue, UV-A, and UV-B on the formation of phenolic compounds in general and anthocyanins, as well as the positive impacts of blue, red, and UV-B on carotenoids in a variety of vegetables. Researchers have shown that UV-C-induced effects generally depend on radiation dose and developmental stage (Poiroux *et al.*, 2010). There is mounting evidence that plants may benefit from UV-C light (Urban *et al.*, 2016). These may result in changes to various primary and secondary metabolites, such as chlorophyll, carotenoids, phenolics, and glucosinolates (Katerova *et al.*, 2012; Vandenbussche *et al.*, 2018; Mewis *et al.*, 2012; Heinze *et al.*, 2018). When exposed to UV radiation, flavonoid accumulation was seen in epidermal cells, leaf hairs, and leaf wax (Harborne and Williams, 2000). UV light greatly stimulates the production of carotenoids such as violaxanthin, neoxanthin, antheraxanthin, luteoxanthin, and lutein (Molnar *et al.*, 2009). Exposure to UV-C radiation increased carotenoids in tomatoes (Bravo *et al.*, 2013). So, it becomes crucial to know what light quality leads to which response in plants. During the daytime, sunlight is the only source of different light qualities, but plants do not need all of them for proper growth. We need to understand and find what kinds of light in plants lead to the proper growth, leaf extension or stem elongation and yield. In greenhouse horticulture, this information could be used to improve crop yield, quality, and production schedules (Paradiso and Proietti, 2022).

The employment of advanced plant cultivation technologies in a controlled environment to assure good crop yield, even in the face of harsh external conditions or in high-density cultivation systems, has become more common in modern agriculture. Although some previous research was carried out to investigate the response of spinach to drought stress (Gilani *et al.*, 2020) and light intensity (Nguyen *et al.*, 2022), to our knowledge no research focused on whether different light qualities and mild water deprivation can affect the morpho-physiology of spinach plants. The present experiment was carried out to study how different qualities of light influenced the growth parameters (root and shoot length, fresh and dry root and shoot weight, root volume), photosynthetic pigments and MDA content of spinach in general and in combination with mild water deprivation.

Materials and Methods

Experimental location and conditions

The Department of Applied Plant Biology, Institute of Crop Sciences, University of Debrecen, Hungary, conducted this experiment in 2021. Garafarm Trade Kft. in Budapest provided the spinach seeds, which were surface sterilized with 6% [v/v] H₂O₂ for 20 minutes before being properly washed and rinsed with deionized water. Three replicates of the 2.5% concentration PEG 6000 solution (VWR International bvba Geldenaaksebaan, Leuven, Belgium) were utilized to compare it to the control treatment, which only employed the nutritional solution. The nutrient mixture included 0.7 mM K₂SO₄, 0.5 mM MgSO₄, 0.1 mM KH₂PO₄, 0.1 mM KCl, 0.5 μM MnSO₄, 0.5 μM ZnSO₄, 10 μM H₃BO₃, and 0.2 μM CuSO₄ along with 2.0 mM Ca (NO₃)₂. Additionally, iron was given in the form of 10⁻⁴ M Fe-EDTA (Marschner *et al.*, 1990). The seeds were grown geotropically at 24 °C between moist filter sheets. The humidity was preserved at 62.5% ± 0.5% while the room temperature was maintained at 24 °C ± 0.2 °C. After they developed two true leaves, seedlings of the same size were transplanted into plastic pots (1.7 liters) in a hydroponic environment. The racks on which pots were placed were equipped with LED lamp sets (Tungsram Ltd, Hungary) with 300 μmol m⁻² s⁻¹ and three different light qualities, Light 1 with higher percentage of blue light (Blue >>Red), Light 2 with normal white light taken as control and Light 3 with higher red light (Red > Blue >) along with flashes of UV-C once a week (Table 1).

Table 1. Percentage of different light qualities used in the experiment

LED Lamps	Light quality						Explanation
	Blue	Green	Red	Far-red	UV-C	Total PPF	
Light 1	20%	40%	66%	10%	0%	245 μmol/s	Blue >>Red
Light 2	35%	48%	15%	1%	0%	350 μmol/s	Normal White
Light 3	12%	4%	75%	10%	1%	308 μmol/s	Red > Blue > Flashes of UV-C radiations once a week for 5 minutes.

In the experiment, six pots were set up for each LED lamp set, with each pot containing four spinach plants. Among these pots, three were filled with a nutrient solution, and the other three were filled with a nutrient solution mixed with 2.5% polyethylene glycol (PEG) to induce mild water deprivation.

Throughout the experiment, the nutrient solution was replaced every three days until harvest. On the 25th day after sowing, two plants were harvested from each pot containing only the nutrient solution to examine the impact of various light qualities on spinach growth.

For further investigation, the remaining two plants from each pot (with only nutrient solution) were allowed to grow until the 35th day after sowing, serving as the control group to compare against the plants grown with the nutrient solution mixed with 2.5% PEG under different light qualities. These plants were

harvested on the 35th day (Full harvest) to analyze the effects of mild drought and different light qualities on spinach growth and development.

Growth parameters

Measurements were taken from three replicates to determine the length of roots and shoots. The first measurement was conducted on the 25th day after sowing, referred to as the first harvest, while the second measurement was taken on the 35th day after sowing, named as the full harvest. Root and shoot lengths in three replicates were measured using a ruler after both harvests. To assess the volume of the roots, the water displacement technique was employed, whereby the sample volume was determined by calculating the difference between the initial and final water levels. Fresh weights of both roots and shoots were measured using an electronic scale manufactured by Ohaus in Switzerland. To obtain the dry weights, three replicate samples from each treatment were dried in an oven set at 70 °C for four days, until a consistent weight was achieved.

Relative chlorophyll content

SPAD values were recorded using a SPAD-502Plus device manufactured by Konica Minolta in Japan to assess the relative chlorophyll content. Measurements were taken every other day over a period of ten days during the first harvest, and at two-day intervals during the full harvest. Three repetitions were performed for each treatment, with replicates taken from fully-grown leaves.

*Total chlorophyll content (Chl-*a*, Chl-*b*) and total carotenoids (Chl-*x+c*)*

Fresh leaf disks (0.05 g) were collected and transferred into plastic vials containing 5ml of N, N-Dimethylformamide for extraction of chlorophyll and carotenoids (Moran and Porath, 1980) and placed in the dark at 4 °C for three days. Chlorophyll and total carotenoid concentration in the extract was measured with V-VIS spectrophotometry (Metertech SP-830 PLUS, Taiwan) at three wavelengths; 480, 647 and 664 nm (Wellburn, 1994).

Malondialdehyde (MDA) Content

Trichloroacetic acid (TCA) was used as an extraction buffer to assess the MDA content of the lyophilized leaf samples (Christ, Germany) (Baryla *et al.*, 2000). 0.1 g of crushed leaf powder was homogenized in 1 ml of 0.1% (w/v) TCA. Following homogenization, the samples were centrifuged at 10,000 rpm for 10 min. and the supernatant was transferred to a tube containing 4 ml of 20% TCA and 0.5% thiobarbituric acid (TBA). The resulting mixture was heated for 30 min. at 95 °C before being promptly cooled on ice. The recovered material was centrifuged again at 10,000 rpm for 5 min. Absorbance was determined using a spectrophotometer at 532 nm (Metertech SP-830 PLUS, Taiwan) (Baryla *et al.*, 2000).

Antheraxanthin, lutein, and violaxanthin content

To 0.05 g of leaf powder, 1.5 ml of 80% acetone was added and stored inside the fridge (4 °C) for one day. Samples were centrifuged for 10 min. at 13000 rpm, and the supernatant was separated from the pellet. The supernatant was stored inside the fridge (4 °C), and 1 ml of 80% acetone was added to the pellet and stored in the fridge (4 °C) for one day. Again, the samples were centrifuged at 13000 rpm for 10 min. Supernatants obtained were mixed and the pellet was discarded. The method used to determine the Antheraxanthin, Lutein, and Violaxanthin content in the extract was HPLC (ABL[®]E-Jasco Hungary Ltd.) (Goodwin *et al.*, 1988), specifically using a Nucleosil C18 column. The eluents used were acetonitrile and water in a 9:1 ratio, with 0.01% triethylamine, as well as ethyl acetate. The HPLC-system used for the analysis was equipped with a UV/VIS detector manufactured by JASCO in Japan. To identify peaks in the chromatogram and calculate pigment contents, zeaxanthin was regularly injected as a standard compound during the analyses (Toth *et al.*, 2002).

Experimental design and statistical analysis

A randomized complete block design with three replications and four plants per replica was used to arrange the pots. Analysis of variance (ANOVA) and the Tukey Post Hoc test was used by the SPSS program (Version 26, IBM, USA) to indicate the significantly different means of each treatment.

Results*Effect of different light qualities on the morpho-physiology of spinach*Growth parameters

The growth parameters of spinach (root and shoot length, root volume, fresh and dry mass of root and shoot) were observed to decrease significantly when exposed to Light 3 compared to both Light 2 and Light 1 at both harvest times, as presented in Table 2.

At the first harvest, compared to the control group (Light 2), significant reductions were recorded under Light 3: 48.8% decrease in root length, 73.6% decrease in shoot length, 93.8% decrease in root volume, 95.7% decrease in fresh root weight, 93.5% decrease in fresh shoot weight, 93.7% decrease in dry root weight, and 91.6% decrease in dry shoot weight.

At the full harvest, under Light 3 compared to Light 2, there was a significant decrease in shoot length by 32.0%, fresh root weight by 84.4%, and fresh shoot weight by 71.2%. However, root length increased by 9.5%, and there were decreases in root volume by 55.2%, dry root weight by 9.2%, and dry shoot weight by 28.7%.

Under Light 1, compared to Light 2 (control), there was a significant decrease in root length by 25.9% at the first harvest. However, at the full harvest, there was a significant increase in root volume by 91.7% and fresh root weight by 37.5%.

Furthermore, under Light 1 compared to Light 2, at the first harvest, there was an increase in shoot length by 4.0%, root volume by 6.6%, fresh root weight by 17.6%, fresh shoot weight by 16.7%, and dry root weight by 130%. At the full harvest, there was a decrease in shoot length by 9.7%, an increase in fresh shoot weight by 5.13%, and an increase in dry root weight by 18.3%.

In summary, Light 3 negatively impacted the growth parameters of spinach the most, while Light 1 showed some positive effects on certain growth aspects at both harvest times when compared to the control group (Light 2).

Table 2. Root length (cm), shoot length (cm), root volume (ml), fresh root and shoot, dry root, and shoot weight (g) under different light qualities

Trait	First Harvest (25 th day after sowing)			Full Harvest (35 th day after sowing)		
	Light 1 (Blue >>Red)	Light 2 (Normal White)	Light 3 (Red > Blue > Flashes of UV-C radiations once a week for 5 minutes)	Light 1 (Blue >>Red)	Light 2 (Normal White)	Light 3 (Red > Blue > Flashes of UV-C radiations once a week for 5 minutes)
Root Length (cm)	33.8 ^B	45.6 ^A	23.3 ^B	30.4 ^A	35.7 ^A	39.1 ^A
Shoot Length (cm)	12.9 ^A	12.4 ^A	3.2 ^B	15.7 ^A	17.4 ^A	11.8 ^B
Root volume (ml)	6.58 ^A	6.1 ^A	0.3 ^B	33.6 ^A	17.5 ^B	7.8 ^A
Fresh root weight (g)	5.26 ^A	4.4 ^A	0.19 ^B	29.4 ^A	21.4 ^B	3.3 ^C
Fresh shoot weight (g)	6.77 ^A	5.8 ^A	0.38 ^B	20.5 ^A	19.5 ^A	5.6 ^B
Dry root weight (g)	0.23 ^A	0.1 ^A	0.01 ^B	0.71 ^A	0.6 ^A	0.5 ^A
Dry shoot weight (g)	0.60 ^A	0.4 ^B	0.04 ^C	1.83 ^A	1.9 ^A	1.36 ^A

*Note: Different letters among light qualities within a specific harvest indicate significant differences ($p \leq 0.05$).

Relative chlorophyll content

Relative chlorophyll content (SPAD values) significantly decreased by 44.5, 38.5, 25.5, 30.6, and 22.0% in Light 3 compared to Light 2 respectively, on each day of measurement. Under Light 1, it first decreased by 0.9 and 7.9% (Figure 1) on the first and second days of measurement, later significantly increased by 12.8 and 25.8% on the fourth and fifth day of measurement compared to Light 2.




Figure 1. Relative chlorophyll content (SPAD) during five days under different light qualities (First harvest)

Note: Different letters in each day correspond to significant differences among treatments ($p \leq 0.05$). $n=3$, \pm S.E.

Contents of photosynthetic pigments

Chlorophyll-a ($Chl-a$) significantly decreased by 49.4% under Light 3 and increased by 12.0% under Light 1 compared to Light 2. Chlorophyll- b, also significantly decreased by 56.8% under Light 3 and increased by 25.7% under Light 1 compared to Light 2. Similarly, total carotenoids significantly decreased by 47.1% under Light 3 and increased by 18.1% under Light 1 compared to Light 2 (Figure 2).




Figure 2. Contents of photosynthetic pigments [chlorophyll- a (A), chlorophyll-b (B) and total carotenoids (C) (mg. g^{-1})] in spinach leaves grown under different light qualities provided by LED Lamps. Note: Different letters correspond to significant differences ($p \leq 0.05$) among lights in a particular trait. $n=3$, \pm S.E.

Contents of different carotenoids (antheraxanthin, lutein, violaxanthin)

Under Light 3, antheraxanthin increased by 24.8%, while in Light 1, it decreased by 24.5% compared to Light 2. Lutein increased by 1.32% under Light 3 and 43.4% under Light 1 compared to Light 2. Similarly, violaxanthin increased by 1.9% under Light 3 and 16.4% under Light 1 compared to Light 2 (Table 3).

Table 3. Contents of different carotenoids [antheraxanthin, lutein, and violaxanthin ($\mu\text{g } \mu\text{L}^{-1}$)] and MDA (nmol g^{-1}) under different light qualities

Trait	Light 1	Light 2	Light 3
Antheraxanthin ($\mu\text{g } \mu\text{L}^{-1}$)	3.19 ^A	4.23 ^A	5.2 ^A
Lutein ($\mu\text{g } \mu\text{L}^{-1}$)	136.2 ^A	94.9 ^A	96.2 ^A
Violaxanthin ($\mu\text{g } \mu\text{L}^{-1}$)	57.0 ^A	49.0 ^A	49.0 ^A
Malondialdehyde (nmol g^{-1})	11.9 ^A	13.2 ^A	13.3 ^A

*Note: Different letters correspond to significant differences ($p \leq 0.05$).

Malondialdehyde content

The MDA levels in nmol g^{-1} increased from 11.9 under Light 1 to 13.3 in Light 3, indicating an increase of 11.2%. Similarly, under Light 2, the MDA levels increased to 13.2 nmol g^{-1} , which is a 10.4% increase compared to Light 1 (Table 3).

*Effect of mild drought stress at different light qualities (Full harvest)*Growth parameters

In the presence of 2.5% PEG under Light 1, the following parameters were significantly decreased compared to the control of Light 1: root length by 36.15%, shoot length by 72.2%, root volume by 53.1%, fresh root weight by 97.5%, fresh shoot weight by 97.7%, dry root weight by 95.6%, and dry shoot weight by 91.6%. Similarly, under Light 2, treatment with 2.5% PEG resulted in a significant decrease in the following parameters compared to the control of Light 2: root length by 53.2%, shoot length by 52.3%, root volume by 94%, fresh root weight by 90%, fresh shoot weight by 89.7%, dry root weight by 81.2%, and dry shoot weight by 68.7% (Table 4).

Under Light 3, when exposed to flashes of UV-C radiation and treated with PEG, the plants could not survive. Compared to the PEG treatment of Light 1, it was recorded that root length decreased by 1.25% and root volume decreased by 87.9% under the PEG treatment of Light 2. However, shoot length increased by 65.6%, fresh root weight increased by 238%, fresh shoot weight increased by 300%, and dry root and shoot weight increased by 200% under Light 2 compared to Light 1 (Table 4).

Table 4. Root length(cm), shoot length(cm), root volume (ml), fresh root and shoot weight, dry root and shoot weight (g), SPAD value per day and MDA (nmol g^{-1}) after mild water deprivation under two different light qualities

Trait	Light 1		Light 2	
	Control	2.5% PEG	Control	2.5% PEG
Root length (cm)	33.83 ^{B,1}	21.60 ^{A,2}	45.67 ^{A,1}	21.33 ^{A,2}
Shoot length (cm)	12.92 ^{A,1}	3.58 ^{B,2}	12.42 ^{A,1}	5.92 ^{B,2}
Root volume (ml)	6.58 ^{A,1}	3.08 ^{A,2}	6.17 ^{A,1}	0.37 ^{A,2}
Fresh root weight (g)	5.26 ^{A,1}	0.13 ^{A,2}	4.47 ^{A,1}	0.44 ^{A,2}
Fresh shoot weight (g)	6.77 ^{A,1}	0.15 ^{A,2}	5.88 ^{A,1}	0.60 ^{A,2}
Dry root weight (g)	0.23 ^{A,1}	0.01 ^{A,2}	0.16 ^{A,1}	0.03 ^{A,2}
Dry shoot weight (g)	0.60 ^{A,1}	0.05 ^{A,2}	0.48 ^{B,1}	0.15 ^{A,2}
SPAD day 1	39.4 ^{A,1}	21.2 ^{A,1}	39.7 ^{A,1}	41.9 ^{A,1}
SPAD day 2	39.8 ^{A,1}	14.6 ^{B,2}	43.2 ^{A,1}	39.3 ^{A,1}
Malondialdehyde (nmol g^{-1})	13.20 ^{A,1}	16.22 ^{A,1}	11.95 ^{A,1}	16.41 ^{A,2}

*Note: Different letters indicate the significant difference at $p \leq 0.05$ within treatments in a different light and the different numbers indicate the significant difference at $p \leq 0.05$ within treatments in a particular light.

Relative chlorophyll content

When treated with 2.5% PEG under Light 1, the SPAD value decreased by 46% and 63.3% (measured at 2-day intervals) compared to control. In contrast, under Light 2, there was an initial increase of 5.5% in SPAD value, followed by a decrease of 8.9%. In PEG treatment under Light 2, an increase of 97.6% compared to PEG treatment of Light 1 on the first day and 69% on day 2 was recorded (Table 4).

Malondialdehyde (MDA) content

In the presence of 2.5% PEG under Light 1, the MDA levels increased from 13.2 to 16.2 nmol/g , indicating an increase of 22.8% compared to control. Similarly, under Light 2, the MDA levels increased from 11.9 to 16.4 nmol/g , indicating an increase of 37.3% when treated with 2.5% PEG. However, on comparing with light + PEG treatment, under light 2, 1.8% increase was recorded compared to Light 1 (Table 4).

Discussion

Light and water are two crucial needs of plants regulating their survival, morpho-physiology, and productivity (McElrone *et al.*, 2013; Bayat *et al.*, 2018; Klem *et al.*, 2019), and abundance or scarcity of any of them imposes deleterious effects on plants and limit crop productivity (Fahad *et al.*, 2017). In nature, plants are exposed to a combination of stresses, and their impact differs from the individual impact of stress (Francini and Sebastiani, 2019). Combinatorial stresses may be antagonistic or synergistic to each other. For instance, heat and drought stress acts synergistically and cause more damage than their individual treatments (Pandey *et al.*, 2017). In contrast, some stresses in combination (drought and ozone, drought, and pathogen) act antagonistically. Some stress combinations have a net neutral or beneficial effect on plants because they cancel out one another (Pandey *et al.*, 2017). In our results the combination of mild water deprivation and flashes of UV-C acted synergistically and proved fatal to spinach. Yield is not always negatively impacted since one stress may give rise to another's resistance.

Individual drought and ozone stresses hinder the growth of *Medicago truncatula* (alfalfa), but the two together boost the plant's ability to withstand the combined stress (Puckette *et al.*, 2007). The present study showed a differential response of spinach exposed to different light qualities and the response of spinach under mild water deprivation in different light qualities. Both light (quality and quantity) and drought induce various physiological, molecular, and biochemical responses in plants, thereby inducing the generation of active oxygen species (AOS), resulting in oxidative damage at the cellular level (Katerova *et al.*, 2017; Mewis *et al.*, 2012; Foyer and Noctor, 2002; Czczuga, 1987; Muraoka *et al.*, 2002).

Plant growth is a complex process that can be significantly influenced by various environmental factors, including the quality and quantity of light. These factors play a crucial role in determining the growth characteristics of plants, such as root and shoot length, root volume, fresh root and shoot weight, as well as dry root and shoot weight. When plants are subjected to unfavorable light conditions, such as insufficient or poor-quality light, their growth is adversely affected. In the present experiment, growth parameters (root and shoot length, root volume, fresh root and shoot weight, dry root and shoot weight) were affected in plants raised under different light qualities, and maximum reduction was observed in Light 3 (Red > Blue > along with flashes of UV-C once a week) compared to Light 2 (Red > Blue > taken as control) and Light 1 (Blue >>Red) (Table 2). The reduction could be due to oxidative stress generated by a higher accumulation of AOS and MDA content, leading to a decline in chlorophyll content (Figure 2) and consequently decreasing the growth in stressed plants. Moreover, UV-C light has also been shown to be harmful to chloroplasts, mitochondria, and, more broadly, membranes in higher plants. Starting with chloroplasts, UV-C light has been shown to directly destroy plastoquinone (Shavit and Avron, 1963). According to Wituszynska *et al.* (2014), UV-C radiation damages the chloroplasts in *Arabidopsis thaliana* and causes morphological alterations in mesophyll cells that mimic apoptosis. UV radiation degrades the endogenous Phytohormones such as auxin levels, thus inferring normal growth and developmental processes in plants (Vandenbussche *et al.*, 2014). Similar results were recorded in *Vigna unguiculata* L. (Lingakumar and Kulandaivelu, 1993) and *Beta vulgaris* L. (Panagopoulos *et al.*, 1990). UV exposure also interferes with various critical physiological processes associated with biomass production, thus reducing the growth and biomass of crop plants, as observed in lettuce (Paul *et al.*, 2012), rice (Hidema *et al.*, 2005), wheat and sorghum (Kataria and Guruprasad, 2012). Plant height reduction in vegetables was also recorded earlier (Ruhland *et al.*, 2007; Singh *et al.*, 2011; Kataria *et al.*, 2013; Baroniya *et al.*, 2014; Zhang *et al.*, 2014). UV-induced plant height reductions are more or less due to the direct damage to macromolecules such as DNA (Britt, 1999) and proteins (Gerhardt *et al.*, 1999). UV exposure consequently leads to changes in the morphology and growth, e.g., smaller but thicker leaves, less plant height and lower biomass (Suchar and Robberecht, 2015). This may be the reason that in present experiment plants could not survive under combinatorial stress of drought and UV-C radiation.

In our experiment, the presence of 2.5% PEG (Polyethylene Glycol) under different light conditions (Light 1, Light 2, and Light 3) has been investigated. The results indicate that under Light 1, the parameters related to plant growth, including root length, shoot length, root volume, fresh root weight, fresh shoot weight, dry root weight, and dry shoot weight, showed a significant decrease compared to the control. Similarly, under Light 2, the same growth characteristics exhibited a significant decrease when plants were treated with 2.5% PEG. These findings suggest that in both light conditions, the presence of PEG negatively impacts plant growth, resulting in reduced root and shoot length, decreased root volume, as well as lower fresh and dry root and shoot weights. Moreover, under Light 3, the combination of flashes of UV-C radiation and PEG treatment proved to be detrimental to plant survival. Under different light qualities (Light 1 and Light 2), and mild drought along the growth traits, relative chlorophyll content decreased compared to the control (Table 4). According to Qaderi *et al.* (2007), drought stress may cause stunted growth as a result of decreased cell wall flexibility and turgor, changes in plant architecture that result in decreased height, smaller leaves, fewer leaves, decreased leaf water potential, turgor pressure, stomatal closure, and reduced cell growth and expansion. Other crucial physiological mechanisms, including the production of chlorophyll, photosynthesis, respiration, nutrition, and carbohydrate metabolism, are all impacted by drought stress (Jaleel *et al.*, 2008; Farooq *et al.*, 2009; Li *et al.*, 2011). Overall, these findings highlight the sensitivity of plant growth to environmental factors, particularly light quality, and quantity, as well as the impact of PEG treatment. Understanding these relationships can provide valuable insights into optimizing plant growth conditions and mitigating the negative effects of environmental stressors on plant development.

Light for photosynthesis is pursued by the protein pigment complexes, chlorophylls and carotenoids (Jahns and Holzwarth, 2012). Thus, it is a crucial factor in assessing the plant under different stresses (Habiba *et al.*, 2014). In the present experiment, the levels of total chlorophyll content, chlorophyll a and chlorophyll b and SPAD significantly decreased in plants grown under different light qualities and highest reduction was observed in Light 3 compared to Light 2 and Light 1. The decrease in chlorophyll levels can be attributed to the suppression of chlorophyll production or the breakdown of the pigment along with its precursors, protochlorophyll and protochlorophyllide (Santos *et al.*, 2004; Teramura, 1983; Teramura, 1994). The outcomes of our study are consistent with previous findings reported in various plants such as grape vines (Martinez *et al.*, 2013), soybean (Ambasht and Agarwal, 2003), okra (Kumari *et al.*, 2009), sweet wormwood (Rai *et al.*, 2011), Swedish ivy (Vidovic *et al.*, 2014), cotton, sorghum, wheat (Kataria and Guruprasad, 2007), pea (Singh *et al.*, 2015), basil (Sakalauskaite *et al.*, 2012), common privet (Guidi *et al.*, 2016), canola (Tohidi *et al.*, 2011), maize (Gao *et al.*, 2004), and barley (Palmer *et al.*, 2002). Nonetheless, the impact was more harmful when plants were subjected to a combination of UV-C light and drought stress. The results align with Nyachiro *et al.* (2001), who reported a decrease in chlorophyll content caused by drought stress in *Triticum aestivum*. Reducing total chlorophyll under drought stress suggests a reduced capacity for light-harvesting complexes (Teramura, 1983; Teramura, 1994; Santos *et al.*, 2004).

Carotenoids and xanthophylls are essential to light-dependent processes, photoprotection, and potent free radical scavengers (Jahns and Holzwarth, 2012). According to North *et al.* (2007), carotenoids play a significant role in responding to stress. They act as excellent scavengers of AOS (active oxygen species) and are crucial for defending plants against light stress. Carotenoids protect chlorophylls from photooxidative damage resulting from UV irradiation by dissipating excess excitation energy. In the current study, the level of Carotenoids content (antheraxanthin, lutein and violaxanthin) increased under stress conditions, and the maximum was observed in plants raised under Light 3 (Table 3). In agreement with our results, Molnár *et al.* (2009) observed an increase in carotenoids, namely violaxanthin, luteoxanthin, neoxanthin, antheraxanthin, and lutein, in plants exposed to stress conditions. An increase in carotenoid content represents a biochemical response to alleviate UV stress (Palmer *et al.*, 2002), as observed in maize (Ambasht and Agrawal, 1998) and pea (Singh *et al.*, 2015). UV radiation in higher plants can trigger the rise in carotenoids. An increase in antheraxanthin and zeaxanthin, along with a decrease in other carotenoids, was recorded in Swedish Ivy on

UV-B treatment (Vidovic *et al.*, 2014). Lutein decreased in red lettuce after UV-B treatment (Caldwell and Britz, 2006).

The primary target of most stresses is the cellular membranes that endure lipid peroxidation induced by excessive accumulation of AOS, resulting in excess production of MDA, which indicates stress injury (Martinez *et al.*, 2013). AOS production, lipid peroxidation, and membrane damage are known effects of UV-C radiation (Hideg and Vass, 1996; Takeuchi *et al.*, 1996). Lipid peroxidation of a membrane can be directly predicted by MDA content (Health and Packer, 1968) and MDA increase under stress conditions (Lei *et al.*, 2007). In the present study, higher MDA content was observed in plants raised under Light 3, and the effect was synergistic in the case of a combination of different lights and water deprivation. Higher accumulation of MDA indicates more lipid peroxidation in the cell membrane and consequently will lead to cell death. Similar results were also observed in rice (Zhou *et al.*, 2007), *Withania somnifera* (Takshak and Agarwal, 2014), and *Ditylum brightwellii* (Rijstenbil, 2001).

Conclusions

This study emphasizes the significance of blue light in plant growth and the influence of varied light qualities on several aspects of plant development. The presence or absence of blue light may impact root and shoot length, fresh and dry weight, root volume, chlorophyll, carotenoid content, and malondialdehyde (MDA) levels. According to the study, when using LED technology, spinach develops better when exposed to more blue light than red light. UV-C rays, on the other hand, may harm plants, especially when combined with water deprivation. It was found that under such circumstances, mild water stress does not have any beneficial impacts on the development and quality of spinach. The correct UV-C treatment may increase the carotenoid content of spinach, but determining the appropriate doses and knowing how it interacts with the environment will be challenging problems to solve in the future. The findings emphasize the need to consider UV-C radiation and the light spectrum when designing lighting systems for indoor plant growth.

Authors' Contributions

Conceptualization, S.V.; methodology, S.V.; software, O.B.; validation, S.V., O.B. and T.B.Z.; formal analysis, O.B.; investigation, T.B.Z...; resources, S.V.; data curation, S.V.; writing—original draft preparation, T.B.Z. and S.V.; writing—review and editing, O.B. and S.V.; visualization, O.B.; supervision, S.V.; project administration, S.V.; funding acquisition, S.V. All authors have read and agreed to the published version of the manuscript.

Ethical approval (for researches involving animals or humans)

Not applicable.

Acknowledgements

This work was supported by the National Research, Development and Innovation Fund of Hungary, grant number TKP2021-NKTA-32, financed under the TKP2021-NKTA funding scheme. This study was also supported by the National Research, Development, and Innovation Office of Hungary under grant: RRF-2.3.1-21-2022-00008.

Conflict of Interests

The authors declare that there are no conflicts of interest related to this article.

References

- Akinci Ş, Lösel DM (2012). Plant water-stress response mechanisms. *Water Stress*. <https://doi.org/10.5772/29578>
- Ambasht NK and Agrawal M (2003). Effects of enhanced UV-B radiation and tropospheric ozone on physiological and biochemical characteristics of field grown wheat. *Biologia Plantarum* 46(4):625-628. <https://doi.org/10.1023/b:biop.0000041076.95209.c3>
- Ambasht, Navin Kumar, Agrawal M (1998). Physiological and biochemical responses of *Sorghum vulgare* plants to supplemental ultraviolet-B radiation. *Canadian Journal of Botany* 76(7):1290-1294. <https://doi.org/10.1139/b98-137>
- Baroniya SS, Kataria S, Pandey GP, Guruprasad KN (2014). Growth, photosynthesis, and nitrogen metabolism in soybean varieties after exclusion of the UV-B and UV-A/B components of solar radiation. *The Crop Journal* 2(6):388-397. <https://doi.org/10.1016/j.cj.2014.08.002>
- Barros L, Morales P, Carvalho AM, Ferreira IC (2016). Antioxidant potential of wild plant foods. *Mediterranean Wild Edible Plants* 209-232. https://doi.org/10.1007/978-1-4939-3329-7_10
- Barylá A, Laborde C, Montillet JL, Triantaphylidès C, Chagvardieff P (2000). Evaluation of lipid peroxidation as a toxicity bioassay for plants exposed to copper. *Environmental Pollution* 109(1):131-135. [https://doi.org/10.1016/s0269-7491\(99\)00232-8](https://doi.org/10.1016/s0269-7491(99)00232-8)
- Bayat L, Arab M, Aliniaiefard S, Seif M, Lastochkina O, Li T (2018). Effects of growth under different light spectra on the subsequent high light tolerance in rose plants. *AoB Plants* 10(5). <https://doi.org/10.1093/aobpla/ply052>
- Bian ZH, Yang QC, Liu WK (2014). Effects of light quality on the accumulation of phytochemicals in vegetables produced in controlled environments: A Review. *Journal of the Science of Food and Agriculture* 95(5):869-877. <https://doi.org/10.1002/jsfa.6789>
- Bravo S, García-Alonso J, Martín-Pozuelo G, Gómez V, García-Valverde V, Navarro-González I, Periago MJ (2013). Effects of postharvest UV-C treatment on carotenoids and phenolic compounds of vine-ripe tomatoes. *International Journal of Food Science & Technology* 48(8):1744-1749. <https://doi.org/10.1111/ijfs.12146>
- Britt AB (1999). Molecular genetics of DNA repair in higher plants. *Trends in Plant Science* 4(1):20-25. [https://doi.org/10.1016/s1360-1385\(98\)01355-7](https://doi.org/10.1016/s1360-1385(98)01355-7)
- Caldwell CR, Britz SJ (2006). Effect of supplemental ultraviolet radiation on the carotenoid and chlorophyll composition of greenhouse-grown leaf lettuce (*Lactuca sativa* L.) cultivars. *Journal of Food Composition and Analysis* 19(6-7):637-644. <https://doi.org/10.1016/j.jfca.2005.12.016>
- Cogo SLP, Chaves FC, Schirmer MA, Zambiazzi RC, Nora L, Silva JA, Rombaldi CV (2011). Low soil water content during growth contributes to preservation of green colour and bioactive compounds of cold-stored broccoli (*Brassica oleraceae* L.) florets. *Postharvest Biology and Technology* 60(2):158-163. <https://doi.org/10.1016/j.postharvbio.2010.12.008>
- Czczuga B (1987). Carotenoid contents in leaves grown under various light intensities. *Biochemical Systematics and Ecology* 15(5):523-527. [https://doi.org/10.1016/0305-1978\(87\)90098-6](https://doi.org/10.1016/0305-1978(87)90098-6)

- Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, ... Huang J (2017). Crop production under drought and heat stress: Plant Responses and Management Options. *Frontiers in Plant Science* 8. <https://doi.org/10.3389/fpls.2017.01147>
- Farooq M, Wahid A, Kobayashi N, Fujita D, Basra SM (2009). Plant drought stress: Effects, mechanisms and management. *Sustainable Agriculture* 153-188. https://doi.org/10.1007/978-90-481-2666-8_12
- Folta KM (2004). Green light stimulates early stem elongation, antagonizing light-mediated growth inhibition. *Plant Physiology* 135(3):1407-1416. <https://doi.org/10.1104/pp.104.038893>
- Foyer CH, Noctor G (2002). Photosynthetic nitrogen assimilation: Inter-pathway control and signaling. *Advances in Photosynthesis and Respiration* 1-22. https://doi.org/10.1007/0-306-48138-3_1
- Francini A, Sebastiani L (2019). Abiotic stress effects on performance of horticultural crops. *Horticulturae* 5(4):67. <https://doi.org/10.3390/horticulturae5040067>
- Gao W, Zheng Y, Slusser JR, Heisler GM, Grant RH, Xu J, He D (2004). Effects of supplementary ultraviolet-B irradiance on maize yield and qualities: A field experiment. *Photochemistry and Photobiology* 80(1):127. <https://doi.org/10.1562/2004-05-03-ra-156.1>
- Gerhardt KE, Wilson MI, Greenberg BM (1999). Tryptophan photolysis leads to a UVB-induced 66 KDA photoproduct of ribulose-1,5 bisphosphate carboxylase/oxygenase (rubisco) *in vitro* and *in vivo*. *Photochemistry and Photobiology* 70(1):49-56. <https://doi.org/10.1111/j.1751-1097.1999.tb01948.x>
- Gilani M, Danish S, Ahmed N, Rahi AA, Akrem A, Younis U, ... Iqbal RK (2020). Mitigation of drought stress in spinach using individual and combined applications of salicylic acid and potassium. *Pakistan Journal of Botany* 52(5). [https://doi.org/10.30848/pjb2020-5\(18\)](https://doi.org/10.30848/pjb2020-5(18))
- Goodwin TW, Britton GEORGE (1988). Distribution and analysis of carotenoids. *Plant Pigments* 61-132.
- Guidi L, Brunetti C, Fini A, Agati G, Ferrini F, Gori A, Tattini M (2016). UV radiation promotes flavonoid biosynthesis, while negatively affecting the biosynthesis and the de-epoxidation of xanthophylls: Consequence for photoprotection? *Environmental and Experimental Botany* 127:14-25. <https://doi.org/10.1016/j.envexpbot.2016.03.002>
- Habiba U, Ali S, Farid M, Shakoor MB, Rizwan M, Ibrahim M, ... Ali B (2014). EDTA enhanced plant growth, antioxidant defense system, and phytoextraction of copper by *Brassica napus* L. *Environmental Science and Pollution Research* 22(2):1534-1544. <https://doi.org/10.1007/s11356-014-3431-5>
- Harborne JB, Williams CA (2000). Advances in flavonoid research since 1992. *Phytochemistry* 55(6):481-504. [https://doi.org/10.1016/s0031-9422\(00\)00235-1](https://doi.org/10.1016/s0031-9422(00)00235-1)
- Hasan M, Bae H (2017). An overview of stress-induced resveratrol synthesis in grapes: Perspectives for resveratrol-enriched grape products. *Molecules* 22(2):294. <https://doi.org/10.3390/molecules22020294>
- Hassan EA, Al-Zuhairi WSh, Ibrahim WA (2022). Antioxidants and their role in preventing diseases: A Review. *Earthline Journal of Mathematical Sciences* 165-182. <https://doi.org/10.34198/ejcs.7222.165182>
- Heath RL, Packer L (1968). Photoperoxidation in isolated chloroplasts. *Archives of Biochemistry and Biophysics* 125(3):850-857. [https://doi.org/10.1016/0003-9861\(68\)90523-7](https://doi.org/10.1016/0003-9861(68)90523-7)
- Heinze M, Hanschen FS, Wiesner-Reinhold M, Baldermann S, Gräfe J, Schreiner M, Neugart S (2018). Effects of developmental stages and reduced UVB and low UV conditions on plant secondary metabolite profiles in pak choi (*Brassica rapa* subsp. *chinensis*). *Journal of Agricultural and Food Chemistry* 66(7):1678-1692. <https://doi.org/10.1021/acs.jafc.7b03996>
- Hideg É, Vass I (1996). UV-B induced free radical production in plant leaves and isolated thylakoid membranes. *Plant Science* 115(2):251-260. [https://doi.org/10.1016/0168-9452\(96\)04364-6](https://doi.org/10.1016/0168-9452(96)04364-6)
- Hidema J, Teranishi M, Iwamatsu Y, Hirouchi T, Ueda T, Sato T, ... Kumagai T (2005). Spontaneously occurring mutations in the cyclobutane pyrimidine dimer photolyase gene cause different sensitivities to ultraviolet-B in Rice. *The Plant Journal* 43(1):57-67. <https://doi.org/10.1111/j.1365-313x.2005.02428.x>
- Jahns P, Holzwarth AR (2012). The role of the xanthophyll cycle and of Lutein in photoprotection of Photosystem II. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* 1817(1):182-193. <https://doi.org/10.1016/j.bbabi.2011.04.012>
- Jaleel CA, Gopi R, Sankar B, Gomathinayagam M, Panneerselvam R (2008). Differential responses in water use efficiency in two varieties of *Catharanthus roseus* under drought stress. *Comptes Rendus Biologies* 331(1):42-47. <https://doi.org/10.1016/j.crv.2007.11.003>

- Kaiser E, Weerheim K, Schipper R, Dieleman JA (2019). Partial replacement of red and blue by green light increases biomass and yield in tomato. *Scientia Horticulturae* 249:271-279. <https://doi.org/10.1016/j.scienta.2019.02.005>
- Kataria S, Guruprasad KN (2012). Intraspecific variations in growth, yield, and photosynthesis of sorghum varieties to ambient UV (280–400 nm) radiation. *Plant Science* 196:85-92. <https://doi.org/10.1016/j.plantsci.2012.07.011>
- Kataria S, Guruprasad KN, Ahuja S, Singh B (2013). Enhancement of growth, photosynthetic performance, and yield by exclusion of ambient UV components in C3 and C4 plants. *Journal of Photochemistry and Photobiology B: Biology* 127:140-152. <https://doi.org/10.1016/j.jphotobiol.2013.08.013>
- Katerova Z, Todorova D, Sergiev I (2017). Plant secondary metabolites and some plant growth regulators elicited by UV irradiation, light and/or shade. *Medicinal Plants and Environmental Challenges* 97-121. https://doi.org/10.1007/978-3-319-68717-9_6
- Klem K, Ač A, Holub P, Kováč D, Špunda V, Robson TM, Urban O (2012). Interactive effects of Par and UV radiation on the physiology, morphology and leaf optical properties of two barley varieties. *Environmental and Experimental Botany* 75:52-64. <https://doi.org/10.1016/j.envexpbot.2011.08.008>
- Klem K, Gargallo-Garriga A, Rattanapichai W, Oravec M, Holub P, Veselá B, ... Urban O (2019). Distinct morphological, physiological, and biochemical responses to light quality in barley leaves and roots. *Frontiers in Plant Science* 10. <https://doi.org/10.3389/fpls.2019.01026>
- Kumari R, Singh S, Agrawal SB (2009). Combined effects of psoralens and ultraviolet-B on growth, pigmentation, and biochemical parameters of *Abelmoschus esculentus* L. *Ecotoxicology and Environmental Safety* 72(4):1129-1136. <https://doi.org/10.1016/j.ecoenv.2008.12.009>
- Lei Z, Mingyu S, Xiao W, Chao L, Chunxiang Q, Liang C, Fashui H (2007). Antioxidant stress is promoted by nano-anatase in spinach chloroplasts under UV-B radiation. *Biological Trace Element Research* 121(1):69-79. <https://doi.org/10.1007/s12011-007-8028-0>
- Li C, Jiang D, Wollenweber B, Li Y, Dai T, Cao W (2011). Waterlogging pretreatment during vegetative growth improves tolerance to waterlogging after anthesis in wheat. *Plant Science* 180(5):672-678. <https://doi.org/10.1016/j.plantsci.2011.01.009>
- Lingakumar K, Kulandaivelu G (1993). Changes induced by ultraviolet-B radiation in vegetative growth, foliar characteristics, and photosynthetic activities in *Vigna unguiculata*. *Functional Plant Biology* 20(3):299. <https://doi.org/10.1071/pp9930299>
- Marschner H, Oberle H, Cakmak I, Römheld V (1990). Growth enhancement by silicon in cucumber (*Cucumis sativus*) plants depends on imbalance in phosphorus and zinc supply. *Plant and Soil* 124(2):211-219. <https://doi.org/10.1007/bf00009262>
- Martínez-Lüscher J, Morales F, Delrot S, Sánchez-Díaz M, Gomés E, Aguirreola J, Pascual I (2013). Short- and long-term physiological responses of grapevine leaves to UV-B radiation. *Plant Science* 213:114-122. <https://doi.org/10.1016/j.plantsci.2013.08.010>
- McElrone AJ, Choat B, Gambetta GA, Brodersen CR (2013). Water uptake and transport in vascular plants. *Nature Education Knowledge* 4(5):6. <https://doi.org/10.3389/fpls.2013.00108>
- Mewis I, Schreiner M, Nguyen CN, Krumbein A, Ulrichs C, Lohse M, Zrenner R (2012). UV-B irradiation changes specifically the secondary metabolite profile in broccoli sprouts: Induced signalling overlaps with defence response to biotic stressors. *Plant and Cell Physiology* 53(9):1546-1560. <https://doi.org/10.1093/pcp/pcs096>
- Molnár P, Deli J, Tanaka T, Kann Y, Tani S, Gyémánt N, Molnár J, Kawase M (2009). Carotenoids with anti-*Helicobacter pylori* activity from golden delicious apple. *Phytotherapy Research* 24(5):644-648. <https://doi.org/10.1002/ptr.2912>
- Moran R, Porath D (1980). Chlorophyll determination in intact tissues using *N,n*-dimethylformamide. *Plant Physiology* 65(3):478-479. <https://doi.org/10.1104/pp.65.3.478>
- Muraoka H, Tang Y, Koizumi H, Washitani I (2002). Effects of light and soil water availability on leaf photosynthesis and growth of *Arisaema heterophyllum*, a riparian forest understorey plant. *Journal of Plant Research* 115(6):419-427. <https://doi.org/10.1007/s10265-002-0052-8>
- Nguyen TPD, Vu NT, Nguyen QT, Tran TTH, Cao PB, Kim IS, Jang, DC (2022). Growth and quality of hydroponic cultivated spinach (*Spinacia oleracea* L.) affected by the light intensity of red and blue LEDs. *Sains Malaysiana* 51(2):473-483. <https://doi.org/10.17576/jsm-2022-5102-12>

- North HM, Almeida AD, Boutin JP, Frey A, To A, Botran L, Sotta B, Marion-Poll A (2007). The *Arabidopsis* ABA-deficient mutant ABA4 demonstrates that the major route for stress-induced ABA accumulation is via neoxanthin isomers. *The Plant Journal* 50(5):810-824. <https://doi.org/10.1111/j.1365-313x.2007.03094.x>
- Nyachiro JM, Briggs KG, Hoddinott J, Johnson-Flanagan AM (2001). Chlorophyll content, chlorophyll fluorescence and water deficit in spring wheat. *Cereal Research Communications* 29(1-2):135-142. <https://doi.org/10.1007/bf03543653>
- Ouzounis T, Rosenqvist E, Ottosen CO (2015). Spectral effects of artificial light on plant physiology and secondary metabolism: A Review. *HortScience* 50(8):1128-1135. <https://doi.org/10.21273/hortsci.50.8.1128>
- Paim BT, Crizel RL, Tatiane SJ, Rodrigues VR, Rombaldi CV, Galli V (2020). Mild drought stress has potential to improve lettuce yield and quality. *Scientia Horticulturae* 272:109578. <https://doi.org/10.1016/j.scienta.2020.109578>
- Palmer H, Ohta M, Watanabe M, Suzuki T (2002). Oxidative stress-induced cellular damage caused by UV and methyl viologen in *Euglena gracilis* and its suppression with rutin. *Journal of Photochemistry and Photobiology B: Biology* 67(2):116-129. [https://doi.org/10.1016/s1011-1344\(02\)00271-3](https://doi.org/10.1016/s1011-1344(02)00271-3)
- Panagopoulos I, Bornman JF, Björn LO (1990). Effects of ultraviolet radiation and visible light on growth, fluorescence induction, ultraweak luminescence and peroxidase activity in sugar beet plants. *Journal of Photochemistry and Photobiology B: Biology* 8(1):73-87. [https://doi.org/10.1016/1011-1344\(90\)85189-4](https://doi.org/10.1016/1011-1344(90)85189-4)
- Pandey P, Irulappan V, Bagavathiannan MV, Senthil-Kumar M (2017). Impact of combined abiotic and biotic stresses on plant growth and avenues for crop improvement by exploiting physio-morphological traits. *Frontiers in Plant Science* 8. <https://doi.org/10.3389/fpls.2017.00537>
- Paradiso R, Proietti S (2022). Light-quality manipulation to control plant growth and photomorphogenesis in greenhouse horticulture: The state of the art and the opportunities of modern LED systems. *Journal of Plant Growth Regulation* 41(2):742-780. <https://doi.org/10.1007/s00344-021-10337-y>
- Paradiso R, Meinen E, Snel JFH, De Visser P, Van Ieperen W, Hogewoning SW, Marcelis LFM (2011). Spectral dependence of photosynthesis and light absorbance in single leaves and canopy in rose. *Scientia Horticulturae* 127(4):548-554. <https://doi.org/10.1016/j.scienta.2010.11.017>
- Paul ND, Moore JP, McPherson M, Lambourne C, Croft P, Heaton JC, Wargent JJ (2012). Ecological responses to UV radiation: Interactions between the biological effects of UV on plants and on associated organisms. *Physiologia Plantarum* 145(4):565-581. <https://doi.org/10.1111/j.1399-3054.2011.01553.x>
- Poiroux-Gonord F, Bidet LP, Fanciullino AL, Gautier H, Lauri-Lopez F, Urban L (2010). Health benefits of vitamins and secondary metabolites of fruits and vegetables and prospects to increase their concentrations by agronomic approaches. *Journal of Agricultural and Food Chemistry* 58(23):12065-12082. <https://doi.org/10.1021/jf1037745>
- Puckette MC, Weng H, Mahalingam R (2007). Physiological and biochemical responses to acute ozone-induced oxidative stress in *Medicago truncatula*. *Plant Physiology and Biochemistry* 45(1):70-79. <https://doi.org/10.1016/j.plaphy.2006.12.004>
- Qaderi MM, Reid DM, Yeung EC (2007). Morphological and physiological responses of canola (*Brassica napus*) siliques and seeds to UVB and CO₂ under controlled environment conditions. *Environmental and Experimental Botany*, 60(3):428-437. <https://doi.org/10.1016/j.envexpbot.2006.12.019>
- Rai R, Meena RP, Smita SS, Shukla A, Rai SK, Pandey-Rai S (2011). UV-B and UV-C pre-treatments induce physiological changes and artemisinin biosynthesis in *Artemisia annua* L. – an antimalarial plant. *Journal of Photochemistry and Photobiology B: Biology* 105(3):216-225. <https://doi.org/10.1016/j.jphotobiol.2011.09.004>
- Ribera A, Bai Y, Wolters, AMA, van Treuren R, Kik C (2020). A review on the Genetic Resources, domestication and breeding history of spinach (*Spinacia oleracea* L.). *Euphytica* 216(3). <https://doi.org/10.1007/s10681-020-02585-y>
- Rijstenbil J (2001). Effects of periodic, low UVA radiation on cell characteristics and oxidative stress in the marine planktonic *Diatomditylum brightwellii*. *European Journal of Phycology* 36(1):1-8. <https://doi.org/10.1080/09670260110001735138>
- Roberts JL, Moreau R (2016). Functional properties of spinach (*Spinacia oleracea* L.) phytochemicals and bioactives. *Food & Function* 7(8):3337-3353. <https://doi.org/10.1039/c6fo00051g>
- Ruhland C, Fogal M, Buyarski C, Krna M (2007). Solar ultraviolet-B radiation increases phenolic content and ferric reducing antioxidant power in *Avena sativa*. *Molecules* 12(6):1220-1232. <https://doi.org/10.3390/12061220>

- Sakalauskaitė J, Viskelis P, Dambrauskienė E, Sakalauskienė S, Samuolienė G, Brazaitytė A, ...Urbanavičienė D (2012). The effects of different UV-B radiation intensities on morphological and biochemical characteristics in *Ocimum basilicum* L. Journal of the Science of Food and Agriculture 93(6):1266-1271. <https://doi.org/10.1002/jsfa.5879>
- Santos I, Fidalgo F, Almeida JM, Salema R (2004). Biochemical and ultrastructural changes in leaves of potato plants grown under supplementary UV-B radiation. Plant Science 167(4):925-935. <https://doi.org/10.1016/j.plantsci.2004.05.035>
- Shavit N, Avron M (1963). The effect of ultraviolet light on photophosphorylation and the Hill reaction. Biochimica et Biophysica Acta 66:187-195. [https://doi.org/10.1016/0006-3002\(63\)91185-5](https://doi.org/10.1016/0006-3002(63)91185-5)
- Singh S, Agrawal SB, Agrawal M (2015). Responses of pea plants to elevated UV-B radiation at varying nutrient levels: N-metabolism, carbohydrate pool, total phenolics and yield. Functional Plant Biology 42(11):1045. <https://doi.org/10.1071/fp15003>
- Singh S, Kumari R, Agrawal M, Agrawal SB (2011). Modification in growth, biomass and yield of radish under supplemental UV-B at different NPK Levels. Ecotoxicology and Environmental Safety 74(4):897-903. <https://doi.org/10.1016/j.ecoenv.2010.12.011>
- Sorrentino M, Colla G, Roupael Y, Panzarová K, Trtílek M (2020). Lettuce reaction to drought stress: Automated high-throughput phenotyping of plant growth and photosynthetic performance. Acta Horticulturae 1268:133-142. <https://doi.org/10.17660/actahortic.2020.1268.17>
- Suchar VA, Robberecht R (2015) Integration and scaling of UV-B radiation effects on plants: from DNA to leaf. Ecology and Evolution 5(13):2544-2555. <https://doi.org/10.1002/ece3.1332>
- Takeuchi Y, Kubo H, Kasahara, H, Sakaki T (1996). Adaptive alterations in the activities of scavengers of active oxygen in cucumber cotyledons irradiated with UV-B. Journal of Plant Physiology 147(5):589-592. [https://doi.org/10.1016/s0176-1617\(96\)80050-2](https://doi.org/10.1016/s0176-1617(96)80050-2)
- Takshak S, Agrawal SB (2014). Effect of ultraviolet-B radiation on biomass production, lipid peroxidation, reactive oxygen species, and antioxidants in *Withania Somnifera*. Biologia Plantarum 58(2):328-334. <https://doi.org/10.1007/s10535-014-0390-0>
- Teramura AH (1983). Effects of ultraviolet-B radiation on the growth and yield of Crop Plants. Physiologia Plantarum, 58(3):415-427. <https://doi.org/10.1111/j.1399-3054.1983.tb04203.x>
- Teramura AH, Sullivan JH (1994). Effects of UV-B radiation on photosynthesis and growth of terrestrial plants. Photosynthesis Research 39(3):463-473. <https://doi.org/10.1007/bf00014599>
- Tohidi Moghadam HR, Ghooshchi F, Zahedi H (2011). Effect of UV radiation and evaluated CO₂ on morphological traits, yield and yield components of canola (*Brassica napus* L.) grown under water deficit stress. Notulae Botanicae Horti Agrobotanici Cluj-Napoca 39(1):213. <https://doi.org/10.15835/nbha3915534>
- Tóth VR, Mészáros I, Veres S, Nagy J (2002). Effects of the available nitrogen on the photosynthetic activity and xanthophyll cycle pool of maize in field. Journal of Plant Physiology 159(6):627-634. <https://doi.org/10.1078/0176-1617-0640>
- Urban L, Charles F, de Miranda MR, Aarouf J (2016). Understanding the physiological effects of UV-C light and exploiting its agronomic potential before and after harvest. Plant Physiology and Biochemistry 105:1-11. <https://doi.org/10.1016/j.plaphy.2016.04.004>
- Vandenbussche F, Tilbrook K, Fierro AC, Marchal K, Poelman D, Van Der Straeten D, Ulm R (2014). Photoreceptor-mediated bending towards UV-B in *Arabidopsis*. Molecular Plant 7(6):1041-1052. <https://doi.org/10.1093/mp/ssu039>
- Vandenbussche F, Yu N, Li W, Vanhaelewyn L, Hamshou M, Van Der Straeten D, Smagghe G (2018). An ultraviolet B condition that affects growth and defense in *Arabidopsis*. Plant Science 268:54-63. <https://doi.org/10.1016/j.plantsci.2017.12.005>
- Vidović M, Morina F, Milić S, Zechmann B, Albert A, Winkler JB, Veljović jovanović S (2014). Ultraviolet-B component of sunlight stimulates photosynthesis and flavonoid accumulation in variegated *Plectranthus coleoides* leaves depending on background light. Plant, Cell & Environment 38(5):968-979. <https://doi.org/10.1111/pce.12471>
- Wellburn AR (1994). The spectral determination of chlorophylls a and b, as well as total carotenoids, using various solvents with spectrophotometers of different resolution. Journal of Plant Physiology 144(3):307-313. [https://doi.org/10.1016/s0176-1617\(11\)81192-2](https://doi.org/10.1016/s0176-1617(11)81192-2)

- Wituszyńska W, Szechyńska-Hebda M, Sobczak M, Rusaczek A, Kozłowska-makulska A, Witoń D, Karpiński S (2014). Lesion simulating disease 1 and enhanced disease susceptibility 1 differentially regulate UV-C-induced photooxidative stress signalling and programmed cell death in *Arabidopsis thaliana*. *Plant, Cell & Environment* 38(2):315-330. <https://doi.org/10.1111/pce.12288>
- Zhang L, Allen LH, Vaughan MM, Hauser BA, Boote KJ (2014). Solar ultraviolet radiation exclusion increases soybean internode lengths and plant height. *Agricultural and Forest Meteorology* 184:170-178. <https://doi.org/10.1016/j.agrformet.2013.09.011>
- Zheng L, He H, Song W (2019). Application of light-emitting diodes and the effect of light quality on horticultural crops: A Review. *HortScience* 54(10):1656-1661. <https://doi.org/10.21273/hortsci14109-19>
- Zhou Y, Lam HM, Zhang J (2007). Inhibition of photosynthesis and energy dissipation induced by water and high light stresses in rice. *Journal of Experimental Botany* 58(5):1207-1217. <https://doi.org/10.1093/jxb/erl291>



The journal offers free, immediate, and unrestricted access to peer-reviewed research and scholarly work. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles, or use them for any other lawful purpose, without asking prior permission from the publisher or the author.



License - Articles published in *Notulae Botanicae Horti Agrobotanici Cluj-Napoca* are Open-Access, distributed under the terms and conditions of the Creative Commons Attribution (CC BY 4.0) License.

© Articles by the authors; Licensee UASVM and SHST, Cluj-Napoca, Romania. The journal allows the author(s) to hold the copyright/to retain publishing rights without restriction.

Notes:

- **Material disclaimer:** The authors are fully responsible for their work and they hold sole responsibility for the articles published in the journal.
- **Maps and affiliations:** The publisher stay neutral with regard to jurisdictional claims in published maps and institutional affiliations.
- **Responsibilities:** The editors, editorial board and publisher do not assume any responsibility for the article's contents and for the authors' views expressed in their contributions. The statements and opinions published represent the views of the authors or persons to whom they are credited. Publication of research information does not constitute a recommendation or endorsement of products involved.