

Prevalence of E-Cigarettes in Children and Adolescents A Failure of Public Health Prevention

Surendra Anthony Anish*

University of Delhi, Benito Juarez Marg, South Campus, South Moti Bagh, New Delhi, Delhi 110021, India

*: All correspondence should be sent to: Dr. Surendra Anthony Anish

Author's Contact: Dr. Surendra Anthony Anish, Ph.D., E-mail: surendra.anish@gmail.com

DOI: <https://doi.org/10.15354/si.24.re1080>

Funding: No funding source declared.

COI: The author declares no competing interest.

AI Declaration: The author affirms that artificial intelligence did not contribute to the process of preparing the work.

The rising prevalence of e-cigarette use among children and adolescents poses a significant challenge to public health prevention efforts. As these electronic nicotine delivery systems gain popularity among youth, concerns about the health impacts and long-term consequences have escalated. In this article, we delve into the concerning trends of e-cigarette use among the younger population, exploring the factors driving this phenomenon and evaluating the failures in current public health prevention strategies. By examining the regulatory gaps in e-cigarette marketing to the youth and proposing interventions to combat this growing issue, we aim to shed light on the urgent need for stronger policies and enforcement mechanisms to protect the health and well-being of our youth.

Keywords: E-cigarettes; Adolescents; Prevention; Health Enforcement; Health Wellbeing

Science Insights, 2024 October 31; Vol. 45, No. 4, pp.1593-1602.

© 2024 Insights Publisher. All rights reserved.



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the [Creative Commons Attribution-NonCommercial 4.0 License](https://creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed by the Insights Publisher.

THE PREVALENCE of e-cigarettes among children and adolescents is a substantial failure of public health prevention efforts. The allure of these products to young people remains robust, despite efforts to regulate their sales and marketing (Singh et al., 2020). This appeal is fueled by aggressive advertising campaigns and appealing flavors. Studies have demonstrated that the use of e-cigarettes by young individuals is linked to adverse health consequences, such as an elevated risk of nicotine addiction and potential long-term effects on brain development (Leslie, 2020; Padon et al., 2016). Furthermore, the normalization of vaping among young people has resulted in a rise in the use of traditional cigarettes in conjunction with vaping (Costantino et al., 2024; Duke et al., 2014; Glantz et al.,

2024; Hammond et al., 2019; Jackson et al., 2024). To effectively address this issue, public health authorities must prioritize the enforcement of existing regulations, implement stricter policies on sales and marketing that are specifically designed for youth, increase education about the dangers of vaping, and provide increased access to cessation programs for young users. The future health outcomes of our younger generations will be significantly impacted if we fail to take decisive action now.

Current Prevention Efforts

Public health officials are primarily addressing the issue of e-cigarette use among children by stringent regulations on the sale and marketing of these devices (E-Cigarette Use Among

Youth, 2024). Multiple nations have enacted legislation that forbids the sale of e-cigarettes to anyone below a specified age, generally 18 or 21 years (Kennedy et al., 2016; Snell et al., 2021). Furthermore, marketing limits have been instituted to curtail the promotion of e-cigarettes in environments where minors are likely to be exposed, including schools and youth-targeted media (Marynak et al., 2017).

School-based prevention programs constitute a crucial element in mitigating e-cigarette usage among youth (Bialous & Eijk, 2020; Sreeramareddy et al., 2022). These programs inform students about the hazards of e-cigarette usage, the accompanying health risks, and the significance of making health-conscious decisions. These initiatives seek to equip children with precise knowledge regarding e-cigarettes, so enabling them to make informed choices about their health and well-being.

Healthcare providers have a vital role in avoiding e-cigarette usage among youngsters. By integrating inquiries regarding e-cigarette usage into standard evaluations and counseling sessions, healthcare professionals can recognize at-risk youth and provide them with essential help and tools to prevent or cease e-cigarette consumption (Harvey & Chadi, 2016). Moreover, healthcare professionals can inform parents about the hazards of e-cigarette usage and offer advice on how to communicate with their children regarding the associated risks (Singh et al., 2020).

Community-based programs represent a crucial element in the prevention of e-cigarette usage among minors (Liu et al., 2022). Partnerships among community organizations, educational institutions, healthcare professionals, and public health agencies can enhance knowledge regarding the hazards of e-cigarettes and furnish children and parents with the necessary support to make informed health decisions (Khambayat et al., 2023). Through collaboration, these stakeholders can establish a cohesive stance against the utilization of e-cigarettes among children and adolescents.

Gaps in Policy Implementation and Enforcement

Initially, there is a dearth of regulations that govern the marketing and advertising of e-cigarettes to children and adolescents. Many e-cigarette companies employ flashy advertisements, fruity flavors, and vibrant colors to entice young users, resulting in an increase in their use among this vulnerable demographic (Struik et al., 2020). Companies target children and adolescents without facing any consequences due to the absence of regulation, which results in a hazardous situation in which young people are exposed to harmful substances (Bozzola et al., 2022).

A dearth of explicit communication exists regarding the potential hazards of e-cigarette use among children and adolescents. Although some studies indicate that e-cigarettes may be less harmful than traditional cigarettes, they are by no means secure, particularly for young, developing bodies (Marqu s et al., 2021; Public Health Consequences of E-Cigarettes, 2018; Singh et al., 2020; The Harmful Consequences of Vaping: A Public Health Threat, 2020). Children and adolescents may be more inclined to experiment with e-cigarettes without a comprehensive comprehension of the potential repercussions if there is a lack of clear and consistent messaging regarding the risks asso-

ciated with their use.

Additionally, lack of enforcement of age restrictions on the purchase of e-cigarette is another reason. The sale of these products to individuals under the age of 18 or 21 is prohibited by laws in numerous states (Marynak et al., 2017; State Laws Prohibiting Sales to Minors and Indoor Use of Electronic Nicotine Delivery Systems, 2014; Wang et al., 2020); however, these laws are frequently not rigorously enforced (Nguyen et al., 2020). Consequently, children and adolescents can readily obtain e-cigarettes, which can result in increased usage and potential health risks.

No evidence exists on the long-term health consequences of e-cigarette use in children and adolescents. The long-term effects of e-cigarettes on young, developing bodies are not yet completely understood, despite the fact that some short-term studies have demonstrated that they may be less harmful than traditional cigarettes (Eltorai et al., 2018; Kaisar et al., 2016; Leslie, 2020). It is challenging to establish comprehensive policies that safeguard the health of children and adolescents in the absence of additional research on the potential health hazards associated with e-cigarette use.

Furthermore, there is a scarcity of resources and assistance for children and adolescents who are grappling with the addiction to e-cigarettes (E-Cigarette Use Among Youth and Young Adults: A Report of the Surgeon General, n.d; Galderisi et al., 2020). Nicotine addiction is a genuine concern for juvenile e-cigarette users, and many may find it challenging to discontinue their use of these products without the appropriate support (Bold et al., 2021; Kong et al., 2020). Children and adolescents may continue to use e-cigarettes and risk developing long-term health issues if they lack access to resources such as nicotine replacement therapy and counseling.

There is a lack of education about the risks associated with e-cigarettes in educational institutions and other adolescent settings. The risks associated with e-cigarette use may not be completely comprehended by a significant number of children and adolescents, and they may be more inclined to experiment with these products in the absence of adequate education (Collaco et al., 2014; Drummond & Upson, 2014). Schools and other youth settings can assist in the prevention of the use of e-cigarettes among young people by offering comprehensive education on the hazards of these products.

A lack of collaboration between public health organizations and government bodies about the issue of e-cigarette usage among children and adolescents is a serious issue. It is challenging to establish effective policies that safeguard the health of young people in the absence of collaboration and communication among these entities (Galderisi et al., 2020; Murthy, 2016). By collaborating, government agencies and public health organizations can develop comprehensive strategies to resolve the policy gaps associated with the use of e-cigarettes by children and adolescents (Chu et al., 2021).

Therefore, the public health must be safeguarded by addressing the substantial voids in policy regarding the use of e-cigarettes by children and adolescents. There are numerous areas in which adjustments can be made to better safeguard the health of young people, including the absence of regulations on marketing and advertising and the insufficient resources availa-

ble to those who are struggling with addiction (Ferkol et al., 2018; Galderisi et al., 2020; Papaleontiou et al., 2019). We can strive to reduce the use of e-cigarettes among children and adolescents and protect their health for the future by implementing comprehensive policies that address these gaps.

Regulatory Deficiency in E-Cigarette Marketing for Youth

The primary regulatory deficiency in e-cigarette marketing to adolescents is the absence of prohibitions on advertising methods for e-cigarettes (Collins et al., 2017; Snell et al., 2021). Traditional cigarettes are subject to stringent regulations prohibiting marketing strategies that attract youngsters, such as vibrant colors and enticing flavors (Nguyen et al., 2020; Padon et al., 2016); conversely, e-cigarettes are frequently promoted in manners that specifically appeal to adolescents (Dai, 2018; Papaleontiou et al., 2019). Numerous e-cigarette firms have tasted such as cotton candy and gummy bear, which are evidently aimed toward a youthful demographic (Barrington - Trimis et al., 2014; Harrell et al., 2016; Smith et al., 2023). These strategies can render e-cigarettes appear innocuous and enjoyable, despite their retention of addictive nicotine.

A further regulatory deficiency in e-cigarette marketing is the absence of regulations governing the locations where e-cigarettes may be advertised. Although conventional cigarette advertising is predominantly prohibited on television and radio, e-cigarette businesses are permitted to promote their goods on social media, in magazines, and at events frequented by youth (Collins et al., 2017; Paradise, 2014). This indicates that adolescents are perpetually subjected to e-cigarette marketing, increasing the likelihood of their experimentation with these items.

There is an absence of laws concerning the age employed in e-cigarette advertising. Numerous e-cigarette businesses employ youthful, attractive models in their promotional materials, rendering e-cigarettes appear glamorous and enticing to adolescents (Kong et al., 2014; Krugman, 2016; Le et al., 2023; Marynak et al., 2018; Padon et al., 2016). This form of marketing conveys a perilous message that e-cigarette use is fashionable and appealing, hence heightening the probability that adolescents may experiment with these products.

An absence of regulation regarding the health assertions associated with these items is another issue. Although several e-cigarette manufacturers assert that their products represent a safer substitute for conventional cigarettes, the long-term health implications of e-cigarettes remain predominantly uncertain (Carr, 2014; Glantz & Bareham, 2018; Grana et al., 2014; Kaiser et al., 2016). The absence of regulation may lead young individuals to erroneously see e-cigarettes as innocuous, despite their significant health dangers.

A further regulatory deficiency in e-cigarette marketing to adolescents is the absence of supervision over online advertising. E-cigarette businesses frequently promote their products on social media platforms such as Instagram and Facebook, where a significant number of young individuals engage (Alpert et al., 2021; Czaplicki et al., 2019; Massey et al., 2020; Silver et al., 2023). This online advertising is especially effective in targeting youth, as it can be tailored according to their interests and de-

mographics (Hartoyo & Sitorus, 2021). This indicates that adolescents are incessantly inundated with communications regarding e-cigarettes, hence increasing the likelihood of their experimentation with these items.

There is an absence of restrictions concerning marketing strategies that encourage addiction to e-cigarettes. Numerous e-cigarette companies employ strategies such as providing complimentary samples and discounts to get adolescents to sample their products (Ferkol et al., 2018; Marynak et al., 2018; Vallone et al., 2020). This kind of marketing can generate urgency and enthusiasm over e-cigarettes, increasing the likelihood of nicotine addiction among youth. In the absence of adequate laws, e-cigarette companies might employ these strategies to ensnare youth with their goods.

Moreover, there is little enforcement of current restrictions regarding e-cigarette marketing to minors (Galderisi et al., 2020). Although several restrictions exist that limit the marketing of e-cigarettes, their enforcement is frequently inadequate (Bialous & Eijk, 2020; Bonnie et al., 2015). This indicates that e-cigarette producers can persist in directing their advertising towards young individuals, notwithstanding the possible dangers associated with these products.

Proposed Strategies for Tightening Marketing Regulations

One of the primary strategies for enforcing marketing regulations regarding e-cigarettes is to limit the use of specific advertising techniques that may be appealing to young individuals (Bialous & Eijk, 2020). This encompasses the utilization of cartoon characters, celebrity endorsements, and varieties that are likely to appeal to a younger demographic (Padon et al., 2016). Regulators can prevent young individuals from beginning to use e-cigarettes and potentially developing nicotine addiction by disallowing these marketing practices.

Another critical approach to enforcing marketing regulations regarding e-cigarettes is to mandate that companies furnish more comprehensive information regarding the ingredients and potential health hazards of their products (Grana et al., 2014; Krugman, 2016). Ensure that consumers are thoroughly informed about the risks associated with e-cigarette use in order to enable them to make more informed decisions regarding their use (Paradise, 2014; Zhu et al., 2014). Regulators can assist in safeguarding consumers from misleading or false advertising claims by mandating that companies disclose this information in a manner that is both transparent and readily accessible.

To try to prevent young individuals from obtaining these products, regulators may also enforce more stringent age restrictions on the sale of e-cigarettes. Regulators can mitigate the risk of nicotine addiction among young individuals by mandating that retailers verify the age of customers prior to selling e-cigarettes (Murthy, 2016; Winickoff et al., 2014). Furthermore, the implementation of these age restrictions can serve to mitigate the potential health hazards associated with the use of e-cigarettes at a relatively young age (Abouk & Adams, 2017; Morain & Malek, 2017).

Additionally, regulators may establish restrictions on the marketing of these products in specific regions. For instance, by prohibiting the sale of e-cigarettes in schools, playgrounds, and

other locations where children are present, it is possible to mitigate the exposure of young individuals to these products and prevent them from becoming regular users (Astor et al., 2019; Coxe et al., 2014; McCarthy et al., 2009; Miller et al., 2016). Regulators can contribute to the establishment of a safer and healthier environment for children and adolescents by enacting these types of restrictions.

To guarantee that e-cigarette companies adhere to regulations, regulators may also mandate that they submit their marketing materials for approval prior to dissemination to the public. Regulators can safeguard consumers from potential harm and prevent the dissemination of misleading or false advertising claims by evaluating and approving these materials in advance (Hildebr, 1994). This can also assist in the establishment of a level playing field for companies in the e-cigarette industry and the guarantee that all marketing materials comply with the same standards (Chacón et al., 2024; Collins et al., 2017; Klein et al., 2016).

To discourage companies from engaging in deceptive or harmful advertising practices, regulators may also implement fines or other penalties on them that violate marketing regulations (Nguyen et al., 2020; Wagoner et al., 2019). Regulators can assist in the accountability of companies for their actions and the prevention of them exploiting consumers by enforcing these penalties (Brubaker, 2007; Cohen, 1974). This can also act as a potent deterrent for other companies in the industry and ensure that marketing regulations are taken seriously.

Public health organizations and advocacy groups can collaborate with regulators to increase awareness of the potential health risks associated with e-cigarette use and to enlighten consumers about the significance of making informed decisions regarding their use (Chen & Husten, 2014; Paradise, 2014; Sindelar, 2020; Snell et al., 2021; Tremblay et al., 2015). Regulators can assist in the empowerment of consumers to make healthier decisions and the reduction of the prevalence of e-cigarette use in the general population by disseminating accurate and current information about the dangers of e-cigarette use. This can also assist in alleviating the burden on the healthcare system and preventing the long-term health consequences that are associated with the use of e-cigarettes.

In addition, regulators may establish evidence-based regulations that are substantiated by scientific research and data through collaboration with industry stakeholders and researchers (Chacón et al., 2024; Saitta et al., 2014; Singh et al., 2020). Regulators can guarantee that their marketing regulations are effective in safeguarding consumers and advancing public health by collaborating with expert professionals in the field (Burkitt, 2007; Monteiro et al., 2017). This can assist in the establishment of a more robust regulatory framework for e-cigarettes and guarantee that regulations are founded on the most robust available evidence.

Public Health Interventions to Combat E-Cigarette Use among Youth

It is essential to implement public health interventions to address the emerging public health crisis of e-cigarette use among adolescents (Jenssen et al., 2019). The objective of these interventions is to prevent young individuals from initiating the use of

e-cigarettes and to assist those who are currently using them in quitting. Regulatory measures, community-based initiatives, and education and awareness campaigns are among the numerous strategies that can be implemented to accomplish these objectives (Galderisi et al., 2020; Maddu, 2019).

Education and awareness campaigns are essential components of public health interventions that aim to reduce the use of e-cigarettes among young people. These campaigns have the potential to assist young individuals in comprehending the hazards associated with e-cigarette use and to furnish them with precise information regarding the health consequences of vaping (Hunter & Martin, 2019; Vogel et al., 2020). These campaigns have the potential to deter young people from experimenting with e-cigarettes by raising awareness of their potential hazards.

Another critical approach to addressing the prevalence of e-cigarette use among adolescents is the implementation of regulatory measures (Vogel et al., 2020). These measures may encompass policies that restrict the availability of flavored e-cigarette products that are particularly alluring to young people, as well as restrictions on the sale and marketing of e-cigarettes to young people (Cullen et al., 2019; Czaplicki et al., 2019; Park - Lee et al., 2022). Public health officials can contribute to the reduction of vaping among adolescents by enacting regulations that impede their ability to obtain e-cigarettes.

Additionally, community-based initiatives are crucial for the prevention of e-cigarette usage among adolescents. These programs can offer young individuals the support and resources necessary to cease vaping, as well as the opportunity to engage in healthier behaviors. Community-based programs can assist in the reduction of the appeal of e-cigarettes and the promotion of positive adolescent development by providing activities and services that encourage young people to make healthier choices (Drummond & Upson, 2014).

The marketing strategies employed by e-cigarette companies to target young people are one of the most significant obstacles to combating e-cigarette use in youth (Vogel et al., 2020). These companies frequently employ social media and other platforms to advertise their products to youthful consumers, and they employ flavors and packaging that are appealing to this group of population. Public health interventions must combat these marketing strategies by increasing awareness of the hazards of e-cigarettes and advocating for policies that restrict the marketing and sale of these products to young individuals (Marynak et al., 2018; Padon et al., 2016).

Another obstacle to reducing the use of e-cigarettes among young people is the prevalence of dual use, which involves the use of both e-cigarettes and traditional cigarettes. Many young individuals who use e-cigarettes also smoke traditional cigarettes (Abouk & Adams, 2017; Duke et al., 2014), which can elevate their susceptibility to addiction and other health complications. This issue necessitates public health interventions that offer assistance and resources to young individuals who are having trouble quitting both traditional cigarettes and e-cigarettes (Ferkol et al., 2018; United States Public Health Service. Office of the Surgeon General & National Center for Chronic Disease Prevention and Health Promotion (U.S.), Office on Smoking and Health, 2016; Xiang et al., 2023).

Public health interventions to combat e-cigarette use in

adolescents must also consider the influence of peers and social norms, in addition to education, regulation, and community-based programs (Cheng et al., 2023; Tworek et al., 2014; Vogel et al., 2020). Young individuals are frequently influenced by their peers and social norms that endorse the use of e-cigarettes as a normal and acceptable behavior. The prevalence of e-cigarette use among adolescents can be reduced through interventions that promote healthy social norms around vaping and target peer networks.

It is also crucial for public health interventions to adopt a comprehensive approach to the prevention of e-cigarette use among young people, which encompasses not only their individual behaviors but also the broader social, economic, and environmental factors that influence their decisions. Public health interventions can contribute to the establishment of environments that promote healthy lifestyles and behaviors by addressing the underlying causes of e-cigarette use among youth, including stress, peer pressure, and a lack of access to healthy alternatives (Chacan et al., 2024; Cheng et al., 2023; Galderisi et al., 2020).

A Call to Action for Public Health Initiatives

Public health initiatives that are designed to inform young individuals about the risks and hazards of e-cigarettes are one of the most effective methods of combating their use (Galderisi et al., 2020; Murthy, 2016; *The Harmful Consequences of Vaping: A Public Health Threat*, 2020). Empowering young people to make informed decisions about their health and well-being is possible by increasing awareness of the potential harm that e-cigarettes can cause (Wang et al., 2019). This can assist in preventing them from succumbing to nicotine addiction and potentially developing severe health problems in the future.

Additionally, the marketing and availability of e-cigarettes to young individuals can be restricted through public health initiatives (Gaiha & Halpern - Felsher, 2020; Janssen & Boykan, 2019; Singh et al., 2020). At present, e-cigarette manufacturers employ flashy advertisements and flavors to attract young consumers, presenting these devices as innocuous and enjoyable. We can reduce the likelihood of young people becoming regular users of e-cigarettes and restrict their access to these hazardous products by enacting more stringent regulations on the marketing and sale of e-cigarettes.

The implementation of prevention programs in schools and communities is another critical component. These programs have the potential to establish a culture of awareness and prevention regarding e-cigarettes, provide resources and support for individuals who are interested in quitting, and educate young people about the dangers associated with e-cigarette use (Anic et al., 2018; Centers for Disease Control and Prevention (US), 2012). We can prevent the next generation from succumbing to the perils of e-cigarette addiction by educating young people

about the significance of making healthy choices at an early age.

Partnerships with healthcare providers, community organizations, and government agencies can also be implemented as part of public health initiatives to develop a comprehensive strategy for addressing the use of e-cigarettes by young people (Perelman et al., 2019). We can establish a unified front against the e-cigarette epidemic among young people by collaborating across sectors to combine our resources, expertise, and influence (*E-Cigarette Use Among Youth*, 2024; *Stopping the start: our new plan to create a smokefree generation*, 2023). This multi-sectoral approach has the potential to expand our audience, develop interventions that are more effective, and have a lasting impact on the health and well-being of our adolescents.

Advocacy efforts may also be implemented as part of public health initiatives to advocate more stringent policies and regulations regarding the use of e-cigarettes (*E-Cigarette Use Among Youth and Young Adults: A Report of the Surgeon General*, n.d). This may involve advocating for the imposition of higher taxation on e-cigarettes, the implementation of more stringent age restrictions on their sale, and the prohibition of specific flavors and marketing strategies that are directed at young individuals. By striving to alter the social and political environment surrounding e-cigarettes, we can establish a more supportive and protective environment for our youth, thereby reducing their risk of e-cigarette addiction and its associated health risks (Singh et al., 2020).

Research and data collection can also be incorporated into public health initiatives to gain a more comprehensive understanding of the extent and consequences of adolescent e-cigarette use (Mattingly & Hart, 2024; Shamout et al., 2020; Vogel et al., 2020). By collecting and analyzing data on e-cigarette trends, patterns of use, and associated health outcomes, we can develop more effective prevention strategies and obtain a better understanding of the problem overall. In addition, this research can assist us in monitoring our progress over time, identifying areas of concern, and making data-driven decisions regarding the most effective areas for our efforts.

Conclusion

Addressing the prevalence of e-cigarettes in children and adolescents is critical for safeguarding their health and well-being. By identifying the failures in public health prevention strategies and advocating for stronger policies and enforcement measures, we can work towards mitigating the impact of e-cigarette use on our youth. It is crucial for stakeholders at all levels to collaborate and take decisive action to protect the future generation from the harm associated with e-cigarettes. Together, through concerted efforts and a commitment to public health initiatives, we can strive to create a safer and healthier environment for our children and adolescents. ■

References

- Abouk, R., & Adams, S. (2017). Bans on electronic cigarette sales to minors and smoking among high school students. *Journal of Health Economics*, 54, 17-24. DOI: <https://doi.org/10.1016/j.jhealeco.2017.03.003>
- Alpert, J. M., Chen, H., Riddell, H., Chung, Y. J., & Mu, Y. A. (2021). Vaping and Instagram: A Content Analysis of e-Cigarette Posts Using the Content Appealing to Youth (CAY) Index. *Substance Use & Misuse*, 56(6), 879-887. DOI: <https://doi.org/10.1080/10826084.2021.1899233>
- Anic, G. M., Sawdey, M. D., Jamal, A., & Trivers, K. F. (2018). Frequency of Use Among Middle and High School Student Tobacco Product Users — United States, 2015-2017. *MMWR Morbidity and Mortality Weekly Report*, 67(49), 1353-1357. DOI: <https://doi.org/10.15585/mmwr.mm6749a1>
- Astor, R. L., Urman, R., Barrington-Trimis, J. L., Berhane, K., Steinberg, J., Cousineau, M., Leventhal, A. M., Unger, J. B., Cruz, T., Pentz, M. A., Samet, J. M., & McConnell, R. (2019). Tobacco Retail Licensing and Youth Product Use. *Pediatrics*, 143(2), e20173536. DOI: <https://doi.org/10.1542/peds.2017-3536>
- Barrington-Trimis, J. L., Samet, J. M., & McConnell, R. (2014). Flavorings in Electronic Cigarettes. *JAMA*, 312(23), 2493. DOI: <https://doi.org/10.1001/jama.2014.14830>
- Bialous, S. A., & Van Der Eijk, Y. (2020). How Should Global Tobacco Control Efforts Be Prioritized to Protect Children in Resource-Poor Regions? *The AMA Journal of Ethics*, 22(2), E135-146. DOI: <https://doi.org/10.1001/amaethics.2020.135>
- Bold, K., Kong, G., Cavallo, D., Davis, D., Jackson, A., & Krishnan-Sarin, S. (2021). School-based E-cigarette cessation programs: What do youth want? *Addictive Behaviors*, 125, 107167. DOI: <https://doi.org/10.1016/j.addbeh.2021.107167>
- Bonnie, R. J., Stratton, K., & Kwan, L. Y. (2015, July). Restrictions on Youth Access to Tobacco Products. Public Health Implications of Raising the Minimum Age of Legal Access to Tobacco Products - NCBI Bookshelf. <https://www.ncbi.nlm.nih.gov/books/n/nap18997/ch5>
- Bozzola, E., Spina, G., Agostiniani, R., Barni, S., Russo, R., Scarpato, E., Di Mauro, A., Di Stefano, A. V., Caruso, C., Corsello, G., & Staiano, A. (2022). The Use of Social Media in Children and Adolescents: Scoping Review on the Potential Risks. *International Journal of Environmental Research and Public Health*, 19(16), 9960. DOI: <https://doi.org/10.3390/ijerph19169960>
- Brubaker, S. (2007). Ethics and regulation in direct marketing. *Direct Marketing an International Journal*, 1(1), 55-58. DOI: <https://doi.org/10.1108/17505930710734134>
- Burkitt, H. (2007). Responsible drinks marketing: shared rights and responsibilities. *International Journal of Wine Business Research*, 19(3), 159-169. DOI: <https://doi.org/10.1108/17511060710817203>
- Carr, E. R. (2014). E-Cigarettes: Facts, Perceptions, and Marketing Messages. *Clinical Journal of Oncology Nursing*, 18(1), 112-116. DOI: <https://doi.org/10.1188/14.cjon.112-116>
- Centers for Disease Control and Prevention (US). (2012). Preventing Tobacco Use Among Youth and Young Adults. NCBI Bookshelf. <https://www.ncbi.nlm.nih.gov/books/NBK99237/>
- Chacón, L., Mitchell, G., & Golder, S. (2024). The commercial promotion of electronic cigarettes on social media and its influence on positive perceptions of vaping and vaping behaviours in Anglophone countries: A scoping review. *PLOS Global Public Health*, 4(1), e0002736. DOI: <https://doi.org/10.1371/journal.pgph.0002736>
- Chen, I., & Husten, C. G. (2014). Introduction to tobacco control supplement. *Tobacco Control*, 23(suppl 2), ii1-ii3. DOI: <https://doi.org/10.1136/tobaccocontrol-2013-051504>
- Cheng, H. G., Lizhnyak, P. N., & Richter, N. (2023). Mutual pathways between peer and own e-cigarette use among youth in the United States: A cross-lagged model. *BMC Public Health*, 23(1), 1609. DOI: <https://doi.org/10.1186/s12889-023-16470-5>
- Chu, K., Hershey, T. B., & Sidani, J. E. (2021). Collaborative Public Health Strategies to Combat e-Cigarette Regulation Loopholes. *JAMA Pediatrics*, 175(11), 1102. DOI: <https://doi.org/10.1001/jamapediatrics.2021.2432>
- Cohen, D. (1974). The Concept of Unfairness as It Relates to Advertising Legislation. *Journal of Marketing*, 38(3), 8. DOI: <https://doi.org/10.2307/1249844>
- Collaco, J. M., Drummond, M. B., & McGrath-Morrow, S. A. (2014). Electronic Cigarette Use and Exposure in the Pediatric Population. *JAMA Pediatrics*, 169(2), 177. DOI: <https://doi.org/10.1001/jamapediatrics.2014.2898>
- Collins, L., Glasser, A. M., Abudayyeh, H., Pearson, J. L., & Villanti, A. C. (2017). E-Cigarette Marketing and Communication: How E-Cigarette Companies Market E-Cigarettes and the Public Engages with E-cigarette Information. *Nicotine & Tobacco Research*, 21(1), 14-24. DOI: <https://doi.org/10.1093/ntr/ntx284>
- Costantino, S., Torre, A., Randazzese, S. F., Mollica, S. A., Motta, F., Busceti, D., Ferrante, F., Caminiti, L., Crisafulli, G., & Manti, S. (2024). Association between Second-Hand Exposure to E-Cigarettes at Home and Exacerbations in Children with Asthma. *Children*, 11(3), 356. DOI: <https://doi.org/10.3390/children11030356>
- Coxe, N., Webber, W., Burkhart, J., Broderick, B., Yeager, K., Jones, L., & Fenstersheib, M. (2014). Use of tobacco retail permitting to reduce youth access and exposure to tobacco in Santa Clara County, California. *Preventive Medicine*, 67, S46-S50. DOI: <https://doi.org/10.1016/j.ypmed.2014.01.023>
- Cullen, K. A., Liu, S. T., Bernat, J. K., Slavitt, W. I., Tynan, M. A., King, B. A., & Neff, L. J. (2019). Flavored Tobacco Product Use Among Middle and High School Students — United States, 2014-2018. *MMWR Morbidity and Mortality Weekly Report*, 68(39), 839-844. DOI: <https://doi.org/10.15585/mmwr.mm6839a2>
- Czaplicki, L., Kostygina, G., Kim, Y., Perks, S. N., Szczypka, G., Emery, S. L., Vallone, D., & Hair, E. C. (2019). Characterising JUUL-related posts on Instagram. *Tobacco Control*, 29-6, 612-617. DOI: <https://doi.org/10.1136/tobaccocontrol-2018-054824>

- Dai, H. (2018). Single, Dual, and Poly Use of Flavored Tobacco Products Among Youths. *Preventing Chronic Disease*, 15, E87. DOI: <https://doi.org/10.5888/pcd15.170389>
- Drummond, M. B., & Upson, D. (2014). Electronic Cigarettes. Potential Harms and Benefits. *Annals of the American Thoracic Society*, 11(2), 236-242. DOI: <https://doi.org/10.1513/annalsats.201311-391fr>
- Duke, J. C., Lee, Y. O., Kim, A. E., Watson, K. A., Arnold, K. Y., Nonnemaker, J. M., & Porter, L. (2014). Exposure to Electronic Cigarette Television Advertisements Among Youth and Young Adults. *PEDIATRICS*, 134(1), e29-e36. DOI: <https://doi.org/10.1542/peds.2014-0269>
- E-Cigarette Use Among Youth. (2024, October 17). Smoking and Tobacco Use. Retrieved October 20, 2024, from <https://e-cigarettes.surgeongeneral.gov/>
- Eltorai, A. E., Choi, A. R., & Eltorai, A. S. (2018). Impact of Electronic Cigarettes on Various Organ Systems. *Respiratory Care*, 64(3), 328-336. DOI: <https://doi.org/10.4187/respcare.06300>
- Ferkol, T. W., Farber, H. J., La Grutta, S., Leone, F. T., Marshall, H. M., Neptune, E., Pisinger, C., Vanker, A., Wisotzky, M., Zabert, G. E., & Schraufnagel, D. E. (2018). Electronic cigarette use in youths: a position statement of the Forum of International Respiratory Societies. *European Respiratory Journal*, 51(5), 1800278. DOI: <https://doi.org/10.1183/13993003.00278-2018>
- Gaiha, S. M., & Halpern-Felsher, B. (2020). Public Health Considerations for Adolescent Initiation of Electronic Cigarettes. *Pediatrics*, 145(Supplement 2), S175-S180. DOI: <https://doi.org/10.1542/peds.2019-2056e>
- Galderisi, A., Ferraro, V. A., Caserotti, M., Quareni, L., Perilongo, G., & Baraldi, E. (2020). Protecting youth from the vaping epidemic. *Pediatric Allergy and Immunology*, 31(S26), 66-68. DOI: <https://doi.org/10.1111/pai.13348>
- Glantz, S. A., & Bareham, D. W. (2018). E-Cigarettes: Use, Effects on Smoking, Risks, and Policy Implications. *Annual Review of Public Health*, 39(1), 215-235. DOI: <https://doi.org/10.1146/annurev-publhealth-040617-013757>
- Glantz, S. A., Nguyen, N., & Da Silva, A. L. O. (2024). Population-Based Disease Odds for E-Cigarettes and Dual Use versus Cigarettes. *NEJM Evidence*, 3(3), EVIDoA2300229. DOI: <https://doi.org/10.1056/evidoa2300229>
- Grana, R., Benowitz, N., & Glantz, S. A. (2014). E-Cigarettes. *Circulation*, 129(19), 1972-1986. DOI: <https://doi.org/10.1161/circulationaha.114.007667>
- Hammond, D., Reid, J. L., Rynard, V. L., Fong, G. T., Cummings, K. M., McNeill, A., Hitchman, S., Thrasher, J. F., Goniewicz, M. L., Bansal-Travers, M., O'Connor, R., Levy, D., Borland, R., & White, C. M. (2019). Prevalence of vaping and smoking among adolescents in Canada, England, and the United States: repeat national cross sectional surveys. *BMJ (Clinical Research Ed.)*, 365, I2219. DOI: <https://doi.org/10.1136/bmj.I2219>
- Harrell, M., Weaver, S., Loukas, A., Creamer, M., Marti, C., Jackson, C., Heath, J., Nayak, P., Perry, C., Pechacek, T., & Eriksen, M. (2016). Flavored e-cigarette use: Characterizing youth, young adult, and adult users. *Preventive Medicine Reports*, 5, 33-40. DOI: <https://doi.org/10.1016/j.pmedr.2016.11.001>
- Hartoyo, Y. R., & Sitorus, O. F. (2021). Utilization Instagram as a Marketing Communication Media in the Covid-19 Pandemic. *Advances in Social Science, Education and Humanities Research/Advances in Social Science, Education and Humanities Research*. DOI: <https://doi.org/10.2991/assehr.k.210430.030>
- Harvey, J., & Chadi, N. (2016). Preventing smoking in children and adolescents: Recommendations for practice and policy. *Paediatrics & Child Health*, 21(4), 209-214. DOI: <https://doi.org/10.1093/pch/21.4.209>
- Hildebr, D. (1994). Legal Aspects of Euro - marketing. *European Journal of Marketing*, 28(7), 44-54. DOI: <https://doi.org/10.1108/03090569410064454>
- Hunter, S. R., & Martin, C. A. (2019). Teen vaping boom: Electronic cigarette use, associated outcomes, and efforts in prevention. *Journal of the American Academy of Child & Adolescent Psychiatry*, 58(10), S71. DOI: <https://doi.org/10.1016/j.jaac.2019.07.435>
- Jackson, S. E., Tattan-Birch, H., East, K., Cox, S., Shahab, L., & Brown, J. (2024). Trends in Harm Perceptions of E-Cigarettes vs Cigarettes Among Adults Who Smoke in England, 2014-2023. *JAMA Network Open*, 7(2), e240582. DOI: <https://doi.org/10.1001/jamanetworkopen.2024.0582>
- Jenssen, B., & Boykan, R. (2019). Electronic Cigarettes and Youth in the United States: A Call to Action (at the Local, National and Global Levels). *Children*, 6(2), 30. DOI: <https://doi.org/10.3390/children6020030>
- Jenssen, B. P., Walley, S. C., Groner, J. A., Rahmandar, M., Boykan, R., Mih, B., Marbin, J. N., & Caldwell, A. L. (2019). E-Cigarettes and Similar Devices. *Pediatrics*, 143(2), e20183652. DOI: <https://doi.org/10.1542/peds.2018-3652>
- Kaisar, M. A., Prasad, S., Liles, T., & Cucullo, L. (2016). A decade of e-cigarettes: Limited research & unresolved safety concerns. *Toxicology*, 365, 67-75. DOI: <https://doi.org/10.1016/j.tox.2016.07.020>
- Kennedy, R. D., Awopegba, A., De León, E., & Cohen, J. E. (2016). Global approaches to regulating electronic cigarettes. *Tobacco Control*, 26(4), 440-445. DOI: <https://doi.org/10.1136/tobaccocontrol-2016-053179>
- Khambayat, S., Jaiswal, A., Prasad, R., Wanjari, M. B., Sharma, R., & Yelne, S. (2023). Vaping Among Adolescents: An Overview of E-Cigarette Use in Middle and High School Students in India. *Cureus*, 15(5), e38972. DOI: <https://doi.org/10.7759/cureus.38972>
- Klein, E. G., Berman, M., Hemmerich, N., Carlson, C., Htut, S., & Slater, M. (2016). Online E-cigarette Marketing Claims: A Systematic Content and Legal Analysis. *Tobacco Regulatory Science*, 2(3), 252-262. DOI: <https://doi.org/10.18001/trs.2.3.5>
- Kong, G., Bold, K. W., Cavallo, D. A., Davis, D. R., Jackson, A., & Krishnan-Sarin, S. (2020). Informing the development of adolescent e-cigarette cessation interventions: A qualitative study. *Addictive Behaviors*, 114, 106720. DOI: <https://doi.org/10.1016/j.addbeh.2020.106720>
- Kong, G., Morean, M. E., Cavallo, D. A., Camenga, D. R.,

- & Krishnan-Sarin, S. (2014). Reasons for Electronic Cigarette Experimentation and Discontinuation Among Adolescents and Young Adults. *Nicotine & Tobacco Research*, 17(7), 847-854. DOI: <https://doi.org/10.1093/ntr/ntu257>
- Krugman, D. M. (2016). Understanding the Impact That Marketing, Advertising, and Promotion Have on Adolescent E-cigarette Behavior. *Journal of Adolescent Health*, 59(6), 609-611. DOI: <https://doi.org/10.1016/j.jadohealth.2016.09.016>
- Le, T. T. H., Le, T. H., Le, M. D., & Nguyen, T. T. (2023). Exposure to E-Cigarette Advertising and Its Association with E-Cigarette Use Among Youth and Adolescents in Two Largest Cities in Vietnam 2020. *Tobacco Use Insights*, 16, 1179173X2311796. DOI: <https://doi.org/10.1177/1179173x231179676>
- Leslie, F. M. (2020). Unique, long-term effects of nicotine on adolescent brain. *Pharmacology Biochemistry and Behavior*, 197, 173010. DOI: <https://doi.org/10.1016/j.pbb.2020.173010>
- Liu, J., Gaiha, S. M., & Halpern-Felsher, B. (2022). School-based programs to prevent adolescent e-cigarette use: A report card. *Current Problems in Pediatric and Adolescent Health Care*, 52(6), 101204. DOI: <https://doi.org/10.1016/j.cppeds.2022.101204>
- Maddu, N. (2019). What is e-Cigarette and Associated Health Risks. In *IntechOpen eBooks*. DOI: <https://doi.org/10.5772/intechopen.84747>
- Marqués, P., Piqueras, L., & Sanz, M. (2021). An updated overview of e-cigarette impact on human health. *Respiratory Research*, 22(1), 151. DOI: <https://doi.org/10.1186/s12931-021-01737-5>
- Marynak, K., Gentzke, A., Wang, T. W., Neff, L., & King, B. A. (2018). Exposure to Electronic Cigarette Advertising Among Middle and High School Students — United States, 2014-2016. *MMWR Morbidity and Mortality Weekly Report*, 67(10), 294-299. DOI: <https://doi.org/10.15585/mmwr.mm6710a3>
- Marynak, K., Kenemer, B., King, B. A., Tynan, M. A., MacNeil, A., & Reimels, E. (2017). State Laws Regarding Indoor Public Use, Retail Sales, and Prices of Electronic Cigarettes — U.S. States, Guam, Puerto Rico, and U.S. Virgin Islands, September 30, 2017. *MMWR Morbidity and Mortality Weekly Report*, 66(49), 1341-1346. DOI: <https://doi.org/10.15585/mmwr.mm6649a1>
- Massey, Z. B., Brockenberry, L. O., & Harrell, P. T. (2020). Vaping, smartphones, and social media use among young adults: Snapchat is the platform of choice for young adult vapers. *Addictive Behaviors*, 112, 106576. DOI: <https://doi.org/10.1016/j.addbeh.2020.106576>
- Mattingly, D. T., & Hart, J. L. (2024). Trends in Current Electronic Cigarette Use Among Youths by Age, Sex, and Race and Ethnicity. *JAMA Network Open*, 7(2), e2354872. DOI: <https://doi.org/10.1001/jamanetworkopen.2023.54872>
- McCarthy, W. J., Mistry, R., Lu, Y., Patel, M., Zheng, H., & Dietsch, B. (2009). Density of Tobacco Retailers Near Schools: Effects on Tobacco Use Among Students. *American Journal of Public Health*, 99(11), 2006-2013. DOI: <https://doi.org/10.2105/ajph.2008.145128>
- Miller, S., Pike, J., Chapman, J., Xie, B., Hilton, B. N., Ames, S. L., & Stacy, A. W. (2016). The Prevalence and Marketing of Electronic Cigarettes in Proximity to At-Risk Youths: An Investigation of Point-of-Sale Practices Near Alternative High Schools. *Journal of Child & Adolescent Substance Abuse*, 26(2), 119-124. DOI: <https://doi.org/10.1080/1067828x.2016.1230529>
- Monteiro, M. G., Babor, T. F., Jernigan, D., & Brookes, C. (2017). Alcohol marketing regulation: from research to public policy. *Addiction*, 112(S1), 3-6. DOI: <https://doi.org/10.1111/add.13660>
- Morain, S. R., & Malek, J. (2017). Minimum Age of Sale for Tobacco Products and Electronic Cigarettes: Ethical Acceptability of US "Tobacco 21 Laws." *American Journal of Public Health*, 107(9), 1401-1405. DOI: <https://doi.org/10.2105/ajph.2017.303900>
- Murthy, V. H. (2016). E-Cigarette Use Among Youth and Young Adults. *JAMA Pediatrics*, 171(3), 209. DOI: <https://doi.org/10.1001/jamapediatrics.2016.4662>
- Nguyen, H., Dennehy, C. E., & Tsourounis, C. (2020). Violation of US regulations regarding online marketing and sale of e-cigarettes: FDA warnings and retailer responses. *Tobacco Control*, 29(e1), e4-e9. DOI: <https://doi.org/10.1136/tobaccocontrol-2019-055106>
- Padon, A. A., Maloney, E. K., & Cappella, J. N. (2016). Youth-Targeted E-cigarette Marketing in the US. *Tobacco Regulatory Science*, 3(1), 95-101. DOI: <https://doi.org/10.18001/trs.3.1.9>
- Papaleontiou, L., Agaku, I. T., & Filippidis, F. T. (2019). Effects of Exposure to Tobacco and Electronic Cigarette Advertisements on Tobacco Use: An Analysis of the 2015 National Youth Tobacco Survey. *Journal of Adolescent Health*, 66(1), 64-71. DOI: <https://doi.org/10.1016/j.jadohealth.2019.05.022>
- Paradise, J. (2014). Electronic Cigarettes: Smoke-Free Laws, Sale Restrictions, and the Public Health. *American Journal of Public Health*, 104(6), e17-e18. DOI: <https://doi.org/10.2105/ajph.2014.301890>
- Park-Lee, E., Ren, C., Cooper, M., Cornelius, M., Jamal, A., & Cullen, K. A. (2022). Tobacco Product Use Among Middle and High School Students — United States, 2022. *MMWR Morbidity and Mortality Weekly Report*, 71(45), 1429-1435. DOI: <https://doi.org/10.15585/mmwr.mm7145a1>
- Perelman, J., Leão, T., & Kunst, A. E. (2019). Smoking and school absenteeism among 15- to 16-year-old adolescents: a cross-section analysis on 36 European countries. *European Journal of Public Health*, 29(4), 778-784. DOI: <https://doi.org/10.1093/eurpub/ckz110>
- Public Health Consequences of E-Cigarettes. (2018). In *National Academies Press eBooks*. DOI: <https://doi.org/10.17226/24952>
- Saitta, D., Ferro, G. A., & Polosa, R. (2014). Achieving appropriate regulations for electronic cigarettes. *Therapeutic Advances in Chronic Disease*, 5(2), 50-61. DOI: <https://doi.org/10.1177/2040622314521271>
- Shamout, M., Tanz, L., Herzig, C., Oakley, L. P., Peak, C. M., Heinzerling, A., Hast, M., McGowan, E., Williams, R. J., Hess, C., Wang, C., Planche, S., Herndon, S., Martin, J., Kansagra, S. M., Al-Shawaf, M., Melstrom, P., Marynak, K., Tynan, M. A., . . . King, B. A. (2020). Notes from the Field: Characteristics of E-cigarette, or Vaping, Products Confiscated in Public High Schools in California and North

- Carolina — March and May 2019. *MMWR Morbidity and Mortality Weekly Report*, 69(42), 1552-1554. DOI: <https://doi.org/10.15585/mmwr.mm6942a7>
- Silver, N. A., Bertrand, A., Kucherlapaty, P., & Schillo, B. A. (2023). Examining influencer compliance with advertising regulations in branded vaping content on Instagram. *Frontiers in Public Health*, 10, 1001115. DOI: <https://doi.org/10.3389/fpubh.2022.1001115>
- Sindelar, J. L. (2020). Regulating Vaping — Policies, Possibilities, and Perils. *New England Journal of Medicine*, 382(20), e54. DOI: <https://doi.org/10.1056/nejmp1917065>
- Singh, S., Windle, S. B., Filion, K. B., Thombs, B. D., O'Loughlin, J. L., Grad, R., & Eisenberg, M. J. (2020). E-cigarettes and youth: Patterns of use, potential harms, and recommendations. *Preventive Medicine*, 133, 106009. DOI: <https://doi.org/10.1016/j.ypmed.2020.106009>
- Smith, M. J., Buckton, C., Patterson, C., & Hilton, S. (2023). User-generated content and influencer marketing involving e-cigarettes on social media: A scoping review and content analysis of YouTube and Instagram. *BMC Public Health*, 23(1), 530. DOI: <https://doi.org/10.1186/s12889-023-15389-1>
- Snell, L. M., Nicksic, N., Panteli, D., Burke, S., Eissenberg, T., Fattore, G., Gauci, C., Koprivnikar, H., Murauskiene, L., Reinap, M., & Barnes, A. J. (2021). Emerging electronic cigarette policies in European member states, Canada, and the United States. *Health Policy*, 125(4), 425-435. DOI: <https://doi.org/10.1016/j.healthpol.2021.02.003>
- Sreeramareddy, C. T., Acharya, K., & Manoharan, A. (2022). Electronic cigarettes use and 'dual use' among the youth in 75 countries: estimates from Global Youth Tobacco Surveys (2014-2019). *Scientific Reports*, 12(1), 20967. DOI: <https://doi.org/10.1038/s41598-022-25594-4>
- State Laws Prohibiting Sales to Minors and Indoor Use of Electronic Nicotine Delivery Systems — United States, November 2014. (2014, December 12). Retrieved October 2, 2024, from <https://www.cdc.gov/mmwr/preview/mmwrhtml/mm6349a1.htm>
- Stopping the start: our new plan to create a smokefree generation. (2023, November 8). GOV.UK. Retrieved September 25, 2024, from <https://www.gov.uk/government/publications/stopping-the-start-our-new-plan-to-create-a-smokefree-generation/stopping-the-start-our-new-plan-to-create-a-smokefree-generation>
- Struik, L. L., Dow-Fleisner, S., Belliveau, M., Thompson, D., & Janke, R. (2020). Tactics for Drawing Youth to Vaping: Content Analysis of Electronic Cigarette Advertisements. *Journal of Medical Internet Research*, 22(8), e18943. DOI: <https://doi.org/10.2196/18943>
- The Harmful Consequences of Vaping: A Public Health Threat. (2020). *Journal of Addictions Nursing*, 31(2), E3-E4. DOI: <https://doi.org/10.1097/jan.0000000000000342>
- Tremblay, M., Pluye, P., Gore, G., Granikov, V., Filion, K. B., & Eisenberg, M. J. (2015). Regulation profiles of e-cigarettes in the United States: a critical review with qualitative synthesis. *BMC Medicine*, 13(1), 130. DOI: <https://doi.org/10.1186/s12916-015-0370-z>
- Tworek, C., Schauer, G. L., Wu, C. C., Malarcher, A. M., Jackson, K. J., & Hoffman, A. C. (2014). Youth Tobacco Cessation. *American Journal of Preventive Medicine*, 47(2), S15-S27. DOI: <https://doi.org/10.1016/j.amepre.2014.05.009>
- United States Public Health Service. Office of the Surgeon General & National Center for Chronic Disease Prevention and Health Promotion (U.S.). Office on Smoking and Health. (2016). E-Cigarette Use Among Youth and Young Adults. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. https://www.cdc.gov/tobacco/data_statistics/sgr/e-cigarettes/pdfs/2016_sgr_entire_report_508.pdf
- Vallone, D. M., Cuccia, A. F., Briggs, J., Xiao, H., Schillo, B. A., & Hair, E. C. (2020). Electronic Cigarette and JUUL Use Among Adolescents and Young Adults. *JAMA Pediatrics*, 174(3), 277. DOI: <https://doi.org/10.1001/jamapediatrics.2019.5436>
- Vogel, E. A., Ramo, D. E., Rubinstein, M. L., Delucchi, K. L., Darrow, S. M., Costello, C., & Prochaska, J. J. (2020). Effects of Social Media on Adolescents' Willingness and Intention to Use E-Cigarettes: An Experimental Investigation. *Nicotine & Tobacco Research*, 23(4), 694-701. DOI: <https://doi.org/10.1093/ntr/ntaa003>
- Wagoner, K. G., Reboussin, D. M., King, J. L., Orlan, E., Ross, J. C., & Sutfin, E. L. (2019). Who Is Exposed to E-Cigarette Advertising and Where? Differences between Adolescents, Young Adults and Older Adults. *International Journal of Environmental Research and Public Health*, 16(14), 2533. DOI: <https://doi.org/10.3390/ijerph16142533>
- Wang, T. W., Gentzke, A. S., Creamer, M. R., Cullen, K. A., Holder-Hayes, E., Sawdey, M. D., Anic, G. M., Portnoy, D. B., Hu, S., Homa, D. M., Jamal, A., & Neff, L. J. (2019). Tobacco Product Use and Associated Factors Among Middle and High School Students — United States, 2019. *MMWR Surveillance Summaries*, 68(12), 1-22. DOI: <https://doi.org/10.15585/mmwr.ss6812a1>
- Wang, T. W., Neff, L. J., Park-Lee, E., Ren, C., Cullen, K. A., & King, B. A. (2020). E-cigarette Use Among Middle and High School Students — United States, 2020. *MMWR Morbidity and Mortality Weekly Report*, 69(37), 1310-1312. DOI: <https://doi.org/10.15585/mmwr.mm6937e1>
- Winickoff, J. P., Hartman, L., Chen, M. L., Gottlieb, M., Nabi-Burza, E., & DiFranza, J. R. (2014). Retail Impact of Raising Tobacco Sales Age to 21 Years. *American Journal of Public Health*, 104(11), e18-e21. DOI: <https://doi.org/10.2105/ajph.2014.302174>
- Xiang, S., Jia, T., Xie, C., Cheng, W., Chaarani, B., Banaschewski, T., Barker, G. J., Bokde, A. L. W., Büchel, C., Desrivieres, S., Flor, H., Grigis, A., Gowland, P. A., Brühl, R., Martinot, J., Martinot, M. P., Nees, F., Orfanos, D. P., Poustka, L., . . . Feng, J. (2023). Association between vmPFC gray matter volume and smoking initiation in adolescents. *Nature Communications*, 14(1), 4684. DOI: <https://doi.org/10.1038/s41467-023-40079-2>
- Zhu, S., Sun, J. Y., Bonnevie, E., Cummins, S. E., Gamst, A., Yin, L., & Lee, M. (2014). Four hundred and sixty brands of e-cigarettes and counting: implications for product regulation. *Tobacco Control*, 23(suppl 3),

iii3-iii9. DOI:

<https://doi.org/10.1136/tobaccocontrol-2014-051670>

Received: September 01, 2024 | Revised: September 22, 2024 | Accepted: September 30, 2024
