


ORAL CONTRAST IN DIAGNOSING ACUTE APPENDICITIS WITH CT SCAN; IS IT NECESSARY?



Ebtehal Althobaiti¹, Rozan Mirza¹, Ahmed Brinji¹, Nawal Barnawi¹, Awdah Akbar², Amar Akbar², Nourah ALSaleh², Ahmed Hawsawi² and Maher Ghazawi¹✉

¹Al-Noor Specialist Hospital, Department of Radiology, Makkah, Saudi Arabia

²Al-Noor Specialist Hospital, Department of General Surgery, Makkah, Saudi Arabia



ABSTRACT

BACKGROUND: The diagnosis of acute appendicitis clinically is challenging due to variant age groups at presentation; multidetector computed tomography (MDCT) has become the standard modality with high sensitivity and specificity. The American College of Radiology recommend the use of intravenous contrast-enhanced computed tomography for suspected cases but no explicit comment on the use of enteral contrast which create diversity in the literature for the appropriate MDCT protocols worldwide. The objective of this study is to compare the accuracy, sensitivity and specificity of variant MDCT protocols in diagnosing acute appendicitis.

METHODOLOGY: A retrospective cross-sectional study was conducted at Alnoor Specialist Hospital – Makkah between January 2016 and December 2020 including all patients who presented to the emergency department with clinical suspicion of acute appendicitis and performed MDCT scans with subsequent surgical intervention. A net of 385 MDCT scans were retrieved from hospital System (PACS). Data processing was made on exported coded excel sheet data and analyzed by SPSS 26.0 statistical software.

RESULTS: 385 MDCT scans with different protocols were collected (NECT, ORAL only, IV only, ORAL and IV). Excluding ones without final diagnose 355 MDCT scans were analyzed. Radiologists'ability to detect acute appendicitis was higher in patients administered both oral and IV contrast (sensitivity =92.44%), IV only (89.61%), NECT (82.35%), and oral contrast (73.47%), respectively. Diagnostic accuracy was higher in patients administered both oral and IV contrast (accuracy = 89.23%).

CONCLUSION: The present study demonstrates that the diagnostic accuracy and sensitivity in diagnosing acute appendicitis utilizing IV contrast are comparable to both IV and oral contrast; therefore, oral contrast administration can be omitted in routine appendicitis protocol. A higher specificity was found with the use of oral contrast due to the luminal filling of the appendix. Yet, it can be reserved for contraindicated cases to IV iodinated contrast administration. CT scan showed a hun-

Received 19 January 2023

Revised 5 February 2023

Accepted 7 February 2023

Published 5 May 2023

Corresponding Author

Maher Ghazawi, Al-Noor Specialist Hospital, Department Of Radiology, Makkah, Saudi Arabia. Contact: +966125665000 Ext. 1445, Mobile: +966503507706, Email: mghazawi@moh.gov.sa

Copyright: © 2023 The Author(s).

This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

eISSN: 1658-8959



dred percent sensitivity in diagnosing perforated appendicitis regardless of the used protocol.

ملخص البحث

نبذة:

يعد التشخيص السريري لالتهاب الزائدة الدودية الحاد أمراً صعباً نظراً لاختلاف الفئات العمرية عند المرضى; أصبح التصوير المقطعي المحوسب متعدد الأجهزة (MDCT) هو الطريقة القياسية ذات الحساسية والنوعية العالية. توصي المبادئ التوجيهية القائمة على الأدلة للكلية الأمريكية لعلم الأشعة (ACR) باستخدام التصوير المقطعي المحسن باستخدام الصبغة الوريدية (CECT) (IV) للحالات المشتبه بها من التهاب الزائدة الدودية الحاد ولكن لا يوجد توصيات صريحة تتعلق باستخدام الصبغة المعوية. الهدف من هذه الدراسة هو مقارنة دقة وحساسية ونوعية بروتوكولات MDCT المتغيرة في تشخيص التهاب الزائدة الدودية الحاد.

المنهجية المتبعة:

تم إجراء دراسة استعادية مقطعية في مستشفى النور التخصصي بمكة المكرمة بين يناير ٢٠١٦ وديسمبر ٢٠٢٠. تضمنت الدراسة جميع مرضى قسم الطوارئ المشتبهين بالتهاب الزائدة الدودية الحاد بناء على الفحص السريري والذين أجروا الأشعة المقطعية (MDCT) قبل الجراحة عن طريق استئصال الزائدة الدودية المفتوح أو بالمنظار. تم الحصول على ما مجموعه ٣٨٥ تصوير مقطعي (MDCT) من نظام أرشفة الصور والاتصالات بالمستشفى (PACS). تمت معالجة جميع البيانات وتصديرها على جداول (excel) المشفرة وتم تحليلها بواسطة حزمة البرامج الإحصائية (SPSS).

النتائج:

تم جمع ٣٨٥ فحص مقطعي (MDCT) بروتوكولات مختلفة تتضمن: الفحوصات بدون صبغة (NECT), الفحوصات المتباينة عن طريق الصبغة المعوية فقط (Oral), عن طريق الصبغة الوريدية فقط (IV), وعن طريق الصبغة الوريدية والصبغة المعوية معاً (IV and oral). تم استبعاد فحوصات الأشعة المقطعية التي احتوت على نتائج غير مؤكدة أو غير متوفرة ولم تحتسب ضمن قيم التحليل النهائية الخاصة بالبحث. ولذلك, انحصر التحليل على ٣٥٥ فحصاً مقطعيًا (MDCT). كانت قدرة أطباء الأشعة التشخيصية على اكتشاف التهاب الزائدة الدودية الحاد أعلى في المرضى الذين تم إعطاؤهم صبغة معوية ووريدية (IV, Oral). (الحساسية = 92,44%), يليها المرضى الذين تم إعطاؤهم صبغة وريدية IV فقط (89,61%), ومن ثم المرضى الذين لم يعطوا أي نوع من أنواع الصبغة (NECT) (82,35%), وأقل نسبة كانت للمرضى الذين أعطوا صبغة معوية (Oral) فقط (73,47%). كانت الدقة التشخيصية أعلى في المرضى الذين تم إعطاؤهم صبغة وريدية و معوية معاً (الدقة = 89,23%).

الخلاصة:

قيم الدقة والحساسية الخاصة بالتشخيص عند استخدام الصبغة الوريدية والمعوية, مقارنة لقيم التشخيص باستخدام الصبغة الوريدية فقط. ولذلك قد يمكن الاستغناء عن الصبغة المعوية في فحوصات الأشعة المقطعية الروتينية لالتهاب الزائدة الدودية الحاد. مع العلم بأن القيمة النوعية عند إعطاء صبغة معوية للتشخيص كانت أعلى لمساعدتها على ملء تجويف الزائدة الدودية. ويمكن تقنين الطريقة هذه على المرضى الذين لديهم حساسية تجاه استخدام الصبغات الوريدية.

Keywords: Appendicitis, contrast, computed tomography (CT)

1. INTRODUCTION

Diagnosing acute appendicitis clinically is challenging due to variant age groups at presentation; variable clinical scenarios were around 20% presented with atypical clinical features and the possibility of associated different abdominal pathology [1–3]. Furthermore, the number of removals of a normal appendix or so-called “negative appendectomy rates” was as high as 40 - 50% [4–6]. However, this rate has decreased since the introduction of radiological imaging due to the ability to detect normal appendix or alternative medical conditions. While still high, the acceptable negative appendectomy rates are around 15 – 25% [7–10].

Ultrasonography (US) has been used as the initial imaging modality for establishing the diagnosis of acute appendicitis in children and women of reproductive age for potential gynecological entities with a sensitivity range from 44 – 95% [11, 12]. However, widely known limitations of the US include operator dependence, being affected by gaseous distension of the abdomen, and not reliable in detecting the normal or perforated appendix [12, 13].

Therefore, multidetector computed tomography (MDCT) has become the standard modality with sensitivity and specificity ranging from 88 – 100% and 91 – 99%, respectively [12]. The American College of Radiology (ACR) appropriateness criteria evidence-based guidelines recommend the use of intravenous (IV) contrast-enhanced computed tomography (CECT) for suspected cases of acute appendicitis but no explicit comment on the use of enteral contrast and refer it to the propensity of the institutions which create diversity in the literature for the appropriate MDCT protocols worldwide with different institutional experiences [14–16].

Unfortunately, there was no available local data to compare different MDCT scan protocols when acute appendicitis is suspected. Therefore, the present study aims to determine the sensitivity, specificity, and accuracy of MDCT in diagnosing acute appendicitis with variant-used protocols, as well as building a local institutional experience to protocol acute appendicitis cases.

2. MATERIALS AND METHODS

2.1 Study Population and Setting

The investigation was approved by the Institutional Board Review (IRB) of the general directorate of health affairs in Makkah – Saudi Arabia. A retrospective cross-sectional study was conducted at Alnoor Specialist Hospital – Makkah between January 2016 and December 2020. The study included all the patients who presented to the emergency

department with a clinical impression of acute appendicitis and performed preoperative MDCT scans with subsequent surgical intervention by either open or laparoscopic appendectomy. A total number of 385 MDCT scans were retrieved from the hospital Picture Archiving and Communication System (PACS).

2.2 Study Technique and Image Interpretation

All patients were scanned using a 256-slice MDCT scanner (SIEMENS, SOMATOM Definition Flash). Image acquisition is made from lung bases to the greater trochanter of the femur. A slice thickness of 2 mm axial images with 3 mm multiplanar coronal and sagittal reformats were obtained. Different protocols were used, including non-enhanced CT (NECT) scans, Oral only, Intravenous (IV) only, or both oral and IV contrast agents. Per institutional protocol, a (2 ml/kg of iohexol 350 mg I/ml or iodixanol 320 mg I/mL, GE Healthcare) of IV contrast was administered by a power injector (Stellant CT Injection System, Medrad, Indianola, PA) at a rate of 3-4 mL/s with a time delay of 70 s after injection followed by a 30 mL bolus of chasing normal saline. For patients who received oral contrast, preparation of (30 mL of Gastrografin 37% Solution 12x120 ML, Bracco Diagnostic) was diluted in 1 – 1.2 L of water and administered 2 hours before image acquisition.

Two Saudi board-certified radiologists were enrolled in this research. They reviewed all the MDCT scans and completed a pre-designed structural report containing the used protocol, demographic data, specific data regarding the appendix, and their final impression regarding each case. The examiners were blinded to the final diagnosis, intraoperative findings, and histopathology report. Finally, the data was exported to an excel sheet and Statistical Package for the Social Sciences (SPSS) for analysis.

The presence of acute appendicitis depends on the following CT findings: Dilated appendix (diameter > 6 mm), mural thickening, mucosal hyperenhancement, appendicolith, absence of intraluminal gas or contrast, peri-appendiceal fat stranding, and free fluid. Complicated appendicitis with inflammatory mass, perforation, or collection was also documented. The final impression was fallen into one of the following categories; acute non-complicated appendicitis, perforated appendicitis, appendicitis with collection, appendicitis with malignancy, equivocal appendicitis, stump appendicitis, or normal appendix. Final CT diagnoses were compared with operative and pathology reports.

2.3 Statistical Analysis

All data processing was made on exported coded excel sheet data and analyzed by SPSS 26.0 statistical software package. Data were presented as frequencies or means \pm standard deviation. The diagnostic sensitivity, specificity, positive predictive value (PPV), negative

predictive value (NPV), and diagnostic accuracy were calculated from the four different protocols. A value of $p < 0.05$ was considered to indicate statistical significance.

3. RESULTS

385 MDCT scans of the abdomen and pelvis utilizing different protocols were collected (NECT, ORAL only, IV only, ORAL and IV). Final impressions of diagnoses that were not available were excluded from the final analysis. Therefore, 355 MDCT scans were analyzed: 89 (25.07%) in the NECT group, 54 (15.21%) in the oral group, 82 (23.01%) in the IV group, and 130 (36.62%) in the oral and IV contrast groups. A total of 232 (65.35%) male patients and 123 (34.65%) female patients data were analyzed in this study. No significant difference was observed in the gender and age mean differences with different protocols used (Table 1).

Table 1 Characterization of patients across the four protocols:

	NECT (89)	ORAL ONLY (54)	IV ONLY (82)	ORAL+IV(130)
Gender				
Male	55 (61.8%)	35 (64.8%)	57 (69.5%)	85 (65.4%)
Female	34 (38.2%)	19 (35.2%)	25 (30.5%)	45 (34.6%)
Age				
Mean± SD	36.42 ± 13.98	38.70 ± 15.84	37.87 ± 13.23	38.11 ± 14.76

All MDCT scans of the abdomen and pelvis were assessed for acute appendicitis. True positives and negatives, false positives and negatives were determined by comparing the CT scan findings to the final histopathology reports. Of the 295 patients with positive CT findings for appendicitis, 285 were confirmed by correlating the radiological findings with histopathological findings. Among the 285 true positive results, 70 were NECT, 36 were oral, 69 were IV, and 110 were both oral and IV, respectively. There were 58 patients with a normal appendix. Other patients were diagnosed with perforated appendicitis, appendicitis with collection, and an appendix with malignancy. Radiologists' ability to detect acute appendicitis was higher in patients administered both oral and IV contrast (sensitivity = 92.44%), following IV only (89.61%), NECT (82.35%), and oral contrast (73.47%), respectively (Table 2). Diagnostic accuracy was higher in patients administered both oral and IV contrast (89.23%).

The MDCT findings of patients who received oral contrast showed more true negative results (specificity =80%) than other protocols. In the final impression of radiologists correlating radiological and histopathological findings, the most frequently diagnosed conditions were acute non-complicated appendicitis (309 patients) and perforated appendicitis

Table 2 Diagnostic performance of the utilized protocol (NECT, oral only, IV only, and both oral and IV contrast):

	NECT (95% CI)	ORAL ONLY (95% CI)	IV ONLY (95% CI)	ORAL+IV (95% CI)
Sensitivity	82.35% (72.57-89.77)	73.47% (58.92-85.05)	89.61% (80.55-95.41)	92.44% (86.13-96.48)
Specificity	75% (19.41-99.37)	80% (28.36-99.49)	40% (5.27-85.34)	54.5% (23.38-83.25)
Accuracy	82.02% (72.45-89.36)	74.07% (60.35- 85.04)	86.59% (77.26-93.11)	89.23% (82.59-93.99)
PPV	98.59% (92.75-99.74)	97.30% (86.09-99.52)	95.83% (91.80 -97.93)	95.65% (91.99-97.68)
NPV	16.67% (72.45-89.36)	23.53% (13.96-36.84)	20.00% (6.63-46.80)	40.00% (22.56-60.41)

PPV; Positive Predictive Value; NPV; Negative Predictive Value; CI; Confidence Interval.

(12 cases). All 12 patients with perforated appendices were correctly diagnosed by CT scan. Hence, the sensitivity of diagnosing the perforated appendix was 100% in each protocol (Table 3). There was a total of 45 false-negative findings of MDCT. Fifteen were administered NECT, thirteen were oral, eight were IV contrast and nine were oral and IV administered patients (Table 4).

Table 3 True positive rates of diagnosing acute non-complicated appendicitis and perforated appendicitis:

	NECT (95% CI)	ORAL ONLY (95% CI)	IV ONLY (95% CI)	ORAL+IV (95% CI)
Acute non-complicated appendicitis	82.93 (73.02–90.34)	75% (59.66–86.81%)	89.04% (79.54–95.15)	92.73 (86.17–96.81)
Perforated appendicitis	100%	100%	100%	100%

Table 4 False negative diagnoses in different protocols:

	No. of patients
NECT	
Acute non-complicated appendix	14
Acute appendix with malignancy	1
ORAL only	
Acute non-complicated appendix	11
Acute appendix with malignancy	2
IV only	
Acute non-complicated appendix	8
ORAL +IV	
Acute non-complicated appendix	8
Acute appendix with malignancy	1

4. DISCUSSION

The value of MDCT in diagnosing acute appendicitis has been the subject of many studies worldwide. Various protocols have been used, including NECT, enteral (oral or rectal), IV or combination IV, and enteral contrast [17]. Reducing the field of view using a focused CT scan at the right lower quadrant to decrease radiation dose has also been suggested [16].

The present study shows a higher ability to detect acute appendicitis in patients administered both oral and IV contrast with a sensitivity of 92.44%, followed by IV only (sensitivity of 89.61%); however, there was no significant difference (P-value = 0.43) which is consistent with the established literature that a CT scan utilizing oral contrast does not improve the diagnostic accuracy of acute appendicitis [7, 17, 18]. A study published in 2016 concluded that the use of oral contrast in diagnosing acute appendicitis might be associated with a higher pulmonary complication rate, including pneumonia and acute respiratory distress syndrome as 93.75% of those who had complications received oral contrast compared to those who had IV contrast only (P > 0.0001). It also delays patients' transition from the emergency department to the operating room by an average of 4 hours [19].

In our study, the overall specificity values were low, ranging from 40 – 80%, which is expected as the inclusion criteria were all the patients who performed appendectomy (the true negative cases were only 58 from a total of 355). The specificity in the literature ranged from 97 to 100% as they included all the patients who presented to the emergency department with acute non-traumatic abdominal pain and were suspected clinically to have acute appendicitis. The standard reference for the final diagnosis was either surgical and histopathological correlation or specific follow-up criteria to ensure the resolution of symptoms [14, 20, 21]. The highest specificity in our study was 80% in the orally administered contrast agent. The reason mentioned in the literature is that the opacification of the appendix with oral contrast helps exclude acute appendicitis [5, 17, 22]. Keyzer et al. found that the visualization of the normal appendix in the oral-contrast group was 81 – 96% compared to 74 – 85% in the no-oral contrast group [21].

Perforated appendicitis is considered when there is discontinuity of the appendiceal wall, extraluminal air or appendicolith, contrast leak, or abscess formation. Our study demonstrates 100% sensitivity in detecting perforated appendicitis with no statistical difference between different used protocols. A 94.9% sensitivity was documented by Horrow et al. in diagnosing perforated appendicitis, considering a focal defect in the enhancing appendiceal wall is the most sensitive indicator of perforation [23].

This study has several limitations, including its retrospective nature. The pre-knowledge of participated radiologists about the research aims in diagnosing acute appendicitis might guide them toward the diagnosis rather than if it was a general indication as acute

abdomen. Lastly, the inclusion criteria limit the number of the normal appendix as we only included all the patients who did appendectomy following high clinical suspicion, laboratory and radiological investigations and ruled out a possible differential diagnosis, resulting in only a small number of true negative cases which affect the specificity in our study when compared to the literature. However, the chosen criteria allowed us to actually compare the sensitivity of each protocol to a final diagnosis confirmed operatively and histopathologically.

5. CONCLUSION

The present study demonstrates that the diagnostic accuracy and sensitivity in diagnosing acute appendicitis utilizing IV contrast are comparable to both IV and oral contrast; therefore, oral contrast administration can be omitted in routine appendicitis protocol. A higher specificity was found with the use of oral contrast due to the luminal filling of the appendix. Yet, it can be reserved for contraindicated cases to IV iodinated contrast administration. CT scan showed a hundred percent sensitivity in diagnosing perforated appendicitis regardless of the used protocol.

CONFLICT OF INTEREST

There is no conflict of interest to declare

ACKNOWLEDGEMENTS

N/A

REFERENCES

- [1] Iwahashi N, Kitagawa Y, Mayumi T, Kohno H. Intravenous Contrast-enhanced Computed Tomography in the Diagnosis of Acute Appendicitis. *World J Surg.* 2005;29(1):83–7.
- [2] Raman SS, Lu DSK, Kadell BM, Vodopich DJ, Sayre J, Cryer H. Accuracy of Nonfocused Helical CT for the Diagnosis of Acute Appendicitis: A 5-Year Review. *Am J Roentgenol.* 2002;178(6):1319–25.
- [3] Naffaa LN, Ishak GE, Haddad MC. The value of contrast-enhanced helical CT scan with rectal contrast enema in the diagnosis of acute appendicitis. *Clin Imaging.*

2005;29(4):255–8.

- [4] Glauser J, Siff J, Emerman C. Emergency Department Experience With Nonoral Contrast Computed Tomography in the Evaluation of Patients for Appendicitis. *J Patient Saf.* 2014;10(3):154–8.
- [5] Drake FT, Alfonso R, Bhargava P, Cuevas C, Dighe MK, Florence MG et al. Enteral Contrast in the Computed Tomography Diagnosis of Appendicitis: Comparative Effectiveness in a Prospective Surgical Cohort. *Ann Surg.* 2014;260(2):311–6.
- [6] Tang SJ, Pickhardt PJ, Kim DU, Pooler BD. Positive Oral Contrast Solution at MDCT for Suspected Acute Appendicitis in Adults: Rate of Appendiceal Luminal Filling of Normal and Inflamed Appendixes. *Am J Roentgenol.* 2019;213(5):W211–7.
- [7] Anderson BA, Salem L, Flum DR. A systematic review of whether oral contrast is necessary for the computed tomography diagnosis of appendicitis in adults. *Am J Surg.* 2005;190(3):474–8.
- [8] Hershko DD, Awad N, Fischer D, Mahajna A, Guralnik L, Israelit SH et al. Focused Helical CT Using Rectal Contrast Material Only as the Preferred Technique for the Diagnosis of Suspected Acute Appendicitis: A Prospective, Randomized, Controlled Study Comparing Three Different Techniques. *Dis Colon Rectum.* 2007;50(8):1223–9.
- [9] Hlibczuk V, Dattaro JA, Jin Z, Falzon L, Brown MD. Diagnostic Accuracy of Non-contrast Computed Tomography for Appendicitis in Adults: A Systematic Review. *Ann Emerg Med.* 2010;55(1):51–59.e1.
- [10] Tamburrini S, Brunetti A, Brown M, Sirlin C, Casola G. Acute appendicitis: diagnostic value of nonenhanced CT with selective use of contrast in routine clinical settings. *Eur Radiol.* 2007;17(8):2055–61.
- [11] Mullins ME, Kircher MF, Ryan DP, Doody D, Mullins TC, Rhea JT et al. Evaluation of Suspected Appendicitis in Children Using Limited Helical CT and Colonic Contrast Material. *Am J Roentgenol.* 2001;176(1):37–41.
- [12] Pinto Leite N, Pereira JM, Cunha R, Pinto P, Sirlin C. CT Evaluation of Appendicitis and Its Complications: Imaging Techniques and Key Diagnostic Findings. *Am J Roentgenol.* 2005;185(2):406–17.
- [13] Fefferman NR, Roche KJ, Pinkney LP, Ambrosino MM, Genieser NB. Suspected Appendicitis in Children: Focused CT Technique for Evaluation. *Radiology.* 2001;220(3):691–5.

- [14] Kepner AM, Bacasnot JV, Stahlman BA. Intravenous contrast alone vs intravenous and oral contrast computed tomography for the diagnosis of appendicitis in adult ED patients. *Am J Emerg Med.* 2012;30(9):1765–73.
- [15] Farrell CR, Bezinque AD, Tucker JM, Michiels EA, Betz BW. Acute appendicitis in childhood: oral contrast does not improve CT diagnosis. *Emerg Radiol.* 2018;25(3):257–63.
- [16] Tarulli M, Rezende-Neto J, Vlachou PA. Focused CT for the evaluation of suspected appendicitis. *Abdom Radiol.* 2019;44(6):2081–8.
- [17] Hekimoglu K, Yildirim UM, Karabulut E, Coskun M. Comparison of combined oral and iv contrast-enhanced versus single iv contrast-enhanced mdct for the detection of acute appendicitis. *J Belg Soc Radiol.* 2011;94(5):278.
- [18] Ramalingam V, Bates DDB, Buch K, Uyeda J, Zhao KM, Storer LA et al. Diagnosing acute appendicitis using a nonoral contrast CT protocol in patients with a BMI of less than 25. *Emerg Radiol.* 2016;23(5):455–62.
- [19] Saadat L, Helenowski I, Mahvi D, Boller AMM. Should Oral Contrast Be Omitted in Patients with Suspected Appendicitis? *J Gastrointest Surg.* 2016;20(6):1150–3.
- [20] Wadhvani A, Guo L, Saude E, Els H, Lang E, McRae A et al. Intravenous and Oral Contrast vs Intravenous Contrast Alone Computed Tomography for the Visualization of Appendix and Diagnosis of Appendicitis in Adult Emergency Department Patients. *Can Assoc Radiol J.* 2016;67(3):234–41.
- [21] Keyzer C, Cullus P, Tack D, De Maertelaer V, Bohy P, Gevenois PA. MDCT for Suspected Acute Appendicitis in Adults: Impact of Oral and IV Contrast Media at Standard-Dose and Simulated Low-Dose Techniques. *Am J Roentgenol.* 2009;193(5):1272–81.
- [22] Naeger DM, Chang SD, Kolli P, Shah V, Huang W, Thoeni RF. Neutral vs positive oral contrast in diagnosing acute appendicitis with contrast-enhanced CT: sensitivity, specificity, reader confidence and interpretation time. *Br J Radiol.* 2011;84(1001):418–26.
- [23] Horrow MM, White DS, Horrow JC. Differentiation of Perforated from Nonperforated Appendicitis at CT. *Radiology.* 2003;227(1):46–51.