

Guy's Stone Scoring in Predicting Postoperative Residual Stone

Muhammad Waqas Iqbal, Farhan Jamshed, Adeen Akram, Nauman Khalid,
Muhammad Saifullah, Muhammad Akram Malik

ABSTRACT

Objective: To determine the frequency of postoperative residual stone based on preprocedural Guy's scoring system.

Methodology: This cross-sectional descriptive study was conducted from 1st June 2018 to 30th November 2019 at University Medical College Faisalabad. All patients with stones in the kidney undergoing PCNL, 20 to 60 years of age of both genders were included. Patients with secondary stones due to pelviureteric junction obstruction, CRF and recurrent stones were excluded. In all patients Guy's scoring was measured. Then all patients underwent PCNL. After 24 hours of operation, CT scan was done in each patient and residual stone (present/absent) was noted.

Results: This study was comprised of 91 patients of age ranged from 20- 60 years. Fifty-three patients (58.24%) were males and 38 (41.76%) females. Seven (7.69%) were of Guy's grade I, 48(52.75%) grade II, 21(23.08%) grade III and 15 of grade IV. Number of patients with residual stone was 09 (9.89%), 0.0% of GS 1, 12.5% of GS 2, 4.76% of GS 3 and 13.33% of GS 4 patients had residual stone.

Conclusion: Pre-PCNL Guy's scoring is easy, rapid and useful for predicting the residual stone after percutaneous nephrolithotomy.

KEYWORDS: Residual stone, Guy's scoring system, Extracorporeal shockwave lithotripsy

INTRODUCTION

Nephrolithiasis is a common disorder with estimated prevalence of 1-13% in various regions of the world.¹ Precipitating factors of kidney stone disease include age, gender, diet, lifestyle, socioeconomic status, geographical location and race/ethnicity resulting in various incidence rates in different countries.^{2,3} Urinary stone disease has affected mankind since antiquity. It has been even found in Egyptian Mummies.⁴ Nephrolithiasis is a recurrent disorder. In some studies recurrence rate has been

reported from 21% to 53%. Most of the stones are of calcium oxalate (75%-90%). Other components were uric acid (5%-20%), calcium phosphate (6%-13%) and struvite (2%-15%).^{5,6}

New techniques have been developed for the management of renal stones with the advances in technology in this modern era. Extracorporeal shockwave lithotripsy (ESWL), Percutaneous nephrolithotomy (PCNL), Retrograde intrarenal surgery (RIRS), and Laparoscopic stone surgeries have replaced open surgical procedures in majority of cases.⁷ Percutaneous Nephrolithotomy (PCNL) is now considered as first line standard treatment for large renal stones >2cm. As the incidence of renal stone is increasing, procedure of PCNL has become increasingly popular. Stone free rate after PCNL and its complications are topic of discussion due to increasing number of procedures in the world. The objective and goal of PCNL is the provision of maximum stone clearance with minimal rate of complications.⁷ Different preoperative factors which affect the outcome of PCNL are size & site of the stone, patient's characteristics and renal anatomy.⁸

To date computed tomography (CT) is considered as the most important imaging investigation in renal stones prior to PCNL. CT plays a very important role in preoperative assessment of stone's characteristics and renal anatomy. Various scoring systems like Guy's scoring, CROES nomogram and S.T.O.N.E., are based on findings of preoperative CT.^{9,10} Aims of the study was to determine the frequency of

Dr. Muhammad Waqas Iqbal MBBS, FCPS

Assistant Professor

University Medical College, FSD

Dr. Farhan Jamshed MBBS, FCPS

Senior Registrar

University Medical College, FSD

Dr. Adeen Akram MBBS

Postgraduate Resident (PGR)

District Headquarter (DHQ) Hospital, FSD

Dr. Nauman Khalid MBBS, FCPS

Assistant Professor

University Medical & Dental College, FSD

Dr. Muhammad Saifullah MBBS, FCPS

Senior Registrar

District Headquarter (DHQ) Hospital, FSD

Dr Muhammad Akram Malik MBBS, FCPS

Professor

University Medical College, FSD

Correspondence:

Dr. Muhammad Akram Malik

Email: akrammlk@yahoo.com

postoperative residual stone based on pre-procedural Guy's scoring system.

METHODOLOGY

It was a descriptive study conducted at University Medical College Faisalabad from June, 01, 2018 to November, 30, 2019. All patients with kidney stones of either gender, with age range 20-60 years were included into this study. Diagnosis was confirmed by plain computerized tomography (CT). Ethical approval was taken from ethical approval committee (TUF/Dean/2018 dated May, 15, 2018).

Preoperatively Guy's scoring was done for each patient as follows: Guy's stone score 1 (GSS1): a solitary stone in the mid/and or lower pole or in the renal pelvis with a normal anatomy and simple collecting system Guy's stone score 2 (GSS2): a solitary stone in the upper pole; multiple stones in patients with simple anatomy; or a solitary stone in a patient with abnormal anatomy Guy's stone score 3 (GSS3): multiple stones in a patient with abnormal anatomy or in a calyceal diverticulum or partial staghorn calculus Guy's stone score 4 (GSS4): a complete staghorn calculus or any stone in a patient with spinal bifida or a spinal injury, calculus in patients with clinical neurological alternations (spinal cord injury, myelomeningocele) All patients underwent PCNL by experienced consultant. CT-scan was performed on 1st postoperative day to see any residual stone >2mm. Patients with secondary stones due to pelviureteric junction obstruction as assessed on Computed Tomography Urography (CTU), of chronic renal failure (s/creatinine >1.5 mg/dl) and with recurrent stones were excluded from study. Morbidly obese patients (BMI >37 kg/m²) and the Patients in which procedure was converted to open pyelolithotomy were also excluded.

Statistical analysis: Social Science (SPSS) Version 20.0 was used to analyze the entered data. Quantitative data were presented as mean ± SD. Qualitative data were presented as frequency and percentage. Pre-operative Guy's grading (1/2/3/4) and effect modifiers like age, gender & BMI were stratified. Post-stratification chi-square was used to assess their effect on postoperative residual stone and p-value ≤0.05 was taken as significant.

RESULTS

Total no of patients in this study was 91. Age ranged from 20- 60 years with mean age of 35.64 ± 8.12 years.


Characteristics of Patients according to age, gender and BMI is shown in table 1. Male to female ratio was 1.4:1. Mean BMI was 27.73 ± 2.86 kg/m².

Table 1: Characteristics of Patients According to Age, Gender and BMI (N=91)

| Groups | | No. of Patients (N) | Percentages (%) |
|--------------------------|--------|---------------------|-----------------|
| Age | 20-40 | 69 | 75.82 |
| | 41-60 | 22 | 24.18 |
| Gender | Male | 53 | 58 |
| | Female | 38 | 42 |
| BMI (kg/m ²) | ≤27 | 44 | 48.35 |
| | >27 | 47 | 51.65 |

Fifty three patients (58.24%) were males and 38 (41.76%) females. Male to female ratio was 1.4:1. Mean BMI was 27.73 ± 2.86 kg/m² (Table I). Mean preoperative Guy's grading was 2.48 ± 0.86. Distribution of patients according to Preoperative Guy's grading is shown in (Figure 1).

Figure 1: Distribution of Patients According to Pre-Operative Guy's Grading (n=91).



GSS=Guy's stone score

Residual stone was present in 09 (9.89%) with none in grade 1 and 2(15.38%) in grade 4(table 2).

Table 2: Stratification of Residual Stone with Respect to Pre-Operative Guy's Grading (n=91).

| Pre-operative Guy's grading | Residual stone | | P-Value |
|-----------------------------|----------------|--------------|---------|
| | Present N (%) | Absent N (%) | |
| Grade 1 | 00 | 7 (100) | 0.582 |
| Grade 2 | 06 (12.5) | 42 (87.5) | |
| Grade 3 | 01 (4.7) | 20 (95.2) | |
| Grade 4 | 02 (13.3) | 13 (86.6) | |

p-value ≤ 0.05 taken as significant.

Stratification of residual stone with respect to age, gender and BMI shows greater number of residual stones in age group of 41-60 years (Table 3).

Table 3: Stratification of Postoperative Residual Stone with Respect to Age, Gender and BMI Groups (n=91).

| Groups | Residual stone | | P-values |
|--------------------------|----------------|----------|----------|
| | Present N | Absent N | |
| Age | 20-40 | 04 | 0.021* |
| | 41-60 | 05 | |
| Gender | Male | 04 | 0.377 |
| | Female | 05 | |
| BMI (kg/m ²) | ≤ 27 | 06 | 0.247 |
| | > 27 | 03 | |

p-value ≤ 0.05 taken as significant.

DISCUSSION

Percutaneous nephrolithotomy (PCNL) is being considered as procedure of choice for large and complex renal calculi and has replaced the open surgical procedures to greater extent.¹¹ Although PCNL is an endoscopic procedure with higher success rate and minimal complications, however stone free rate is not 100%.¹² Stone size, burden, and location, renal calyceal anatomy and associated hydronephrosis are different parameters which predict the outcome of PCNL. However, when these parameters are used individually, precise outcome of the procedure cannot be predicted.¹³ Due to this reason various scoring systems have been invented to predict the outcome of PCNL. These scoring systems are based on preoperative stone size/burden, renal calyceal anatomy and patient's characteristics (age/gender/body built/BMI).¹⁴ Besides prediction of stone free rate, counseling of patient, refinement of procedure and adjustments of various training programs can be managed by these scoring.¹⁵ Various scoring systems have been developed to predict stone free rate and complications. Guy's scoring system, the S.T.O.N.E. scoring system, the S-ReSC Scoring System of the

Seoul National University and the CROES, are among the most commonly applied and evaluated. Guy's scoring is easy to perform, simple and rapid to predict the outcome of PCNL.¹⁶⁻¹⁹ In our study, the percentage of patients with residual stone was 09 (9.89%). 0.0% of GS 1, 12.5% of GS 2, 4.76% of GS 3 patients and 13.33% of GS 4 had residual stone. In a study, 75% of the subjects were GS 1, 21% GS 2 and 3% GS 3. In this study, 66% of GS 3, 4% of GS 2 and 10% of GS had residual stone.²⁰ Another published series reported residual stones in 70% of GS 1, 65% of GS 2 and 38% of GS 3 persons.²¹ Results of our study are comparable with these studies, however with better outcome in present study. Vicentini et al reported rates of 95% and 79%, 59% and 41% of stone-free patients in GS1, GS2, GS 3 and GS 4 scoring system respectively in their series involving 147 PCNL patients, and elaborated a significant association of Guy scores and stone-free rate.¹³ Kay Thomas evaluated residual stones in 100 patients who underwent PCNL, and found residual stones in 19% of grade I, 28% of grade II, 65% of grade III and 71% of grade IV.¹⁸ Results obtained by Kay Thomas and Vicentini et al are contradictory to the study conducted by Mandal et al¹⁹, who found stone free rate of 0%, 56%, 68%, 74%, for grade 1-4 respectively. As reported in all above mentioned series, Guy's scoring system precisely discriminates between the two endpoints. Guy's scoring system briefly creates two group of grades, good and bad. Grades 1 and 2 are considered as good and grades 3 and 4 are relatively bad group. Our study has age range from 20 - 60 years (mean = 35.64 ± 8.12 years). As the age and clearance of renal stone is concerned there is significant difference in terms of residual stone among two age groups. 29% of patients in age group 41-60 have residual stone as compared to 6% in age group 20-40 years ($p < 0.021$). In a study conducted by Rahul Gupta there is no difference in different age groups as far as stone free rate is considered (94% vs 92% respectively in older & young).²² Most probable causative factor for higher residual stone in the older age group in our study is obesity.²³ No statistically significant difference in terms of residual stone with respect to gender (8% in Male and 13% in Female) and weight (14% in BMI ≤ 27 and 6% in BMI > 27). These results are comparable to other series.^{23, 24} The Guy score has been validated externally in a large number of published series. In two separate series by Mandal and Ingimarsson, Guy's scoring system effectively predicted SFR and has been externally validated.^{19,25} The Guy score has excellent interobserver reliability.²⁵

CONCLUSION

Pre-PCNL Guy's is useful for predicting the residual stone after the procedure. We recommend that this scoring system can be applied routinely in patients undergoing PCNL for pre-operative evaluation of surgical outcome so that patients can be counseled pre-operatively for chances of residual stone.

Disclaimer: This study was a part of dissertation for FCPS part II, a mandatory requirement of eligibility to appear in examination.

Conflict of Interest: None

Funding Source: None

REFERENCES

- Sorokin I, Mamoulakis C, Miyazawa K, Rodgers A, Talati J, Lotan Y. Epidemiology of stone disease across the world. *World J Urol.* 2017;35(9):1301-1320. doi:10.1007/s00345-017-2008-6
- Fakhoury MQ, Gordon B, Shorter B, Renson A, Borofsky MS, Cohn MR et al. Perceptions of dietary factors promoting and preventing nephrolithiasis: a cross sectional survey. *World J Urol.* 2019;1;37(8):1723-1731. doi: 10.1007/s00345-018-2562-6
- Jadoon S, Wang J, Mahmood Q, Li X D, Zeb BS, Naseem I, et al. Association of nephrolithiasis with drinking water quality and diet in Pakistan. *Environmental Engineering and Management Journal.* 2020;19(8), 1289-1297
- Ferakis N, Stavropoulos M. Mini percutaneous nephrolithotomy in the treatment of renal and upper ureteral stones: Lessons learned from a review of the literature. *Urol Ann.* 2015;7(2):141-8. doi: 10.4103/0974-7796.152927
- Liu Y, Chen Y, Liao B, et al. Epidemiology of urolithiasis in Asia. *Asian J Urol.* 2018;5(4):205-214. doi: 10.1016/j.ajur.2018.08.007
- Skolarikos A, Straub M, Knoll T, Sarica K, Seitz C, Petřík A, et al. Metabolic Evaluation and Recurrence Prevention for Urinary Stone Patients :EAU Guidelines .*EurUrol.* 2015;67(4): 750-763. doi: 10.1016/j.eururo.2014.10.029
- Rizvi, S.A.H., Hussain, M., Askari, S.H., Hashmi, A., Lal, M. and Zafar, M.N. Surgical outcomes of percutaneous nephrolithotomy in 3402 patients and results of stone analysis in 1559 patients. *BJU Int.* 2017;120:702-709. doi:10.1111/bju.13848
- Poudyal S, Pradhan M, Chapagain S, Luitel BR, Chalise PR. Comparison of Guy's Stone Score and S.T.O.N.E Nephrolithometry Score for Predicting Outcome of Percutaneous Nephrolithotomy. *J Urol Res.* 2017;4(2):1082.
- Tailly TO, Okhunov Z, Nadeau BR. Multicenter external validation and comparison of stone scoring systems in predicting outcomes after percutaneous nephrolithotomy. *J Endourol.* 2016;30:594-601. doi: 10.1089/end.2015.0700
- Jiang, K., Sun, F., Zhu, J. et al. Evaluation of three stone-scoring systems for predicting SFR and complications after percutaneous nephrolithotomy: a systematic review and meta-analysis. *BMC Urol* 2019 19, 57 doi:10.1186/s12894-019-0488-y
- Mak DK, Smith Y, Buchholz N, El-Husseiny T. What is better in percutaneous nephrolithotomy - Prone or supine? A systematic review. *Arab J Urol.* 2016;14:101-107. doi:10.1016/j.aju.2016.01.005
- Jeong CW, Jung JW, Cha WH, Lee BK, Lee S, Jeong SJ, et al. Seoul National University Renal Stone Complexity Score for Predicting Stone-Free Rate after Percutaneous Nephrolithotomy. *PLoS One.* 2013;18(6):e0065888. doi: 10.1371/journal.pone.0065888
- Vicentini FC, Marchini GS, Mazzucchi E, Claro JF, Srougi M. Utility of the Guy's stone score based on computed tomographic scan findings for predicting percutaneous nephrolithotomy outcomes. *Urology* 2014;83:1248-1253. doi:10.1016/j.urology.2013.12.041
- Choi SW, Bae WJ, Ha US, Hong SH, Lee JY, Kim SW, et al. Prediction of stone-free status and complication rates after tubeless percutaneous nephrolithotomy: a comparative and retrospective study using three stone-scoring systems and preoperative parameters. *World J Urol* 2017 35(3):449-457. doi: 10.1007/s00345-016-1891-6
- Omer Rashid A, Khalid H, Friad G, Hamed R, Y, Buchholz N: Guy's Stone Score as a Predictor for Stone-Free Rate and Complications in Percutaneous Nephrolithotomy: A Single-Center Report from a Stone Belt Country. *Urol Int* 2020;104:746-751. doi: 10.1159/000506211
- M. Khalil, H. Sherif, A Mohey R. Omar. Utility of the Guy's Stone Score in predicting different aspects of percutaneous nephrolithotomy. *Afr J U*, 2018,24,3,,191-196. doi:10.1016/j.afju.2018.06.005
- Okhunov Z, Friedlander JI, George AK, Duty BD, Moreira DM, Srinivasan AK, et al. S.T.O.N.E. nephrolithometry: novel surgical classification system for kidney calculi. *Urology.* 2013; 81:1154-1159. doi: 10.1016/j.urology.2012.10.083
- Thomas K, Smith NC, Hegarty N, Glass JM. The Guy's stone score--grading the complexity of percutaneous nephrolithotomy procedures. *Urology* 2011;78:277-281. doi: 10.1016/j.urology.2010.12.026
- Mandal S, Goel A, Kathpalia R, Sankhwar S, Singh V, Sinha RJ, et al. Prospective evaluation of complications using the modified Clavien grading system and of stone free rates of percutaneous nephrolithotomy using Guy's Stone Score: A single-center experience. *Indian J Urol* 2012; 28:392-8. doi: 10.4103/0970-1591.105749.
- Kumsar S, Aydemir H, Halis F, Köse O, Gökçe A, Adsan O. Value of preoperative stone scoring systems in predicting the results of percutaneous nephrolithotomy. *Cent European J Urol.* 2015;68: 353-357. doi: 10.5173/cej.2015.552
- Labadie K, Okhunov Z, Akhavein A, Moreira DM, Moreno-Palacios J, del Junco M, et al. Evaluation and comparison of urolithiasis scoring systems in percutaneous kidney stone surgery. *J Urol.* 2015;193:154-159.
- Gupta R, Mahajan A. Outcomes of percutaneous nephrolithotomy in elderly versus young patients under regional anesthesia: A comparative study. *Urol Ann.* 2020;12(3):254-258. doi: 10.4103/UA.UA_69_19
- Abdrabuh, A.M. Impact of weight on stone-free rate during percutaneous nephrolithotomy. *Afr J Urol,* 2020;26,65. Doi: 10.1186/s12301-020-00071-4
- Immad MI, Jabbour T Zgheib J. Prone PCNL: Safety and efficacy in obese and morbidly obese patients. *Urol Nephrol Open Access J.* 2018;6(3):84-88. doi: 10.15406/unoaj.2018.06.00209
- Ingimarsson JP, Dagrosa LM, Hyams ES. External validation of a preoperative renal stone grading system: reproducibility and inter-rater concordance of the Guy's stone score using preoperative computed tomography and rigorous postoperative stone-free criteria. *Urology.* 2014;83(1):4549. doi: 10.1016/j.urology.2013.09.008

Authors' Contribution:

| | |
|---------------------------|---|
| Dr. M. Waqas Iqbal | Study design, manuscript writing. Revised and approved the articles. |
| Dr. Farhan Jamshed | Study design, acquisition of data and manuscript writing. Revised and approved the articles |
| Dr. Adeen Akram | Study design, acquisition of data, data analysis and interpretation and write up of results |
| Dr. Nauman Khalid | data analysis and interpretation and write up of results approve the final version of article. |
| Dr. M. Saifullah | Study design. Revising manuscript critically for important intellectual content and approve the manuscript |
| Dr. M. Akram Malik | Study design, critically Revising manuscript critically for important intellectual content and approve the manuscript |
| | All authors are equally accountable for research work and integrity |

Date of submission: 10-04-2021,
 Revised received: 16-12-2021,
 Accepted: 22-01-2022