

Breeding possibilities of Booroola Merino in East-Europe

L. VERESS¹, F. HOREK², T. KOMLOSI¹, A. JAVOR¹ and K. MAGYAR¹

¹University of Agriculture, Debrecen, Hungary

²University of Agriculture, Brno, Czechoslovakia

Abstract. Booroola Merino (BM) rams and ewes were imported from New Zealand to Hungary and Czechoslovakia in the 1980's. Part of the imported animals were proved to be homozygous for the F gene. Frozen semen of these has been used for spreading this gene to Hungarian (HM) and Czechish (CM) Merinos. The crossbred lambs were 4—7 % lighter at birth and at 30 d., 6—24 % lighter at 100 d. and 5—10 % lighter at 1-yr than HM. The 1-yr weight was 10—22 % lighter than that of CM. Greasy fleece weights were 5—10 % higher, staple lengths 7—11 % longer than in HM or CM, while fibres were finer. The litter size of BM × HM ewes at 1st lambing was 1.42, at 2nd 1.52 and at 3rd 1.69, while the last figure for HM ewes was 1.15. The % of twins born to the BM × HM ewes was ca. 35 %, when HM ewes had 12 %. The corresponding figures for triplets were 9 and 0.2 %. The BM × HM crosses had shorter anestrus periods after lambing than HM ewes, except after the first lambing. The BM × HM ewes produced 2.4 lambs/yr. in a continuous lambing system, while HM ewes gave 1.9 lambs.

Index words: Booroola, Merino, Hungarian Merino, Czechish Merino, crossbreeding, live weight, fleece weight, litter size, lambing interval

Introduction

By breeding strains viz. Elektoral in Germany, Negretti in Bohemia and in Hungary, and Mazajevi in Russia, these countries played a significant role in Merino (M) breeding in the third part of the 19th century in Europe. Nowadays, most Ms in the East-European countries are bred in Germany and the G.D.R., Hungary and in the Soviet Union, where primarily meat, meat + fleece and

fleece + meat producing strains, respectively, are kept, while in other countries such as Czechoslovakia, Poland, Rumania and Bulgaria fleece + meat producing strains are dominant. In Hungary, Rumania and Bulgaria some of the Ms are milked as well. Since 1968, accelerated lambing has shown a rapid increase in Hungary (14).

For the past decades an increase in mature body weight has become general in almost every strain of M. Although the direct and com-

binative crossing for production of M has yielded promising results (5, 9), it has failed to become prevalent in large-scale sheep farming.

In England, for the last decades, pure breeding has been reduced to the indispensable reproductional ratio, giving way to a wide range of combinative productional crossing. In comparison with the pure breeding of meat producing strains, this sort of crossing has proved to be far more productive and economical (16).

Booroola Merino (BM) is likely to prove an excellent partner for crossing in those regions of East Europe that are suitable for breeding M (13, 15). Therefore we wish to highlight here some of our plans pertaining to the utilization of this breed.

Subject and method

In response to our recommendation, 20 rams and 3 ewes were brought from Haldon Station, New-Zealand (N.Z.), in the years of 1980, 1982 and 1986, to enlarge the available experimental stock. In 1985, Czechoslovakia bought 7 rams from the same source. This stock was the offspring of 19 rams altogether which descended from 9 original rams imported from Robertson's stock in Australia. While still in N.Z., four of them were qualified as homozygotes on the F gene, five other sires were either not assessed or were labelled as heterozygotes. Our intention is to promote F-gene homozygote populations in several breeding stocks, which could then subsequently be developed into nucleus.

Besides focusing on producing F-gene homozygous stocks, we endeavour to improve maternal traits (i.e. oestrus aseasonality, perfect conceiving and lamb rearing capability) as well as the parameters of fleece producing capacity (i.e. clean-fleece yield, ratio of fine and long staples).

For the purpose of achieving this, the sperm of the best imported rams has been frozen so that »post mortem» fertilization could be carried out. The sperm of heterozygous rams is

going to be used only when a sufficient number of homozygous ewes are available so that homozygous lines could be obtained from them, as it is done in N.Z. and France as well (4, 8). Mating heterozygous partners is also preferable because, on the one hand, 1/4 of the offspring is homozygous, and their phenotype is closer to that of Hungarian M (HM), on the other hand, the gene frequency of the breed is extended as well. The deep-frozen sperm is introduced into the uterine horn by means of an endoscope. The fertility rate, however, has not exceeded the 30—40 percent level and it is still far from the results of 60 % reported by SALAMON et al. (11).

Parallel with new-breed crossing three-breed crossing has also been initiated where partner A is HM on Czech M and partner B is a BM, and the terminal partner is a Suffolk which, owing to its high growth capacity, compensates for the lower growth rate of the BM, and moreover it contributes to improving meat ratio and feed conversion.

There is no doubt, that ewes obtained from this crossing are smaller in weight, but this is not considered to be a shortcoming. Their phenotype, however, is the same as that of HM, their relative fleece production is more favourable. Moreover, this sort of fleece can be marketed at a 20 % higher price as it is finer, longer stapled and cleaner as well (15).

Achieved results

The body weight of the imported rams alternated between 50—70 kg. Due to indoor breeding it has increased by 30 %, the fibre diameter has become 3 microns coarser, staple length and rendement have remained constant while clean fleece yield has increased by 5 % (7).

The lambs of imported BM rams weigh 4—7 % less, at the age of 30 days they lag behind by 5—7 % and when they were 100 days old they weighed 6—24 % less than the HM (15). When they were one year old, their weight decreased by 5—10 % in Hungary and by 10—22 % in Czechoslovakia in compari-

son with pure bred control animals. Grease fleece yield decreased by 1—18 % in Hungary and by 10—22 % in Czechoslovakia.

Grease fleece production in relation to the control group increased by 5—10 % together with the increase of staple length (7—11 %), whilst fur became finer and the rendement improved (6, 14). The influence of BM crossings on decreasing growth and mature body weight has been pointed out by several authors (4). It has also been reported that this effect is of the same degree in both F-gene-bearing homozygous and heterozygous rams (11).

This is true in respect of slaughtering and fleece production characteristics as well (6).

We are firmly convinced that the variance in mature body weight is partly of phenotypic origin which is caused, on one hand, by prenatal and postnatal state of nourishment and, on the other hand, by the age when the animals are used for breeding. Our observations indicate that the BM can be classified as

a slowly developing and growing type. Despite this fact a bigger genotype could be produced, if it were so desired.

Sperms from 15 homozygous and 4 presumably heterozygous rams originating from 6 lines have been frozen so far. We have to store sperm of 10 rams from two further lines.

Fecundity results seem to be satisfactory, though, 3 of the rams that could be appraised so far are merely heterozygous carriers of the F gene. The first lambings happened when the ewes were on the average of 21—22 months old (15), but in spite of this the rate of twinings amounted to 30—37 %, the frequency of litters with 3 lambs alternated between 3 and 13 %. In the course of the second and third lambings this rate grew by an additional 10 %. In comparison with AM ewes with repeated lambings, the rate of productivity was 54 % better.

Ewes are lambing continually on the farm from which these data are obtained. Thus it

Table 1. Progeny test of 4 rams for their daughters' prolificacy.

Rams	lambing n	lambing rate %	twins		triplets and more	
			n	%	n	%
f i r s t						
502/78	162	145,0	61	37,6	6	3,0
62/80	119	140,0	39	32,8	6	5,0
51/80	222	138,0	74	33,3	6	2,7
183/80	15	206,6	5	30,0	2	13,3
Σx, or \bar{x}	518	142,0	179	34,5	20	3,86
s e c o n d						
502/78	147	158,0	57	38,7	13	8,0
62/80	95	168,0	38	40,0	13	13,8
51/80	191	153,0	57	29,8	22	11,4
183/80	—	—	—	—	—	—
Σx, or \bar{x}	433	152,0	152	35,1	48	11,0
t h i r d						
502/78	71	163,0	30	42,2	7	9,4
62/80	51	182,0	16	31,1	13	25,5
51/80	111	166,0	38	34,2	17	16,2
183/80	—	—	—	—	—	—
Σx, or \bar{x}	233	169,0	84	36,0	37	15,8
Hungarian Merino (control)	451	114,5	56	12,4	1	0,2




Fig. 1. The distribution of lambings per month of Hungarian Merinos and Booroolas on one farm in 1987.

has become possible to compare the abilities to accelerated lambing of HM and BM crossings (Table 1). The anoestrus period of the BM is a bit shorter than that of the HM as it is referred to in previous studies (1, 2). Significant difference was found between the average interval of the second and third lambing of BM crossings as far as the main season is concerned ($P < 1\%$), and in the pre-and post-seasons as well as out-of-season ($P < 0,1\%$). The re-lambing interval of M used for control was naturally more favourable, since there was a disparity of 54 % in the rate of reproduction, that made the recovery period necessary for the re-cycling of the F-gene heterozygote carrier ewes longer. Nevertheless, considering the average interval between the 2nd and 3rd lambings a significant difference was found to be advantageous for BM crossings ($P < 0,1\%$).

The distribution of the monthly lambings of M and BM crossings on the same farm in 1987 is shown in Figure 1.

The distribution is were even regarding lambings of M in periods of a quarter of a year. While the average lambings of the M was 1.9 for 1987, in the case of the BM crossings this value came to 2.4.

Currently the most difficult long-lasting technological task is to increase the vitality of the born lambs and to keep them alive.

DYRMUNDSSON and ADALSTEINSSON (3) pointed out that concerning Icelandic sheep there is a definite relationship between the pigmentation of the fur and fecundity capabilities. Pigment spots on the face and on the end of the legs were found more frequent in the case of the Hungarian daughters of BM rams than in that of Hungarian M.

Table 2. The relambing interval of Hungarian Merino and heterozygote (F+) Booroola crossings depending on the conceiving time.

Conceiving time		Main season (1.9—30.11)		Pre-post-season (1.12—31.1) (1.7—31.8)		Out of season (1.2—30.6)		Total	
		HM (control)	BM (F+)	HM (control)	BM (F+)	HM (control)	BM (F+)	HM (control)	BM (F+)
first- second	n	26	53	10	180	8	39	44	272
	\bar{x}	351	323	294	353	324	404	333	354
	s	111	86	111	93	119	92	115	91
	CV%	31.5	26.9	37.6	26.4	36.7	22.8	34.4	38.7
second- third	n	6	32	119	88	155	12	280	132
	\bar{x}	363	292	331	302	293	285	310	298
	s	136	45	94	41	97	105	99	50
	CV%	37.4	15.3	28.5	13.5	33.2	36.9	31.9	59.3
total	n	32	85	129	268	163	51	324	404
	\bar{x}	353	311	328	333	294	376	313	336
	s	113	75	95	85	98	107	101	80
	CV%	31.1	24.1	34.6	25.4	30.1	28.5	30.9	41.8

Conclusions

Specific yields can be increased significantly in East-European territories suitable for Merino (M) keeping by means of crossing first with Booroola (BM) rams and with Suffolk used as terminal partners in comparison with fleece, meat and meat-fleece producing strains of the M. On a Hungarian farm with 3,000 ewes, where continual lambing goes on throughout the year, ewes crossed with BM obtained a productivity rate of 142% for the first lambing, 152% for the second and 169% for the third one. M ewes lambing more than

three times produced a productivity rate of 114,5%.

BM crossed ewes pregnant for the first time conceived again later than Hungarian, but their following interval of re-lambing on the average was more favourable than that of Hungarian M. This deviation is sustained statistically ($P < 0.1\%$). As regards fecundity data, BM crossings are inevitably better; in 1987, against the background of 2,000 M ewes, the annual value of annual increase came to 1.9, while for 1000 crossed ewes this indicator was found 2.4 lambs.

References

1. BINDON B.M., PIPER, L.R. & EVANS, R., 1982. Reproductive Biology of the Booroola Merino. Proc. Worksh. at Armidale NSW. Aug. Div. of. Anim. Prod., CSIRO, Austr.
2. BINDON, B.M., 1984. Reproductive biology of the Booroola Merino Sheep. Austr. J. Biol. Sci. 37. 163—189. p.
3. DYRMUNDSSON, O.R. & ADALSTEINSSON, S. 1980. Coat colour supresses sexuelle activity in Icelandic sheep. J. Heredity 71: 363.
4. ELSÉN, J.M. & ORTAVANT, R., 1984. Le gene Booroola mise en evidence fonctionnement perspectives d'utilisation. 9-ième Journ. Rech. Ovine et Capr. (Paris) INRA-ITOVIC: 415—451.
5. GÖHLER, H., 1978. Untersuchungen zur Bestimmung der Körperentwicklung und des Schlahtkörperwertes bei Lämmern. Diss. B. K.-M. Univ. Leipzig.
6. HINCH, G.N., KELLY, R.W., DAVIS, G.H. OWENS, J.L. & CROSBIE, S.T. 1985. Factors affecting lambs birth weights from high fecundity Booroola ewes. Anim. Repr. Sci. Amsterdam. 8.1/2 53—60. p.
7. HORAK, F., 1987. Ergebnisse aus der Zucht der Booroola Schafe in der Tschechoslowach. Sozialist. Republik. Wiss. Tag. Humboldt Univ. 3—4. Dez. Berlin.
8. KLEEMANN, D.O., PONZONI, R.W., STAFFORD, J.E., CUTTEN, I.N. & CRIMSON, R.Y., 1985. Growth and

- carcass characters of the South Australian Merino and its Crosses with the Booroola and Tharghee fertility Merino. *Austr. J. Exp. Agric.*, Melbourne. 25. 4: 750—757. p.
9. KELLY, R.W., DAVIS, G.H. & ALLISON, A.J., 1982. Productive changes in longwool breeds in New Zealand following crossbreeding with Booroola type rams. *Proc. Austr. Soc. Anim. prod. CSIRO. Armidale* 13. 413—416.
 10. KÖNIG, K.H., 1978. Möglichkeiten zur Verbesserung der Fortpflanzungsleistung weiblicher Schafe durch Massnahmen der Züchtung und durch gesteuerte Tageslichtlängen. *Diss. B. K.-M. Univ. Leipzig*.
 11. PONZONI, R.W., FLEET, N.R., WALKLEY, I.R.W. & WALKER, R.K., 1985. A note on the effect of the F gene on wool production and live weight of Booroola × South Australian Merino rams. *Anim. Prod.* 40: 367—369. p.
 12. SALAMON, St., MAXWELL, W.M.C. & EVANS, G., 1985. Fertility of ram semen frozen-stored for 16 years. *Proc. 7.th. Annual Conf. Austr. Repr. Biology. Adelaide* 62. p.
 13. VERESS, L., 1983. Adatok a booroola merino tenyésztési programjához.
 14. VERESS, L., VÉGH, J., TURAI, I., TARNÓCZI, T. & ECSEDY, J., 1987. Some conclusions concerning the large-scale accelerated lambing of Hungarian Merino ewes. *Acta Agr. Hung. Budapest.* 37./1—2/ 111—121. p.
 15. VERESS, L. & ALLISON, A.J. 1988. Improvement of Merino sheep for prolificacy. *Acta Agr. Hung. Budapest.* 37./1—2/ 173—182. p.
 16. WASSMUTH, R., 1971. Optimale Verwendung des genetischen Materials unter spezieller Berücksichtigung der Kreuzung im Vergleich zu anderen Zuchtverfahren. *Schafe und Ziegen. X. Intern. Tierzucht Kongr. Paris, Versailles: 97—115. p.*