

Finnsheep and their utilization in crosses with the Merino under range conditions of South-Africa

J. C. GREEFF and J. H. HOFMEYR

Animal and Dairy Science Research Institute, Irene, RSA

Abstract. Two field trials were performed at two localities in a semiarid region to evaluate the performance of Finn (F) × Merino (M) crossbred females. The trials involved 781 F × M ewes and 657 M ewes. Ile de France, S.A. Mutton Merino, Dohne Merino, M, and F × M rams were used for matings. The F × M ewes were superior to M ewes in lambing-%, especially after mating as lambs, and the fecundities were 187 and 107 %, resp. Multiple lambs of F × M ewes had better survival rates than those of M ewes. Lambs born from F × M ewes grew faster, had a larger weaning mass, over 10 % longer staples, ca. 2 mm thicker fibres, 30 % lower clean fleece weights, and a little less crimps/25 mm than those born from M ewes. Thus, F × M seemed to provide a suitable and acceptable composite showing remarkable adaptability to harsh environmental conditions. The results from the F × M wool and its special qualities can be regarded as useful attributes. Establishment of a composite F × M female line appeared well-founded.

Index words: Finnsheep, S.A. Mutton Merino, crossbreeding, mutton production, wool production, reproduction, daily gain, synthetic

Introduction

Improvement in the reproductive rate through accelerated lambing and/or higher fecundity rate of the ewe flock offers the greatest single opportunity for increasing the biological efficiency of lamb as well as mutton and wool production. The infusion of high fertility genes through crossbreeding has proved to be the most rapid breeding procedure to bring about such biological efficiency and for this reason Finnsheep (F) has been

introduced into the sheep flocks of many countries. Numerous trials have been conducted in recent years to evaluate its suitability in crossbreeding with local breeds and different environmental conditions.

F were imported into the RSA in 1968 and have been used in several crossbreeding experiments (3, 4). A small nucleus experimental flock is also maintained to observe the performance of pure-breds under local conditions.

This short paper covers the preliminary

results of two field trials set up at two localities in a semi-arid region to evaluate the performance of F × Merino (M) crossbred females. As is so often the case it is not always possible to plan and conduct field trials as well as in formal experiments, but in spite of certain limitations useful results do accrue. M controls were included in both trials.

Both trials were launched during an extended drought period which lasted from the previous winter season till last year (1987 July), and the performance of the animals, particularly in the first trial initiated in 1983, suffered seriously as a result of poor nutritional conditions.

Animals in the second trial were supported by irrigated pastures and were not notably affected by the drought conditions.

Experimental

First trial. This trial involved 600 F × M crossbred ewes and 600 M ewes, subdivided into groups and inseminated with semen from Ile de France (IdF), South African Mutton Merino (SAMM) and Dohne Merino (DM) rams. This was followed by group mating of the M and F × M ewes with M and F × M rams. Owing to deteriorating environmental conditions, these latter groups were treated slightly better during the last three weeks of gestation.

Second trial. In the second trial 181 M × M and 57 M ewes were involved in a comparative performance evaluation.

Results

Trial 1

Reproduction: Owing to environmental conditions, lambing percentages for natural mating following AI were poor in both groups but clearly in favour of the F × M with 61 % vs. M 48 %. Fecundity was 118 % for F × M and 102 % for M.

Survival rate of lambs: No significant

differences of survival rate occurred between groups in the case of single born lambs, but the survival rate from birth to weaning was significantly ($P < 0,05$) better for lambs born as multiples to F × M ewes (Table 1).

Growth: Lambs from F × M ewes had significantly ($P < 0,01$) higher growth rates from birth to five weeks of age and significantly ($P < 0,01$) higher weaning masses than lambs born from M ewes at 100 days of age (Table 2). The differences between sire-breed groups were not significant except for the Dohne Merino.

Wool at 9 months of age: Except for the Mutton Merino × M, Dohne Merino × M and Ile de France × M, although not significant, had on average slightly lighter fleeces than their related crosses born from F × M ewes (Table 3). As expected the M and M offspring produced the finest fibres while F × M offspring produced longer wool staples.

Trial 2

Reproduction: Of the 181 F × M ewes exposed to rams (F × M) between 4–10 months of age, 65 % lambled before 12 months of age with a fecundity rate of 136 % (Table 4). None of the M ewes lambled. F × M ewes lambled within a period of 38 days.

All ewes (both groups) were again exposed to rams (F × M) at 18 months of age after some culling on wool faults had been applied (Table 5).

Significant differences were found between: lambing % (92 vs. 75 % in favour of F × M); and fecundity (187 vs. 107 % also in favour of F × M).

Survival rate of lambs: The first group of offspring of the F × M ewes had a survival rate of 77 % (Table 4) at weaning (100-days) and the second group, 98,9 % vs. the M's 80,6 % (Table 5).

Growth: Lambs of the F × M ewes were significantly ($P < 0,01$) heavier at weaning (100-days) than lambs born from M ewes (19,2 vs. 14,9 kg).

Table 1. Survival rate (%) of lambs at different stages from birth to weaning (Trial 1).

	Ewe breed					
	Finn × Merino Type of birth*			Merino Type of birth*		
	1	2	3	1	2	3
Number of lambs born	349	220	39	424	134	21
Survival rate (%)						
at birth	99,1	98,6	100	99,1	98,5	95,2
birth — 5 weeks	89,9	82,8	66,0	90,7	50,0	55,0
5—10 weeks	92,7	88,2	76,9	91,8	85,1	63,6
10 weeks — weaning (100 days)	94,8	97,3	80,0	95,4	92,5	100
Total survival rate	79,4	65,9	40,0	78,3	55,2	33,3

* 1 = Singleton; 2 = Twins; 3 = Triplets

Table 2. Average daily gain (ADG) from birth to 5 weeks of age and weaning mass of crossbred lambs (g/day)² (Trial 1).

Ram breed	Ewe breed			
	n	Finn × Merino	n	Merino
ADG (birth — 5 weeks of age: g)		(x ± S.E.)		(x ± S.E.)
Ile de France	147	299 ± 9,3 ^a	171	277 ± 2,4 ^a
Dohne Merino	24	269 ± 38,6 ^b	95	266 ± 21,5 ^b
SA Mutton Merino	108	290 ± 6,2 ^a	89	276 ± 8,6 ^a
Merino ¹	83	295 ± 16,5 ^a	37	274 ± 16,3 ^a
Finn × Merino ¹	98	292 ± 3,1 ^a	44	274 ± 6,7 ^a
Weaning mass at 100 days (kg)				
Ile de France	98	18,8 ± 0,59 ^a	127	16,7 ± 0,13 ^a
Dohne Merino	9	16,2 ± 3,79 ^b	66	16,1 ± 1,34 ^a
SA Mutton Merino	79	18,5 ± 0,35 ^a	54	16,7 ± 0,59 ^a
Merino ¹	61	18,8 ± 0,95 ^a	19	14,3 ± 1,35 ^b
Finn × Merino ¹	71	18,2 ± 0,19 ^a	30	15,1 ± 0,42 ^b

¹ Lambs born from Merino and Finn × Merino rams received slightly different treatments. See text.

² Means with different superscripts in the same column differ significantly from each other.

Table 3. Wool characteristics of crossbred lambs shorn at 9 months of age in 1987.¹ (Trial 1).

Breed/Genotype ²	n	Clean fleece (kg)	Staple length (mm)	Fibre diameter (micron)	Crimps per 25 mm	Crimp deviation ³
<i>Lambs</i>						
Merino	21	1,40 ^b	88,3 ^a	15,7 ^a	13,3 ^{ac}	79 ^a
F/M × M	25	1,24 ^a	100,5 ^d	16,4 ^a	13,6 ^{ac}	83 ^{ab}
M × F/M	50	1,45 ^b	101,8 ^d	17,2 ^b	12,5 ^a	84 ^{ab}
F/M × F/M	37	1,24 ^a	106,9 ^a	17,7 ^b	12,8 ^a	87 ^b
IdF × M	111	1,38 ^b	91,1 ^c	18,2 ^b	16,0 ^d	99 ^d
IdF × F/M	125	1,42 ^b	96,9 ^f	19,9 ^c	14,4 ^c	103 ^c
SAMM × F/M	128	1,26 ^a	93,4 ^c	18,2 ^b	14,2 ^{bc}	94 ^c
SAMM × M	61	1,31 ^a	80,5 ^b	16,8 ^{ab}	14,3 ^{bc}	87 ^b
DM × F/M	16	1,38 ^b	103,3 ^d	18,2 ^b	13,0 ^{ab}	91 ^b
DM × M	59	1,33 ^{ab}	80,2 ^b	16,4 ^a	13,2 ^{ab}	82 ^a

¹ Means with the same superscript in the same column do not differ significantly from each other.

² Ram breed mentioned first.

³ Duerden standard = 100.

Wool: Ewes were shorn at 11 months of age (Table 6). As in Trial 1 the M produced significantly ($P < 0,01$) more clean wool than the F × M crossbreds, the latter producing an amount equivalent to 71 % of the M.

Conclusion/Remarks

In a country such as South Africa, where both lamb and wool production are important, environmental circumstances dictate the relative emphasis placed on the two commodities.

In general, extensive range conditions favour wool production and here the M is likely to remain the dominant breed in production systems. This breed constitutes approximately 65 % of the current sheep population of 35 million (1). The M is, however, not suitable for intensive lamb and wool production systems (2, 4) and with growing emphasis on more intensive production systems, particularly in areas of higher potential it is obvious that a more suitable genotype is required.

Results obtained from a comprehensive crossbreeding experiment and several field trials, including the two reported on in this paper, have led to the conclusion that the halfbred F × M provides a suitable and acceptable composite showing remarkable adaptability to even comparatively harsh environmental conditions.

The favourable results obtained from the crossbred (F × M) wool and its special qualities (low felting and low creasing features) are regarded as useful attributes.

There appears to be sufficient evidence available now to support the establishment of a composite F × M female line for deployment in the industry, certainly under conditions where favourable nutritional conditions exist.

Table 4. Reproduction of Finn × Merino and Merino ewes exposed to rams from 4–10 months of age (Trial 2).

	Ewe breed	
	Finn × Merino	Merino
Number of ewes available	181	57
Lambing (%)	65,7	—
Fecundity (%)	136,1	—
Survival rate of weaning ¹ (%)	77,1	—
Mass at conception (kg)	29,4–30,9	(27,8–28,6) ²

¹ Lambs weaned per total number of lambs born.

² Mass range concurrent with Finn × Merino conception.

Table 5. Reproduction of F × M and M ewes mated at 18 months of age and survival rates of their lambs (Trial 2).

	Ewe breed	
	Finn × Merino	Merino
Number of ewes available	157	37
Ewe mass at mating kg ± S.E.	47,8 ± 0,33	43,2 ± 0,68
Lambing (%)	92,4	75,5
Fecundity (%)	186,9	107,1
Survival rate at birth ¹ (%)	99,3	86,6
Survival rate from birth to weaning ² (%)	98,9	80,6

¹ Lambs born alive per total number of lambs born.

² Lambs weaned per number of lambs born alive.

Table 6. Wool characteristics of the F × M and M (Trial 2) measured at 11 months of age.

	Genotype	
	Finn × Merino	Merino
Number of ewes shorn (n)	168	55
Mass at shearing (kg)	38,7	33,3
Greasy fleece mass (kg)	3,4	4,8
Clean yield (%)	69,6	71,7
Clean fleece mass (kg)	2,4	3,4
Staple length (mm)	114,7	106,8
Crimps per 25 mm	8,9	9,0
Fibre diameter (micron)	21,3	19,9
Crimp deviation	92	86

References

1. ABSTRACT OF AGRICULTURAL STATISTICS, 1988. Directorate of Agricultural Economics and Marketing. Pretoria. ISBN 0621114235.
2. BONSMAN, F.N., 1939. Factors influencing the growth and development of lambs with special reference to crossbreeding of Merino sheep for fat lamb production in South Africa. Univ. Pretoria Publ., Sec. 1., NO. 48, Rep. South Africa.
3. HOFMEYR, J.H., UECKERMANN, L., KÜHN, G.P. & ROUX, C.Z., 1974. Intensive sheep production: Preliminary observations on the effect of crossbreeding as exhibited by the production qualities of certain Merino crosses. Proc. of 5th Congress. S. Afr. Genet. Soc., Stellenbosch Univ. February, 1974. 140—144.
4. HOFMEYR, J.H. 1980. Implications of experimental results of crossbreeding sheep in the Republic of South Africa. Proc. World Congr. Sheep and Beef Cattle Breed. Vol I. Ed. R A Barton and W C Smith. The Dunmore Press, New Zealand.