

# Inclusion of wheat bran in barley-soybean meal diets with different phosphorus levels for growing-finishing pigs

## I. Effects on nutrient digestibility and mineral balances in finishing pigs

EIJA HELANDER, MATTI NÄSI and KIRSI PARTANEN

HELANDER, E., NÄSI, M. & PARTANEN, K. 1994. **Inclusion of wheat bran in barley-soybean meal diets with different phosphorus levels for growing-finishing pigs. I. Effects on nutrient digestibility and mineral balances in finishing pigs.** *Agricultural Science in Finland* 3: 27-39. (Department of Animal Science, P.O. Box 28, FIN-00014 University of Helsinki, Finland. Present address: Eija Helander, Cultor Ltd, P.O. Box 105, FIN-00241 Helsinki, Finland.)

The effect of the intrinsic phytase of wheat bran (WB) on phytin-phosphorus availability to pigs on barley-soybean meal diets was measured in a digestibility and balance experiment using a 2x3 factorial arrangement in a 6x5 cyclic change-over design. The factors were WB inclusion (0 or 100 g per kg, later referred to as WB- and WB+, respectively) and three phosphorus (P) levels: high (HP), medium (MP) and low (LP). The inclusion of WB in the diet did not significantly improve dietary P utilization. However, the absorption and retention of P appeared to be slightly improved by WB inclusion in the LP diet. This improvement may be due to WB phytase. The effects of WB on the digestibility and balance of other minerals remained relatively small. The P level, on the other hand, had a greater effect on mineral balances. Ash digestibility was not affected by the treatments. Dry matter and organic matter digestibilities were impaired when WB was included in the diet. N absorption of intake was higher ( $p < 0.01$ ) on WB+ diets. The retention of N appeared to decrease on HP and MP diets due to WB, but on LP diets no decrease was observed. Regarding the polluting P-emission, the LPWB+ diet appeared to be the most favourable, since the faecal excretion of P was the lowest and the overall retention coefficient of P was the highest. On the basis of these results, an addition of 100 g WB to the diet in order to improve P utilization does not seem very attractive.

---

Key words: phytin, phytase, availability

---

### Introduction

More than half of the phosphorus (P) in feedstuffs of plant origin occurs in the form of phytate, which is poorly available to non-ruminants (CROMWELL 1992). Phytic acid readily forms complexes with several minerals, like calcium (Ca) and iron (Fe) (HARRIS 1955), zinc (Zn) and manganese (Mn) (ref. ERDMAN 1979) as well as with proteins (SCHEUER-MANN et al. 1988a), thus reducing their availability

to animals. High levels of phytate in the diet can also cause magnesium (Mg) deficiency (POMERANZ 1978). Pig diets are generally supplemented with inorganic P to ensure that the P requirement is satisfied. If the phytate P of plants could be made more available, a reduction in inorganic P supplementation of the feeds would be possible, making P-emissions through manure decrease considerably.

Phytase catalyzes the hydrolysis of phytic acid to

inositol and orthophosphate. The intrinsic phytase activity of different plants varies considerably. Wheat is one of the feedstuffs which has a high phytase activity ranging from 300 to 2000 IU/kg (1 IU is 1  $\mu$ mol P liberated from sodium phytate/min at 37°C) (POINTILLART 1988). Most of its phytase appears in the surface layers (POMERANZ 1978). In an experiment by SCHEUERMANN et al. (1988a), wheat phytase was found to hydrolyze maize phytate to the same extent as wheat phytate. BAGHERI and GUEGUEN (1985) have established that the utilization of P improved when an oats-soybean meal diet was supplemented with 200 g of wheat bran (WB) per kg. WB addition also improved the utilization of Mg but decreased that of Ca and Zn.

The present work was undertaken to evaluate whether the intrinsic phytase of WB could improve the digestibility of phytic P in commercial barley-soybean meal diets enough to serve as a partial or complete replacement for the added inorganic P in the diets of growing-finishing pigs.

## Material and methods

The digestibility and balance assay was conducted with six Large White x Landrace barrows using a 2x3 factorial arrangement in a 6x5 cyclic change-

over design. The factors were WB inclusion (0 or 100 g per kg, later referred to as WB- and WB+, respectively) and three P levels, high (HP), medium (MP) and low (LP) corresponding to 4.33 g (3/3), 2.99 g (2/3) and 1.64 g (1/3) digestible P per feed unit (FU = 0.7 kg starch equivalent), respectively. Thus, the experiment consisted of six different diets: HPWB-, HPWB+, MPWB-, MPWB+, LPWB- and LPWB+. The nutrient digestibility, nitrogen (N) balance and the balance of P, calcium (Ca), magnesium (Mg), potassium (K) and zinc (Zn) were determined. The pigs were kept in metabolism cages made of galvanized iron, which allowed separate quantitative collection of faeces and urine. Each period lasted 10 days: five days of adjustment and five days of collection. The initial weight of the pigs was 70.6 kg (SE 3.57) and the final weight 122.5 kg (SE 5.52).

The feeds were pelleted (65°C, 4 mm diameter) barley-soybean meal complete mixtures. The composition of the experimental diets is shown in Table 1. HP diets have been found adequate for pigs with respect to all nutrients (SALO et al. 1990), and the others with respect to all other nutrients except P. The inorganic source of P in the present experiment was dicalciumphosphate. The energy content was 0.99 FU/kg in WB- diets and 0.96 FU/kg in WB+ diets. The intention was to give the same

Table 1. Composition of the experimental diets, g/kg.

Treatment <sup>1</sup>	HPWB-	HPWB+	MPWB-	MPWB+	LPWB-	LPWB+
Barley	756	668	759	670	764	674
Soybean meal	189	178	188	178	187	178
Wheat bran	-	100	-	100	-	100
Molasses	20	20	20	20	20	20
Dicalciumphosphate	16	15	9	8	1	-
Limestone	7	7	12	12	16	16
Serla Bondex	5	5	5	5	5	5
NaCl	3	3	3	3	3	3
Trace mineral mix <sup>2</sup>	2	2	2	2	2	2
Vitamin mix <sup>3</sup>	1	1	1	1	1	1
Lysine	1	1	1	1	1	1

<sup>1</sup> HP = high P, MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> Supplied per kg diet: 20 mg Fe, 21 mg Mn, 21 mg Cu, 73 mg Zn, 0.2 mg I and 0.1 mg Se.

<sup>3</sup> Supplied per kg diet: 5000 IU vitamin A, 800 IU vitamin D, 60 mg vitamin E, 2 mg vitamin K, 2 mg thiamin, 3 mg riboflavin, 20  $\mu$ g vitamin B<sub>12</sub>, 50  $\mu$ g biotin, 10 mg pantothenic acid and 20 mg niacin.

amount of FUs/d to each pig (ranging from 2.6 FU to 3.0 FU/pig/d in periods 1 to 5, respectively) but, in practice, the pigs on WB-diets got 0.16 FU/d more than the other pigs. Just before feeding, the feed was mixed with water (one litre water per kg feed) and after feeding the pigs were given water *ad libitum*.

The feeds and faeces were analyzed by standard methods (AOAC 1984). Amino acids were assayed with a Beckman 6300 amino acid analyzer. ICP-AES equipment was used in phytic acid determination (PLAAMI and KUMPULAINEN 1991). The sample was first burned by inductively coupled plasma and then the P content of the sample was indirectly measured by atomic emission spectrophotometry. P from feeds, water and faeces was analyzed colorimetrically after dry ashing by the vanadomolybdate procedure of TAYSSKY and SHORR (1953). The other minerals of the feeds, drinking water, faeces and urine were measured with a Perkin-Elmer 5100 PC atomic-absorption spectrophotometer, except K, which was assayed with a Corning 435 flame photometer. Phytase activity was measured as free phosphate from phytate after incubating the sample in a 0.1 M sodium acetate buffer, pH 5.0, at 35°C for 30 min. A phytase unit (U) is defined as the amount of enzyme that liberates 1 µmol of inorganic P from sodium-phytate in one minute. Phytase activity was measured from the main raw materials and from the complete feeds.

The data were subjected to an analysis of variance using the following model (SNEDECOR and COCHRAN 1989):

$$Y_{ijk} = \mu + A_i + P_j + T_k + e_{ijk}$$

where  $\mu$  = overall mean,  $A_i$  = the effect of animal  $i$ ,  $P_j$  = the effect of period  $j$ ,  $T_k$  = the effect of treatment  $k$  and  $e_{ijk}$  = residual error.

In analyzing the crude protein digestibility, N intake was used as a covariate. All other results were first corrected by using FU intake as a covariate, but because it did not have any significant effect on the digestibility and retention of the nutrients, this covariate was omitted. The degrees of freedom for treatment effects were further parti-

tioned into single degrees of freedom by the following orthogonal contrasts: C1 = WB- vs. WB+, C2 = linear effect of P level, C3 = quadratic effect of P level, C4 = interaction C1 x C2, C5 = interaction C1 x C3.

## Results and discussion

### Chemical analyses

The analyzed chemical composition of the experimental diets is presented in Table 2. Although the experimental feeds were produced in the normal production line of a big feed factory, not designed for mixing small feed batches, the chemical composition of the diets was relatively close to the target. Only the protein and amino acid contents of the diets were slightly higher than targeted and, therefore, the HPWB+ and MPWB+ diets contained more protein per FU than the other diets. The hemicellulose content varied from 101 g/kg DM to 130 g/kg DM, naturally being higher on WB+ diets.

Table 3 shows the P, Ca and phytase content of the diets and feed ingredients. The P content of barley was 3.5 g/kg DM and two thirds of it was of phytic origin, which is in accordance with the figures reported earlier by POINTILLART (1988) and JONGBLOED et al. (1991). In WB, 0.89 of total P was of phytic origin. POINTILLART (1988) has reported the corresponding figure to be 0.75 for wheat bran and JONGBLOED et al. (1991) 0.80 for wheat middlings. According to a review by HOUSEMAN and de BRUYNE (1989), phytate P is located in the aleurone and pericarp layers in monocotyledons. Thus, the content of phytate P in wheat bran is highly dependent on the technical process it comes from.

The P contents in HPWB- and HPWB+ diets were 8.2 and 8.0 g/kg DM, in MPWB- and MPWB+ diets 6.1 and 6.2 g/kg DM, and in LPWB- and LPWB+ diets 4.4 and 4.9 g/kg DM, respectively. The phytic P content of the experimental diets varied from 0.305 to 0.621 in total P, being always higher on WB+ diets. The digestible P content of the diets was calculated by using the determined digestibility coefficients (Table 6). Digest-

Table 2. Chemical composition (g/kg DM) and calculated energy content of the experimental diets.

Treatment <sup>1</sup>	HPWB-	HPWB+	MPWB-	MPWB+	LPWB-	LPWB+
Dry matter	897	893	896	891	891	900
Ash	61	64	61	62	55	54
Crude protein	199	199	204	202	201	196
Crude fat	26	28	21	24	24	30
Crude fibre	53	53	49	55	56	58
Nitrogen free extract	661	656	665	657	664	662
Hemicellulose	104	130	101	114	124	128
Cellulose	54	48	51	49	50	52
Lysine	10.2	10.1	11.5	10.5	10.7	11.1
Threonine	7.2	7.1	7.4	7.1	7.0	6.8
FU <sup>2</sup> /kg feed	0.99	0.96	0.99	0.96	0.99	0.96
ME MJ/kg feed	12.4	12.0	12.4	12.0	12.4	12.0

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> FU = 0.7 kg starch equivalent.

Table 3. Phosphorus, calcium and phytase contents of the experimental diets and feed ingredients (g/kg DM)

Treatment <sup>1</sup>	HPWB-	HPWB+	MPWB-	MPWB+	LPWB-	LPWB+	Soybean meal	Barley	Wheat bran
Phosphorus	8.2	8.0	6.1	6.2	4.4	4.9	6.8	3.5	11.3
Phytic acid	9.0	11.0	9.3	10.9	8.8	10.7	14.4	8.2	35.6
P from phytic acid	2.5	3.1	2.6	3.1	2.5	3.0	4.0	2.3	10.0
Phytic P % of total P	0.305	0.388	0.426	0.496	0.560	0.621	0.590	0.657	0.891
Phytase, U/kg	< 600	< 700	< 300	< 400	< 300	< 500	n.d. <sup>2</sup>	300	2800
Digestible P (measured)	4.2	4.2	2.7	2.5	1.4	1.8			
Calcium	9.9	9.7	10.5	10.0	10.4	10.7			
Digestible Ca	3.51	3.22	3.66	3.39	3.48	3.77			
Ca:P ratio	1.21	1.21	1.72	1.62	2.35	2.20			
Ca:dig.P ratio	2.36	2.31	3.89	4.0	7.43	5.94			
Dig.Ca:dig.P ratio	0.84	0.77	1.36	1.36	2.49	2.09			

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> n.d. = not detected.

ible P ranged from 4.2 to 1.4 g/kg DM. On a LPWB+ diet, the measured digestible P content was 0.4 g/kg DM (28%) higher than on a LPWB-diet. This difference cannot be explained by the differences in the total P content of the diets, although the LPWB+ diet contained 11% more P than the LPWB- diet. The difference may rather be attributed to WB phytase.

The phytase activity of WB was 2800 U/kg. POINTILLART (1988) has measured activities between 600 and 1700 IU/kg for wheat bran. The

highest enzyme activities in wheat are found from its surface layers (POMERANZ 1978); thus, the processing technology affects the phytase activity of the end product. The phytase activity of barley was the same as in another experiment by Helander (1993, preliminary results). Soybean meal did not show any phytase activity. COSGROVE and IRVING (1980) have reported phytase activity in soybeans, and BAGHERI and GUEGUEN (1982) have also indicated SBM to have some phytase activity. It is likely that the phytase activity in SBM depends on

the temperature of the process SBM comes from.

There is some inaccuracy in the figures concerning the phytase activity of the diets: e.g., 100 g of WB per kg should have increased the phytase activity of WB+ diets by 280 U/kg. No such increase, however, could be found, which lends support to activity loss during pelleting. Steam pelleting at 80°C reduced the phytase activity in an experiment by JONGBLOED and KEMME (1990), and SCHWARZ and SCHÖNER (1991) reported 15-25 % losses in phytase activity when the temperature of the pellets was 70°C. Since the pelleting temperature in the present trial was lower, this does not explain the smaller-than-expected differences between the treatments. It is probably more a question of an analytical problem: when the phytase activity of the feeds is low, the sensitivity of the analytical method may not be sufficient (Puhakka 1993, personal communication). In any case, the phytase activity was always higher on WB+ diets than on other diets.

### Digestibilities and balances

The average dry matter intake of the pigs was 2526 g (82 g/W<sup>0.75</sup>) on WB- diets and 2435 g (79 g/W<sup>0.75</sup>) on WB+ diets. The difference in daily energy intake was 0.16 FU. The pigs on WB+ diets ate on average 38 g more hemicellulose and 8.6 g more cellulose per day in spite of their lower total

feed intake. WB inclusion was found to impair the dry matter digestibility ( $p < 0.001$ ) (Table 4). There tended to be an interaction between WB inclusion and P level in organic matter digestibility: the digestibility decreased linearly on WB+ diets ( $p < 0.05$ ) when the P level of the diets decreased and tended ( $p < 0.1$ ) to decrease quadratically on WB- diets. No effect on ash digestibility could be found due to treatments. DIGGS et al. (1965) have reported WB to decrease the metabolizable energy content of a diet. Also, other fibre sources are known to decrease digestibility: the faecal digestibility of organic matter was reduced by the inclusion of 50 g cellulose or 50 g of straw meal per kg diet in an experiment conducted by DEN HARTOG et al. (1988).

The amounts of excreted N and digested N were affected by the daily N intake. Correcting the digested N by N intake decreased the digestibility coefficient on all WB- diets and increased it on WB+ diets (Table 5). Since phytic acid readily forms complexes with protein, it was interesting to note that the crude protein digestibility improved significantly ( $p < 0.01$ ) by WB inclusion. This may be due to the intrinsic phytase of WB as in the experiment by MROZ et al. (1991), where a significant improvement in ileal amino acid digestibility by microbial phytase addition was found. Neither N excretion in urine nor N retention was significantly affected by daily N intake; however, N excretion in urine was always lower ( $p < 0.05$ ) on WB+ diets. N retention tended to be lower on HPWB+ and MPWB+ diets than on corresponding diets without

Table 4. Nutrient digestibilities of the experimental diets.

Treatment Nutrient digestibility	HP <sup>1</sup>		MP		LP		SEM	Statistical significance <sup>2</sup>				
	WB-	WB+	WB-	WB+	WB-	WB+		C1	C2	C3	C4	C5
Dry matter	0.830	0.826	0.836	0.816	0.828	0.814	0.0035	***	o	NS	NS	NS
Organic matter	0.850	0.846	0.856	0.837	0.848	0.834	0.0031	***	*	NS	NS	o
Ash	0.506	0.505	0.499	0.488	0.501	0.481	0.0125	NS	NS	NS	NS	NS
Crude protein	0.831	0.865	0.814	0.850	0.837	0.858	0.0051	**	NS	*	o	NS

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> (P < 0.10) o, (P < 0.05) \*, (P < 0.01) \*\*, (P < 0.001) \*\*\*, (P > 0.10) NS

C1 = WB- vs. WB+, C2 = P lin, C3 = P quadr, C4 = C1 × C2, C5 = C1 × C3

Table 5. Nitrogen balance and protein utilization in pigs fed on the experimental diets.

Treatment	HP		MP		LP		SEM	Statistical significance				
	WB-	WB+	WB-	WB+	WB-	WB+		C1	C2	C3	C4	C5
N intake, g/d	80.6	77.7	85.2	78.2	80.7	76.5	0.41	***	NS	***	NS	***
N excretion in faeces, g/d	13.1	11.6	12.8	12.6	12.6	12.5	0.26	*	NS	NS	*	NS
N digested, g/d	67.5	66.1	72.4	65.6	68.0	64.0	0.59	***	NS	***	*	**
-of intake	0.836	0.851	0.850	0.839	0.843	0.836	0.0040	NS	NS	NS	*	NS
-of intake (corrected by N intake)	0.831	0.865	0.814	0.850	0.837	0.858	0.0030	**	NS	*	o	NS
N excretion in urine, g/d	31.9	32.7	37.1	33.1	36.0	31.3	1.26	*	NS	o	*	NS
N retained, g/d	35.6	33.4	35.3	32.4	32.0	32.6	1.16	NS	o	NS	NS	NS
-of intake	0.446	0.432	0.417	0.418	0.400	0.428	0.0142	NS	NS	NS	NS	NS
-of digested	0.535	0.508	0.491	0.498	0.475	0.514	0.0174	NS	NS	NS	o	NS
-g/W <sup>0.75</sup> /d	1.158	1.091	1.154	1.065	1.041	1.088	0.0377	NS	NS	NS	NS	NS
Urea N, g/d	24.8	27.1	28.8	27.2	30.8	27.5	1.22	NS	*	NS	*	NS
Urea N, g/W <sup>0.75</sup> /d	0.769	0.862	0.905	0.857	0.970	0.868	0.0399	NS	*	NS	*	NS
Biological value	0.602	0.577	0.556	0.568	0.545	0.584	0.0167	NS	NS	NS	o	NS

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> (P < 0.10) o, (P < 0.05) \*, (P < 0.01) \*\*, (P < 0.001) \*\*\*, (P > 0.10) NS

C1 = WB- vs. WB+, C2 = P lin, C3 = P quadr, C4 = C1 × C2, C5 = C1 × C3

WB. No decrease in N retention due to WB was found on the LP diet. Some interactions between WB inclusion and P level were found. A decreasing P level had a linear ( $p < 0.05$ ) and a curvilinear ( $p < 0.1$ ) increasing effect on N excretion on WB-diets, while no effect due to P was found on WB+ diets. A decreasing P level also seemed to have a linear decreasing effect on N retention ( $p < 0.05$ ) and on the biological value of protein ( $p < 0.05$ ) on WB-diets. On WB+ diets those parameters remained constant. No effect on N digestibility or N retention was found in an earlier experiment where the inorganic P was omitted from the diet (NÄSI and HELANDER 1993).

The average daily intake of P ranged from 20.5 g/d on a HPWB+ diet to 12.5 g/d on a LPWB- diet (Table 6). The dicalciumphosphate content of the diets varied from 16 g/kg in a HPWB- diet to 1 g/kg on a LPWB- diet. On LPWB+ diets, all P was of plant origin. Phytic P ranged from 0.305 to 0.560

and from 0.388 to 0.621 of total P in WB- and WB+ diets, respectively. Although the excretion of P in faeces decreased linearly ( $p < 0.01$ ) with a decreasing P content of the diet, the differences in excretion between treatments were not significant. P excretion ranged from 9.88 g/d on a HPWB- diet to 8.22 g/d on a LPWB+ diet. Also the apparent digestibility of P (% of intake), decreased linearly ( $p < 0.001$ ) with a decreasing P content of the diet. In the previously mentioned digestibility and balance study conducted by NÄSI and HELANDER (1993), the faecal excretion of P was about twice as high in diets with inorganic P supplementation as compared to unsupplemented diets. The absorption of P was then on an equal level (3.8-3.9 g/d) with P absorption on the LPWB- diet (4.0 g/d) in this study, although the daily intake of P in the present experiment was one-third higher. A possible reason for the low absorption rate of P in the present experiment may be related to the age of the pigs. In

Table 6. Excretion, apparent digestibility and retention of phosphorus in pigs fed on the experimental diets.

Treatment	HP		MP		LP		SEM	Statistical significance				
	WB-	WB+	WB-	WB+	WB-	WB+		C1	C2	C3	C4	C5
P intake, g/d	20.32	20.51	16.45	15.82	12.50	12.96	0.136	NS	***	**	NS	**
P excretion in faeces, g/d	9.88	9.75	9.20	9.33	8.51	8.22	0.461	NS	**	NS	NS	NS
P digested, g/d	10.44	10.76	7.24	6.48	3.99	4.74	0.474	NS	***	NS	NS	NS
P digestibility	0.514	0.528	0.442	0.408	0.322	0.369	0.0286	NS	***	NS	NS	NS
Urinary P excretion, g/d	3.99	3.94	2.06	1.82	0.39	0.23	0.164	NS	***	NS	NS	NS
P retained, g/d	6.45	6.82	5.19	4.67	3.59	4.51	0.503	NS	***	NS	NS	NS
-of intake	0.318	0.334	0.318	0.294	0.291	0.350	0.0305	NS	NS	NS	NS	NS
-of digested	0.621	0.620	0.718	0.724	0.875	0.946	0.0272	NS	***	NS	NS	NS
-g/W <sup>0.75</sup>	0.208	0.218	0.168	0.152	0.117	0.149	0.0147	NS	***	NS	NS	NS

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> (P < 0.10) o, (P < 0.05) \*, (P < 0.01) \*\*, (P < 0.001) \*\*\*, (P > 0.10) NS

C1 = WB- vs. WB+, C2 = P lin, C3 = P quadr, C4 = C1 × C2, C5 = C1 × C3

the earlier study, the pigs were younger and, thus, weighed less (weight 36-88 kg) than the pigs in this study (71-122 kg). JONGBLOED (1987) reviewed the adaptation of young animals to widely differing amounts of P and Ca in the diet by varying their intestinal absorption, and found that a decrease in the adaptation of the intestine to dietary P and Ca restriction occurs with increasing age. This is supported by the findings of NÄSI (1990): in older pigs the retention of P remained low in spite of a high P supply on a maize-soybean meal diet. Another explanation for the lower absorption rate of P on a LPWB- diet as compared to the earlier results may be the less favourable Ca:P ratio in this experiments (2.3 vs 1.8). A wide Ca:P ratio has an important negative influence on a low-phosphorus diet (De WILDE and JOURQUIN 1992). JONGBLOED (1987) concluded in his trials that the higher the Ca content of the diet, the more the absorption percentage of P and Ca decreased at increasing live weights.

Phosphorus homeostasis appears to be achieved mainly by regulating excretion via urine (LINDER 1991). When the P supply in the diet is low, the animal tries to retain the plasma P level constant by reabsorbing P from the kidneys. In the present

study, the urinary P excretion decreased linearly ( $p < 0.001$ ) with a decreasing P content of the diet. The proportion of absorbed P that was retained increased linearly ( $p < 0.001$ ) when the P content of the diet decreased, ranging from 0.620 on a HPWB+ diet to 0.946 on a LPWB+ diet. This is in agreement with earlier observations (NÄSI 1990, NÄSI and HELANDER 1993).

The inclusion of WB in the diet did not improve the digestibility of P. However, the amount of P excreted in urine was non-significantly lower on WB+ diets. The absorption of P appeared to be 0.75 g/d (18.8%) higher on the LPWB+ diet compared to the LPWB- diet. The proportion of absorbed P that was retained was 7.1 percentage units, i.e. almost 1 g/d higher on the LPWB+ diet than on the LPWB- diet. The P intake was 0.5 g/d higher on a LPWB- diet, but this hardly explains the differences in absorption and retention. Thus, it can be assumed that WB (phytase) had improved the utilization of phytic P on the low-P diet. The apparent P absorption was found to improve by 21% in a study by NEWTON et al. (1983), when a corn-soybean meal diet with a normal P level was supplemented with 100 g WB. Those results are, however, not fully

Table 7. Calcium excretion, apparent digestibility and retention in pigs fed on the experimental diets.

Treatment	HP		MP		LP		SEM	Statistical significance				
	WB-	WB+	WB-	WB+	WB-	WB+		C1	C2	C3	C4	C5
Ca intake, g/d	22.11	22.41	23.08	22.08	22.85	22.44	0.118	**	**	NS	**	***
Ca supply in water, g/d	0.11	0.11	0.12	0.11	0.12	0.11	0.005	NS	NS	NS	NS	NS
Ca excretion in faeces, g/d	14.34	15.05	15.10	14.66	15.30	14.66	0.546	NS	NS	NS	NS	NS
Ca absorption, g/d	7.88	7.47	8.10	7.53	7.67	7.89	0.530	NS	NS	NS	NS	NS
Ca absorption	0.355	0.332	0.349	0.339	0.335	0.352	0.0240	NS	NS	NS	NS	NS
Urinary Ca excretion, g/d	0.44	0.43	0.48	0.35	0.87	1.04	0.139	NS	**	*	NS	NS
Ca retained, g/d	7.44	7.03	7.62	7.18	6.81	6.84	0.568	NS	NS	NS	NS	NS
- of intake	0.335	0.312	0.328	0.324	0.296	0.304	0.0258	NS	NS	NS	NS	NS
- of absorption	0.943	0.946	0.942	0.955	0.893	0.866	0.0199	NS	**	NS	NS	NS
-g/W <sup>0.75</sup>	0.238	0.227	0.245	0.232	0.217	0.223	0.019	NS	NS	NS	NS	NS

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> (P < 0.10) o, (P < 0.05) \*, (P < 0.01) \*\*, (P < 0.001) \*\*\*, (P > 0.10) NS

C1 = WB- vs. WB+, C2 = P lin, C3 = P quadr, C4 = C1 × C2, C5 = C1 × C3

comparable with ours, because WB inclusion also enhanced the total P intake. POINTILLART (1991) reported that P from a diet to which no inorganic P was added and which was supplemented by 20% of rye bran, was better absorbed (55 vs 36%) and retained (50 vs 36%) by pigs of 12-43 kg live weight than P from the control diet.

The Ca intake of the pigs ranged between 22-23 g/d (Table 7). Ca digestibility was not affected by WB inclusion or by decreasing the P content of the diet, whereas urinary Ca excretion increased linearly (p<0.01) and quadratically (p<0.05) with a decreasing dietary P content. The results of the present study show that Ca balance is mainly regulated through the intestine as was also concluded by FERNÁNDEZ (1992). In an experiment by DEN HARTOG et al. (1988) apparent faecal digestibility of Ca decreased, when the diet was supplemented with 50 g pectin, cellulose or straw meal per kg. No significant differences were found in Ca retention due to treatments. However, the retention tended to be lower on LP than on other diets. JONGBLOED (1987)

reported the daily Ca retention to vary from 6 to 8 g after 55 kg live weight, which is well in accordance with our results. The proportion of digested Ca that was retained decreased linearly (p<0.01) when the P content of the diet decreased. In our previous digestibility and balance study (NÄSI and HELANDER 1993), Ca absorption was lower in a P-supplemented diet compared to an unsupplemented diet, but no significant differences were found in Ca retention because the urinary Ca excretion was also lower. POINTILLART (1991) reported no change in Ca absorption (53 vs. 50%) but higher Ca retention (52 vs. 45%) on a diet with 200 g inclusion of rye bran per kg feed than on a diet without any inorganic P supplement.

Mg intake was higher (p<0.001) on WB+ diets and decreased when the dietary P level decreased (p<0.001) (Table 8). This indicates that both WB and dicalciumphosphate have contributed some Mg to the diet. Daily Mg absorption did not differ between treatments. According to LINDER (1991), dietary Ca and P do not affect the absorption of Mg

Table 8. Magnesium excretion, apparent digestibility and retention in pigs fed on the experimental diets.

Treatment	HP		MP		LP		SEM	Statistical significance				
	WB-	WB+	WB-	WB+	WB-	WB+		C1	C2	C3	C4	C5
Mg intake, g/d	5.09	5.58	5.25	5.41	4.71	5.15	0.028	***	***	***	NS	***
Mg supply in water, g/d	0.50	0.50	0.56	0.50	0.55	0.52	0.022	NS	NS	NS	NS	NS
Mg excretion in faeces, g/d	3.56	3.80	3.58	3.87	3.54	3.59	0.107	*	NS	NS	NS	NS
Mg absorption, g/d	2.03	2.28	2.22	2.04	1.72	2.08	0.109	NS	*	NS	NS	*
Mg absorption	0.361	0.374	0.382	0.344	0.327	0.367	0.0186	NS	NS	NS	NS	NS
Urinary Mg excretion, g/d	0.66	0.66	0.70	0.72	0.87	0.88	0.040	NS	***	NS	NS	NS
Mg retained, g/d	1.37	1.62	1.52	1.32	0.85	1.19	0.127	NS	**	NS	NS	*
-of intake	0.243	0.246	0.261	0.223	0.163	0.212	0.0227	NS	**	NS	NS	NS
-of absorption	0.673	0.704	0.684	0.644	0.464	0.573	0.0431	NS	**	NS	NS	NS
-g/W <sup>0.75</sup>	0.044	0.051	0.049	0.042	0.028	0.039	0.0038	NS	**	NS	NS	*

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> (P < 0.10) o, (P < 0.05) \*, (P < 0.01) \*\*, (P < 0.001) \*\*\*, (P > 0.10) NS

C1 = WB- vs. WB+, C2 = P lin, C3 = P quadr, C4 = C1 × C2, C5 = C1 × C3

in humans (the studies have, however, usually been done with animals). Mg homeostasis occurs similarly as with P, and thus, mainly via adjustments of urinary excretion in humans (LINDER 1991). According to CRENSHAW (1991) the regulation of body Mg levels in pigs is not very well understood. In a study by NEWTON et al. (1983) WB inclusion was not found to affect the apparent digestibility of Mg in pigs in a study by NEWTON et al. (1983). In the present experiment, Mg excretion in urine increased ( $p < 0.001$ ) with a lower dietary P content. An interaction between WB inclusion and P level was recorded in Mg retention: on WB+ diets Mg retention decreased linearly ( $p < 0.05$ ) and on WB- diets both linearly ( $p < 0.05$ ) and curvilinearly ( $p < 0.05$ ) with a decreasing P content. These results contradict those of a study by NÄSI (1990), in which Mg absorption improved and retention remained unchanged on a unsupplemented maize-soybean meal diet compared with a diet supplemented by inorganic P. The results are, however, well in accordance with those of an earlier study by NÄSI and

HELANDER (1993), where barley-soybean meal diets were used.

K digestibility was not affected by WB inclusion (Table 9). This result is in conflict with the results of NEWTON et al. (1983), who reported that the apparent absorption of K decreased significantly when the diet was supplemented by 100 g of WB/kg and non-significantly when supplemented by 200 g of WB/kg. Urinary K excretion was increased slightly when WB was included in the diet. In K retention, an interaction was observed between WB and P level: on WB- diets the retention decreased curvilinearly ( $p < 0.05$ ) while on WB+ diets the retention remained constant when the P level of the diet decreased.

Due to unknown reasons there was a large variation in zinc (Zn) intake between treatments (Table 9). A high Zn intake led to a higher Zn excretion in faeces. The daily Zn absorption and retention were the highest on MPWB- and MPWB+ diets and were unaffected by WB inclusion. In retention, again, an interaction was found between

Table 9. Apparent digestibility and retention of potassium and zinc in pigs fed on the experimental diets.

Treatment	HP <sup>1</sup>		MP		LP		SEM	Statistical significance <sup>2</sup>				
	WB-	WB+	WB-	WB+	WB-	WB+		C1	C2	C3	C4	C5
<b>Potassium</b>												
Intake, g/d	24.70	25.79	27.19	26.07	25.26	26.67	0.399	NS	NS	**	NS	**
Digested	0.766	0.774	0.808	0.768	0.803	0.804	0.0138	NS	*	NS	NS	NS
Excretion in urine, g/d	15.86	15.38	16.93	15.98	17.35	17.52	0.235	*	***	NS	NS	NS
Retention, g/d	3.07	4.60	5.04	4.02	2.92	3.94	0.578	NS	NS	NS	NS	*
-of intake	0.127	0.175	0.187	0.156	0.117	0.148	0.0212	NS	NS	o	NS	NS
-of digested	0.163	0.224	0.229	0.202	0.144	0.183	0.0250	NS	NS	NS	NS	NS
<b>Zinc</b>												
Intake, mg/d	274	276	394	373	255	357	2.1	***	***	***	***	***
Digested	0.296	0.215	0.357	0.343	0.233	0.270	0.0352	NS	NS	**	NS	NS
Excretion in urine, mg/d	6.0	6.2	11.8	8.7	10.1	11.2	1.27	NS	**	NS	NS	NS
Retention, mg/d	75	54	129	121	49	85	9.95	NS	NS	***	*	NS
-of intake	0.274	0.193	0.327	0.319	0.193	0.238	0.0358	NS	NS	**	NS	NS
-of digested	0.924	0.978	0.915	0.939	0.820	0.877	0.0240	*	***	NS	NS	NS

<sup>1</sup> HP = high phosphorus (P), MP = medium P, LP = low P, WB- = no wheat bran, WB+ = 100 g wheat bran/kg diet.

<sup>2</sup> (P < 0.10) o, (P < 0.05) \*, (P < 0.01) \*\*, (P < 0.001) \*\*\*, (P > 0.10) NS

C1 = WB- vs. WB+, C2 = P lin, C3 = P quadr, C4 = C1 × C2, C5 = C1 × C3

WB inclusion and P level: on WB+ diets Zn retention increased linearly (p<0.05) and quadratically (p<0.001) and on WB- diets it decreased quadratically (p<0.001) with a decreasing dietary P level. In a study by NÄSI (1990) Zn absorption and retention were higher in diets without phosphate addition, but remained unaffected in a later study by NÄSI and HELANDER (1993). The proportion of digested Zn that was retained was higher on all WB+ diets than on WB-diets. In a trial made by NEWTON et al. (1983) Zn apparent absorption decreased, when the WB level of the diet increased. The indigestible, fibrous bran fraction was found to accumulate some minerals such as Zn. Unfortunately, the retention of minerals was not determined in that study.

The rather low effect of WB phytase on the digestibility of P and other minerals may be due to several reasons. The optimum temperature for wheat phytase, 55°C, is higher than the animal body temperature and the optimum pH for its activity is 5.15 (POMERANZ 1978). For pigs, the stomach

seems to be the principal place for the hydrolysis of phytate to utilizable P (JONGBLOED 1987). The low pH of the stomach can cause the inactivation of plant phytase. The solubility of wheat phytate has been reported to be lowest within the pH range from 1.5 to 2.5 (SCHEUERMANN et al. 1988a). According to SCHEUERMANN et al. (1988b), pepsin also inhibits the activity of wheat phytase. JONGBLOED (1987) found that an increase in dietary protein level improved the digestibility of P. In the present study, the pigs on WB+ diets ate less protein than the other pigs. This might also explain the low effect of WB on P utilization.

The age of the pigs may also partly explain the results. Older pigs may need less P in tissue metabolism and/or they are able to utilize phytate P better (REINHART and MAHAN 1986).

One reason for the poor effect of WB could be that the inclusion of 100 g WB used in this experiment per kg diet may not have been a sufficient amount. The calculated increase in phytase activity

in the diet was only 280 U/kg. The addition level of microbiologically produced phytase for growing-finishing pigs recommended by BASF (1993) is 500 U per kg feed, having a total P content of 4.0-5.0 g/kg.

The possible decreasing effect of pelleting on enzyme activity was discussed previously. Pelletting may, on the other hand, have a positive effect on the digestibility of P. BAYLEY et al. (1975) have reported improved P absorption due to steam pelleting. It is interesting to note that such an improvement was found only in diets not supplemented with inorganic P. The pelleting temperature, however, was not mentioned. JONGBLOED (1987) has also reported that the absorption and retention of P improved with pelleted diets.

In conclusion, the supplementation of a barley-soybean meal diet with 100 g WB per kg was not found to improve the dietary P utilization significantly. However, the digestion and retention of P appeared to be slightly improved by WB inclusion in the LP diet. This improvement may be due to WB phytase. The effects of WB on the digestibility and

balance of other minerals remained relatively small. The P level had a greater effect on mineral balances. The dry matter and organic matter digestibilities were impaired when 100 g of WB was included in the diet. N absorption was higher on WB+ diets. The retention of N appeared to decrease on HP and MP diets due to WB, but no decrease was found on LP diets. Ash digestibility was not affected by the treatments. The LPWB+ diet appeared to be the most favourable regarding the polluting P-emission, since the faecal excretion of P on that diet was the lowest and the overall retention coefficient of P was the highest. This conclusion can, however, be confirmed only after a growth trial. On the basis of the present results, an addition of 100 g of WB to the diet in order to improve the P utilization of finishing pigs, does not seem very beneficial.

*Acknowledgements.* The authors wish to thank Ms. M. Yli-aho for technical assistance. The financial support provided by the Academy of Finland and Suomen Rehu Ltd. is gratefully acknowledged.

## References

- AOAC 1984. Official methods of analysis. 1141 p. Association of Official Analytical Chemists, Inc., Arlington, Virginia.
- BAGHERI, S. 1985. Effect of wheat bran and pectin on the absorption and retention of phosphorus, calcium, magnesium and zinc by the growing pig. *Reproduction Nutrition Development* 25, 4A: 705-716.
- & GUEGUEN, L. 1982. Bioavailability to rats of calcium, magnesium, phosphorus and zinc in wheat bran diets containing equal amounts of these minerals. *Nutrition Report International* 25: 583-590.
- BAYLEY, H.S., POS, J. & THOMSON, R.G. 1975. Influence of steam pelleting and dietary calcium level on the utilization of phosphorus by the pig. *Journal of Animal Science* 40: 857-863.
- BASF 1993. Provisional technical leaflet of Natuphos<sup>R</sup> 5000. 4 p.
- COSGROVE, D.J. & IRVING, G.C.J. 1980. Inositol phosphates. Their chemistry, biochemistry and physiology. Elsevier. Amsterdam - Oxford - New York. 191 p.
- CRENSHAW, T.D. 1991. Sodium, potassium, magnesium and chloride in swine nutrition. In: Miller, E.R. et al. (eds.). *Swine nutrition*. 1th ed., Butterworth-Heinemann, Stoneham, MA, USA. p. 183-191.
- CROMWELL, G.L. 1992. The biological availability of phosphorus in feedstuffs for pigs. *Pig News and Information* 13, 2: 75N-78N.
- DEN HARTOG, L.A., HUISMAN, J., THIELEN, W.J.G., VAN SCHAYK, G.H.A., BOER, H. & VAN WEERDEN, E.J. 1988. The effect of including various structural polysaccharides in pig diets on ileal and faecal digestibility of amino acids and minerals. *Livestock Production Science* 18: 157-170.
- DIGGS, B.G., BECKER, D.E., JENSEN, A.H. & NORTON, H.W. 1965. Energy value of various feeds for the young pig. *Journal of Animal Science* 24: 555-558.
- ERDMAN, J.W. 1979. Oilseed phytates: Nutritional implications: *Journal of the American Oil Chemists Society* 56: 736-741.
- FERNÁNDEZ, J.A. 1992. Calcium and phosphorus metabolism in growing pigs studied by the balance technique and simultaneous radio-calcium and radio-phosphorus kinetics. The Royal Veterinary and Agricultural University Department of Anatomy and Physiology. Animal Physiology Unit. Thesis. 148 p. Fredriksberg, Denmark.
- HARRIS, R.S. 1955. Phytic acid and its importance in human nutrition: *Nutrition Reviews* 13, 9: 257-259.
- HOUSEMAN, R.A. & BRUYNE, K. de 1989. PhytinPhosphor und Phytase - Ein Überblick. *Kraftfutter* 4: 112-115.

- JONGBLOED, A.W. 1987. Phosphorus in the feeding of pigs. Effect of diet on the absorption and retention of phosphorus by growing pigs. 343 p. I.V.V.O. Rep.No.179. Lelystad, The Netherlands.
- , EVERTS, H. & KEMME, P.A. 1991. Phosphorus availability and requirements in pigs. In: Haresign, W. & Cole, D.J.A. (eds.). Recent Advances in Animal Nutrition 1991. Butterworths, London. p. 65-80.
- & KEMME, P.A. 1990. Effect of pelleting mixed feeds on phytase activity and the apparent absorbability of phosphorus and calcium in pigs. *Animal Feed Science and Technology* 28: 233-242.
- LINDER, M. C. 1991. Nutrition and metabolism of the major minerals. In: Linder, M. C. (ed.). Nutritional biochemistry and metabolism with clinical applications. p. 191-214. 2th ed. Els. Sci. Publ. Amsterdam.
- MROZ, Z., JONGBLOED, A.W., KEMME, P.A. & LENIS, N.P. 1991. Ileal and overall digestibility of nitrogen and amino acids in a diet for pigs as influenced by *Aspergillus niger* phytase and feeding frequency or levels. In: Eggum, B.O. et al. (eds.). Proceedings of the 6th International Symposium on Protein Metabolism and Nutrition. Herning, Denmark. p. 225-227.
- NÄSI, M. 1990. Microbial phytase supplementation for improving availability of plant phosphorus in the diet of the growing pig. *Journal of Agricultural Science in Finland* 62: 435-443.
- & HELANDER, E. 1993. Effects of microbial phytase supplementation and soaking of barley-soybean meal diet for improving availability of plant phosphorus for growing pigs. *Acta Agriculturae Scandinavica, Section A., Animal Science* 43 (in press).
- NELSON, T.S., SHIEH, R.R., WODINSKI, R.J. & WARE, J.H. 1971. Effect of supplemental phytase on the utilization of phytate phosphorus by chicks. *Journal of Nutrition* 101: 1289-1293.
- NEWTON, G.L., HALE, O.M. & PLANK, C.O. 1983. Effect of wheat bran in practical diets on mineral absorption by pigs at two ages. *Canadian Journal of Animal Science* 63: 399-408.
- PLAAMI, S. & KUMPULAINEN, J. 1991. Determination of phytic acid in cereals using ICP-AES to determine phosphorus. *J. Assoc. Off. Anal. Chem.* 74: 32-36.
- POINTILLART, A. 1988. Phytate phosphorus utilization in growing pigs. Proceedings of the 4th International Seminar on Digestive Physiology in Pigs. Polish Academy of Sciences, Jablonna. p. 319-326.
- 1991. Enhancement of phosphorus utilization in growing pigs fed phytate-rich diets by using rye bran. *Journal of Animal Science* 69: 1109-1115.
- POMERANZ, Y. 1978. Wheat Chemistry and Technology. American Association of cereal chemists, Incorporated. 821 p. 3th Ed., St. Paul, Minnesota. USA.
- REINHART, G.A. & MAHAN, D.C. 1986. Effect of various calcium:phosphorus ratios at low and high dietary phosphorus for starter, grower and finishing swine. *Journal of Animal Science* 63: 457-466.
- SALO, M.-L., TUORI, M. & KIISKINEN, T. 1990. Rehtaulukot ja ruokintanormit. 70 p. Yliopistopaino, Helsinki.
- SCHUEERMANN, S.E., LANTZSCH, H.-J. & MENKE, K.H. 1988a. In vitro und in vivo Untersuchungen zur Hydrolyse von Phytat. I. Löslichkeit von Phytat. *Journal of Animal Physiology and Animal Nutrition* 60: 55-63.
- , LANTZSCH, H.-J. & MENKE, K.H. 1988b. In vitro und in vivo Untersuchungen zur Hydrolyse von Phytat. Aktivität pflanzlicher Phytase. *Journal of Animal Physiology and Animal Nutrition* 60: 64-75.
- SCHWARZ, G. & SCHÖNER, F.-J. 1991. Phytase in der Fütterung von Schweinen und Geflügel. *Kraftfutter* 4: 163-168.
- SNEDECOR, G.W. & COCHRAN, W.G. 1989. Statistical methods. 503 p. 8th ed. Iowa University Press, Ames, Iowa.
- TAYSSKY, H.H. & SHORR, E. 1953. A microcolorimetric method for the determination of inorganic phosphorus. *Journal Biological Chemistry* 202: 675-685.
- WILDE, R.O., de & JOURQUIN, J. 1992. Estimation of digestible phosphorus requirements in growing-finishing pigs by carcass analysis. *Journal Animal Physiology and Animal Nutrition* 68: 218-225.

Manuscript received September 1993

SELOSTUS

**Vehnäleseeseen vaikutus kasviperäisen fosforin hyväksikäyttöön lihasian ruokinnassa  
I. Vaikutus ravintoaineiden sulavuuteen ja kivennäistaseisiin**

EIJA HELANDER, MATTI NÄSI ja KIRSI PARTANEN

Helsingin yliopisto

Sulavuus- ja tasekokeessa tutkittiin, parantaako vehnäleseeseen fytaasi ohrasta ja soijasta peräisin olevan fosforin hyväksikäyttöä ja voidaanko sillä näin ollen osittain tai kokonaan korvata rehuun normaalisti lisättävä epäorgaaninen fosfori.

Vehnälese ei parantanut merkittävästi fosforin hyväksikäyttöä. Kuitenkin fosforin sulavuus ja pidättyminen parani hieman ruokinnalla, jonka fosforitaso oli matala, kun siihen oli lisätty vehnäleseettä. Vehnäleseeseen vaikutus muiden kivennäisten sulavuuteen ja pidättymiseen jäi vähäiseksi. Fosforitasolla oli kivennäisten sulavuuteen ja pidättymiseen suurempi vaikutus. Tuhkan sulavuus pysyi muuttumattomana eri

ruokinnoilla. Kuiva-aineen ja orgaanisen aineen sulavuus heikkenivät vehnäleseeseen vaikutuksesta. Typen imeytymis-% oli korkeampi ( $p < 0.01$ ) vehnälesedieeteillä. Vehnälese näytti alentavan typen pidättymistä korkeimmalla ja keskinkertaisella fosforitasolla, mutta matalalla ei.

Fosforipäästöjä ajatellen matalafosforinen, vehnäleseettä sisältävä ruokinta osoittautui parhaaksi: ulosteiden fosforitaso oli alhaisin ja fosforin pidättymis-% oli korkein. Tulosten perusteella vehnäleseeseen käyttö sikojen rehuseoksissa fosforin hyväksikäytön parantamiseksi ei kuitenkaan näytä lupaavalta.