

Physiological performance of quinoa (*Chenopodium quinoa* Willd.) under agricultural climatic conditions in Boyaca, Colombia

Comportamiento fisiológico de la quinua (*Chenopodium quinoa* Willd.) bajo condiciones agroclimáticas de Boyacá, Colombia

Miguel García-Parra^{1,3*}, José García-Molano^{2,3}, and Yuli Deaquiz-Oyola^{2,3}

ABSTRACT

Quinoa (*Chenopodium quinoa* Willd.) is native to South America; it is characterized by its high nutrient contents and high adaptation capacity to diverse edapho-climatic conditions, which highlights its genetic variability expressed as multiple physiological and phenological responses. The objective of this research was to evaluate the physiological response and proximal composition of the grain to three types of fertilization under the environmental conditions of the municipality of Oicata (Boyaca, Colombia) located at 2,875 m a.s.l. The white Soracá variety was planted using a completely randomized design with four treatments and four replicates. It was observed that the fertilization sources have an effect on the physiological and phenological behavior, mainly on the number of leaves, length of stem and chlorophyll content. The reproductive stage and the proximal composition of seeds changed, which is attributed to the application of mineral organic fertilizer that improves the production of quinoa grains, while N-P-K contribution shows greater growth and vegetable development, but less yield.

Key words: chlorophyll, fertilizer, phenology, protein content.

RESUMEN

La quinua (*Chenopodium quinoa* Willd.) es originaria de América del Sur, y está caracterizada por la alta composición nutricional y su fácil adaptabilidad a condiciones edafoclimáticas, lo que resalta su amplia variabilidad genética que se expresa en múltiples respuestas fisiológicas y fenológicas. El objetivo de este estudio fue evaluar la respuesta fisiológica y la composición proximal del grano a tres tipos de fertilización bajo condiciones de clima y suelo propias del municipio de Oicatá (Boyacá, Colombia), localizado a 2.875 msnm. Se sembró quinua variedad blanca de Soracá, utilizando un diseño completamente al azar con cuatro tratamientos y cuatro repeticiones. Se observó que las fuentes de fertilización tienen efecto sobre el comportamiento fisiológico y fenológico, principalmente en el número de hojas, longitud de tallo y contenido de clorofila. Las etapas reproductivas y la composición proximal de las semillas también mostraron cambios, lo que se atribuye al aporte de abono orgánico-mineral que mejora la producción de granos de quinua, mientras que el aporte de N-P-K muestra mayor crecimiento y desarrollo vegetativo, pero menor rendimiento.

Palabras clave: clorofila, abono, fenología, contenido de proteína.

Introduction

Quinoa (*Chenopodium quinoa* Willd.) is considered a crop with great potential because of its high agronomic characteristics and nutritional value, and especially for its inclusion in children and elderly people's diets (Valcárcel-Yamany and Silva, 2012). According to Escuredo *et al.* (2014), this plant has the capacity to produce grains of high quality and protein content. Additionally, it contains amino acids such as lysine, threonine and methionine, which are considered as essential.

These nutritional characteristics are the result of environmental conditions, such as temperature, light intensity, relative humidity and precipitation. These conditions are key factors in the quality and number of grains per panicle (Morales *et al.*, 2017), as well as in the phenological and physiological performance of the plants related to the adaptive capacity to diverse environmental conditions (Winkel *et al.*, 2016). The plant has adaptive advantages that allow it to express a great productive potential. Quinoa (*Chenopodium quinoa*) has undergone diverse evolutionary

Received for publication: 17 November, 2018. Accepted for publication: 21 May, 2019

Doi: 10.15446/agron.colomb.v37n2.76219

¹ Doctorado en Ciencias Agrarias y Agroindustriales, Universidad del Cauca, Popayán (Colombia).

² Facultad de Ciencias Agrarias y Ambientales, Fundación Universitaria Juan de Castellanos (Colombia).

³ Grupo de Investigación Agricultura Organizaciones y Frutos AOF (Colombia).

* Corresponding author: miguelgarcia@unicauca.edu.co



processes, but it has been bred from crosses with *Chenopodium carnosulum* to acquire resistance to salinity problems, *Chenopodium petiolare* to obtain adaptability to droughts, and *Chenopodium pallidicaule* to receive tolerance to frosts (Jarvis *et al.*, 2017).

The quinoa plant has the capacity to develop alternative metabolic plasticity (Bazile *et al.*, 2014). This change is induced by conditions such as temperature, light intensity, nutritional status, relative humidity, and water availability (Morales *et al.*, 2017). In addition, it is able to undergo phenological, morphological, and physiological changes known as phenotypic plasticity.

On the one hand, the soil conditions, weather, and nutrient availability are important factors in the morpho-agronomic performance of the crop. However, the physical, chemical, and microbiological characteristics are specific to each place and mark plant development and the composition, quality and quantity of the quinoa grain (Veloza *et al.*, 2016).

Moreover, in Colombia, the most cultivated varieties of quinoa are Piartal and Tunkahuan which come from Ecuador, SL47 from Nariño, White from Jerico and White from Soraca and Boyaca (Ardila *et al.*, 2006). These quinoa genotypes have an average productivity of 1.5 and 2.6 t ha⁻¹ depending on the variety the fertilization plan (Delgado *et al.*, 2009) and the environmental conditions (García-Parra *et al.*, 2017).

Furthermore, the quinoa is planted at a small scale in the provinces of Nariño, Cundinamarca, Cauca and Boyaca. In Boyaca, there are reports of crops in the central zone (2,538-3,031 m a.s.l.) grown from a mix of seeds that affects crop productivity and grain quality (Veloza *et al.*, 2016).

In that order, it is necessary to evaluate the physiological performance and the composition of the quinoa grain in three types of fertilization and under the environmental conditions of the municipality of Oicata.

Materials and methods

The research was carried out from June 2016 to May 2017 in the municipality of Oicata (Boyaca, Colombia) with coordinates 5°22'48" N and 73°30'09" W and an elevation of 2,875 m a.s.l. The average temperature was 12°C, with 74.1% relative humidity and an average precipitation of 1018.9 mm per year (Tab. 1). The soil in which the experiment was established had Andic Dystrustepts and Vertic Haplustalf association (IGAC and UPTC, 2005).

White quinoa seeds from Soraca were used as plant material, which were stored by the Research Group Agricultura, Organizaciones y Frutos (AOF) for six months. Some of the fertilizers used were: Paz del Rio Fertilizer (PRF, Escoria Thomas, Colombia), Urea (U) and an organic fertilizer from The Agro-Ecological Farm Victoria (AEFV, Colombia).

The experiment area for the research was of 460 m², where four fertilization treatments with four replicates were established. These trials were performed based on the result of the soil analysis carried out by the Chemistry Laboratory of Soil, Water and Plants from Agrosavia (Tab. 2). The treatment T0 was the control, T1 was for the application of 6 kg of AEFV fertilizer, T2 was for the application of 3 kg of AEFV fertilizer plus 100 g of U and 50 g of PRF, and T3 was for the application of 200 g of U and 100 g of PRF (Tab. 3).

The response variables were: number of leaves, height of plants (rigid flex meter), days to reach six true leaves, and days to 50% flowering. In addition, days to milky grain and days to pasty grain state, chlorophyll (SPAD 502 plus, Konica-Minolta, Japan), dry and fresh weights of the plant (Drying oven HSY-75, 24 h at 104°C), grain productivity, protein amount (Kjeldahl Technique, NTC370), neutral detergent fiber in grain (Gravimetric determination, Van Soest AOAC 2002.4) and acid detergent fiber in grain (Gravimetric determination, Van Soest H₂SO₄) were measured every 15 d.

TABLE 1. Climatic conditions at the experimental plot.

Climate variation	June	July	August	September	October	November	December
Precipitation (mm)*	22.9	60.7	32	60.3	99.6	42.1	38.6
Solar brightness (h)**	119.7	108.8	161.7	151.8	163.7	151.8	162.3
Relative humidity (%)	77.1	78.6	75.1	74.7	74.2	74.7	72.3

*Pluviometer data of Oicata at an altitude of 2,645 m a.s.l. Code 24030450 IDEAM.

**Data supplied by the weather station from IDEAM (Universidad Pedagógica y Tecnológica de Colombia).

TABLE 2. Soil characteristics at the study site.

pH		Organic matter (%)		CECe (cmol/kg)	
5.52		3.85		4.83	
Interchangeable bases (cmol/kg)					
Ca		Mg		K	
2.5		1.12		1.07	
Na					
0.14					
Microelements (mg/kg)					
P		Fe		Mn	
10.06		282.4		9.86	
Zn		Cu		B	
3.1		3.66		0.13	
S		8.28			

TABLE 3. Chemical characteristics of fertilizers.

	PRF (%)	U (%)	AEFV (%)
Total Nitrogen	-	46	1.86
Total phosphorus (P ₂ O ₅)	11	-	2.44
Potassium (K)	-	-	2.45
Sodium (Na)	-	-	0.27
Calcium (CaO)	40	-	1.23
Magnesium (MgO)	1.5	-	0.86
Silicon (SiO ₂)	6	-	-
Zinc (Zn)	0.001	-	0.01
Copper (Cu)	0.001	-	-
Cobalt (Co)	0.0002	-	-
Boron (B)	0.0002	-	-
Molybdenum (Mo)	0.0013	-	-
Sulfur (S)	-	-	0.36
pH	12	7.2	7.8
Humidity (%)	0.3	1	17.9

PRF: Paz del Rio fertilizer; U: Urea; AEFV: Agro-Ecological Farm Victoria fertilizer.

Statistical design

A completely randomized design with four replicates and 16 separate experimental units was established. The data obtained in the experiment were tabulated using Excel program; a test of homogeneity of variety with the Bartlett method and a normality test with the Shapiro-Wilk method were performed. An analysis of variance (ANOVA) and a Tukey test of comparison of means with a 0.05 significance level were performed using the program R version 3.3.0.

Results and discussion

Number of leaves

As shown in Figure 1, significant changes, such as an increase in the number of leaves, were observed after day 75. However, after the start of the flowering stage (105 d), the plant showed a loss of leaves to its senescence. The highest average was found in the T3 treatment (90.75),

which presented significant differences compared to the others, whereas T0 had the lowest average (32.75). All the treatments displayed a sigmoid performance. This performance was expressed in the exponentially development of the plant regarding the time (Taiz and Zeiger, 2007) taking into account that the plant induces, in the vegetative stages, the greatest number of foliar buds capable of capturing and housing the assimilates that will influence the reproductive stages and the formation of grains (Atencio *et al.*, 2014).




FIGURE 1. Number of quinoa leaves under different fertilization protocols in Oicata - Boyaca. T0: absolute control, T1: organic fertilizer, T2: organic fertilizer + urea + Paz del Rio fertilizer, and T3: urea + Paz del Rio fertilizer according to the Tukey test ($P \leq 0.05$). ns: not significant, *: significant for the day of sampling.

Although the number of leaves is a determining variable in the production of quinoa, it is not a key factor to crop yield. Similarly, the application of elements such as nitrogen stimulated the excessive development of fodder, which could affect the grain productivity (Kakabouki *et al.*, 2018). Correspondingly, García *et al.* (2017) stated that the application of an increasing dose of N-P-K in quinoa enhanced leaf area but not productivity. Nevertheless, the opposite occurs when organic-mineral fertilizer is used.

This kind of amendment facilitates the absorption of important elements for energetic, metabolic and enzymatic activities of the plant due to the microorganism content (Ramzani *et al.*, 2017).

Plant height

Statistical differences ($P < 0.05$) in height were recorded from day 75 for the T3 treatment. This contrast resulted in the highest average during the days 75, 90, 105, 135 and 150. The final average was 172.25 cm, compared to the absolute control treatment T0, which resulted in 93.25 cm (Fig. 2). Following day 120, the T2 treatment increased its growth due to the fertilization plan that was provided to each treatment. This increase could be due to the fact that nitrogen can stimulate plant elongation and cellular division, which allows the synthesis of auxin and cytokinins that have an effect on stem growth (Basra *et al.*, 2014).




FIGURE 2. Height of quinoa plants under different fertilization protocols in Oicata - Boyaca. T0: control, T1: organic fertilizer, T2: organic fertilizer + urea + Paz del Rio fertilizer, and T3: urea + Paz del Rio fertilizer according to the Tukey test ($P \leq 0.05$). ns: not significant, *: significant for the day of sampling.

As a result, when more nitrogen is applied and there is more precipitation, an increase of the stem elongation is evident. According to Fghire *et al.* (2015), the continuous cell division and elongation increase the permeability of the root meristems and facilitates the absorption of structural minerals such as calcium, considered as fundamental in tissue rigidity and an active agent in the manifestation of plant hormones that stimulate the elongation and cellular division.

Chlorophyll content

This variable presented important statistical differences among days 90, 105, 120 and 130. T3 showed the highest

average (66.4 SPAD units) in the closest phase to the flowering, at the moment in which the plant displayed the highest number of leaves. However, at 150 d, the treatments did not show significant differences (Fig. 3). This could be due to the fact that nitrogen application and chlorophyll content are directly proportional to each other (Liu *et al.*, 2015), along with the application urea in treatments T2 and T3.




FIGURE 3. Chlorophyll content of quinoa under different fertilization protocols in Oicata - Boyaca. T0: Absolute control, T1: organic fertilizer, T2: organic fertilizer + urea + Paz del Rio fertilizer, and T3: urea + Paz del Rio fertilizer according to the Tukey test ($P \leq 0.05$). ns: not significant, *: significant for the day of sampling.

In addition, nitrogen determines the content of assimilates existing in the leaves, and consequently, the color of the leaf through which the chlorophyll meters act on the wave capture. Then, the high nitrogen applications result in high chlorophyll contents (Fghire *et al.*, 2015), as observed in the experiment. For this reason, it is important to measure the chlorophyll content to evaluate the nutritional status of plants at foliar level (Rincón and Ligarreto, 2010).

Therefore, it is evident that the highest chlorophyll content is equivalent to vegetative phases and it is reduced in the reproductive phases (Fig. 3). According to García-Parra *et al.* (2018), the plant captures the highest amount of mineral elements in these phases, while in the filling and maturation phases, the nutrients are finally accumulated in the seeds.

Fresh weight

Each one of the evaluated organs presented significant statistical differences ($P < 0.05$), which evidences that the T2 treatment had greater fresh weight in roots and leaves. In contrast, T3 displayed the highest average in the weight of




FIGURE 4. Fresh weight of stems, roots, leaves and panicles of quinoa under different fertilization protocols in Oicata - Boyaca. T0: control, T1: organic fertilizer, T2: organic fertilizer + urea + Paz del Rio fertilizer, and T3: urea + Paz del Rio fertilizer according to the Tukey test ($P \leq 0.05$). ns: not significant, *: significant for the day of sampling.

the stem and panicle (Fig. 4). These results were obtained as a response to the implementation of mineral organic fertilizers (Garcia-Parra *et al.*, 2017). Moreover, the high precipitations during the final phase of the vegetative stages and at the beginning of the reproductive stages generated the dilution of the phosphate fertilizer, which is indispensable for the development of diverse energetic activities. Similarly, the contribution of calcium is fundamental in the expression of plant hormones, which stimulate the development of leaves, stems and roots.

The fresh weight of plants is an indicator of efficiency in the uptake of nutrients and water (Torres *et al.*, 2000), which can be influenced by environmental conditions. Thus, the biggest part of fresh biomass is found in the panicles followed by the stems, because the panicles support the leaves, glomerulus, and seeds, and are the duct for

important substances in the nutrition of quinoa (Al-Naggar *et al.*, 2017).

Dry weights

The plants treated with different fertilization protocols under the edaphoclimatic conditions of the municipality of Oicata presented significant statistical differences ($P < 0.05$) regarding the dry weight of the roots, stems, leaves, and panicles. These results showed that the treatments T2 and T3 had the highest average during the entire test (Fig. 5) because of the implementation of the organic material that acts as a retainer of moisture (García, 2006). This facilitates the dilution of urea, Paz del Rio fertilizer, and minerals that are present in the AEF fertilizer, all expressed as assimilates in dry matter. In the case of T3, the increment occurred because of the application of soluble fertilizer and the increase of precipitation during the experiment.




FIGURE 5. Dry weights of quinoa stems, roots, leaves and panicles under different fertilization protocols in Oicata - Boyaca. T0: Absolute control, T1: organic fertilizer, T2: organic fertilizer + urea + Paz del Rio fertilizer, and T3: urea + Paz del Rio fertilizer according to the Tukey test ($P \leq 0.05$). ns: not significant, *: significant for the day of sampling.

Furthermore, the dry matter is an indicator of the capacity that plants have to absorb nutrients (Magolbo *et al.*, 2015). Aliro *et al.* (2011) stated that in the vegetative stages the highest dry matter accumulation occurs in leaves, while in grain formation the elements, minerals, and photo-assimilated compounds are transported from the source to the sink organs.

Nevertheless, the high indexes of dry weight do not certainly indicate either the productivity of the plant or the quality of its composition. It is due to the fact that quinoa expresses diverse potentials according to its variety and origin; the nutrients absorbed not only reach sink organs like seeds, but they are also stored in source organs as reserve, which are determining in the dry weight of leaves, roots and stems (Jayme-Oliveira *et al.*, 2017).

Phenological performance

According to Table 4, the differentiation of the phenological performance started at flowering stage when treatments T1 and T2 took less time to get into their reproductive stages. However, after reaching the stages of milky and pasty grain, T0 presented a shorter productive cycle compared to T1, T2 and T3. This effect was due to the availability of nitrogen, which stimulated the longevity of the plant tissue, generating a constant production of new cells that are expressed in longer productive cycles.

The phenological response by the crop occurs through determining factors such as the soil and weather (Vargas *et al.*, 2015). For this reason, quinoa plants allow to display longer productive cycles when fertilization plans are carried out with N-P-K in excess (Gómez and Aguilar, 2016).

TABLE 4. Phenological performance of quinoa according to the treatments.

Treatment	Days to six true leaves	Days to 50% flowering	Days to milky grain	Days to pasty grain
T0	27±0.0 a	119.5±3.3 a	142.2±1.5 c	171.5±1.0 a
T1	25.7±0.9 a	118.5±1.7 ab	143±0.0 c	173±0.8 b
T2	26.5±0.5 a	118.7±3.5 ab	150±0.0 b	188±1.6 c
T3	26.7±0.5 a	124±0.0 b	154.2±3.4 a	202±2.1 d

Different letters in the same column indicate significant differences according to the Tukey test ($P \leq 0.05$). T0: absolute control; T1: organic fertilizer; T2: organic fertilizer + urea + Paz del Rio fertilizer, and T3: urea + Paz del Rio fertilizer.

TABLE 5. Production and composition of the quinoa grain.

Treatment	Grain productivity (g/m ²)	Protein (%)	NDF (%)	ADF (%)
T0	243±1.6 a	13.9±1.6 a	15± 1.1 a	10.79±0.9 a
T1	389±3.9 a	14.7±0.5 a	16.11± 0.2 a	10.24±0.3 a
T2	298.2±1.7 a	14.9±2.8 a	14.08±0.8 a	9.72±0.2 b
T3	374.2±1.7 a	13.6±0.5 a	15.69±0.3 a	11.64±1.2 c

NDF: Neutral Detergent Fiber. ADF: Acid Detergent Fiber. Different letters in the same column indicate significant differences according to the Tukey test ($P \leq 0.05$). T0: absolute control, T1: organic fertilizer, T2: organic fertilizer + urea + Paz del Rio fertilizer, and T3: Urea + Paz del Rio Fertilizer.

Moreover, and regarding the climatic factors, the activity of the Rubisco activase (enzyme) declines as temperatures increase above the thermal optimum of photosynthesis. This loss of activity causes Rubisco deactivation, which in turn is proposed to reduce photosynthetic capacity at elevated temperatures (Raines 2011).

In addition, elements such as nitrogen determine how long the quinoa plant can be harvested (Geren, 2015). This growth effect is triggered when this element stimulates plant hormones that generate the production of foliar buds, which are indispensable in the photosynthesis process. It also composes the chlorophyll molecule that is found between 40-52 SPAD units in the vegetative phases (García-Parra *et al.*, 2017).

Another determining factor is the edapho-climatic conditions during the establishment and development of the crop. The metabolic and phenotypical plasticity processes adapt to the water levels, sun radiation and relative humidity, which are all variables that intervene in the metabolic activities of the plant. These variables have allowed the plant to adapt to diverse regions of the world (Tabaglio *et al.*, 2015).

Production and composition of the grain

Grain production did not show significant statistical differences ($P < 0.05$). However, T1 showed a greater productivity per square meter, while protein, neutral detergent fiber (NDF) and acidic detergent fiber (ADF) had the best results in T2, T1 and T3, respectively (Tab. 5). The protein

of T1 could be higher because it does not contribute to the available nitrogen despite of the organic fertilizer supply; the population of diazotrophic bacterium existing in this kind of fertilizer influences its availability in the plant (Parra-Cota *et al.*, 2014). Additionally, a balance in elements like phosphorus, magnesium and other microelements contribute to the synthesis of protein (Marschner, 2012).

On the one hand, quinoa production is determined by factors such as the availability of elements in the soil. For this reason, even though no significant differences were observed in the grain productivity, treatments T3 and T2 outweigh the amount reported by Delgado *et al.* (2009). Moreover, the amount of protein obtained is between the ranges established by Jacobsen (2003), which are between 13 and 18%. These ranges are higher in comparison to cereals such as rice (7.5%) and corn (13.4%) (Elsohaimy *et al.*, 2015).

Regarding NDF content, the amount was higher in T1 seeds. However, the values for every treatment are higher in comparison to what was obtained by Peiretti *et al.* (2013), who reported values of 12.75%. The amount of ADF in the test was low compared to studies by Simranpreet *et al.* (2017), who reported 77.73%. These results were obtained because the composition and presence of fiber are influenced by the maturity stage of the seed (Reguera *et al.*, 2018). The results in this test are related to the fact that when plants started the grain phenological stage, precipitations caused the elongation of the panicle that formed new seeds. This prolonged the phenological period of the plant

to seven months; in other words, two more months than the phenological average of the variety in previous tests.

It is noteworthy that reports of ADF and NDF are very diverse in different tests reported in around the world. This could be happening because of the variety, maturity stage of the grain and fertilization, as referenced by Peiretti *et al.* (2013), whose results were 12.75% of ADF and 5.49% of NDF. On the other hand, Marmouzi *et al.* (2015) reported ADF of 72.03% and ADF of 27.06%, similar to the results of Simranpreet *et al.* (2017), with values of 77.73% and 27.4% for NDF and ADF, respectively.

Conclusions

Quinoa under the high Andean tropic conditions has a better reaction and performance reflected on grain productivity and quality when it is amended with organic and mineral fertilizers. Nonetheless, from a physiological activity perspective, the plant responded better to mineral fertilization. Even though the organic fertilizer did not contribute to the nitrogen available to the plant, the population of diazotrophic bacteria native to this environment is efficient in the contribution of nitrogen for plant development.

Acknowledgments

We would like to thank the staff members of the farm San Miguel located in the village of Poravita in Oicata, Boyaca for their support. We would also like to express our gratitude to the staff of the Agro-Ecological Farm Victoria located in Ventaquemada, Boyaca for their cooperation in this experiment. This study was supported by Colciencias and Gobernacion de Boyaca, Call 779/2017.

Literature cited

Al-Naggar, A., R. El-Salam, A. Badram, and M. El-Moghazi. 2017. Genotype and drought effects on morphological, physiological and yield traits of quinoa (*Chenopodium quinoa* Willd.). *Asian J. Advanc. Agric. Res.* 3(1), 1-15. Doi: 10.9734/AJAAR/2017/36655

Atencio, L., J. Tapia, S. Mejía, and J. Cadena. 2014. Comportamiento fisiológico de gramíneas forrajeras bajo tres niveles de humedad en condiciones de casa malla. *Temas Agrarios* 19(2), 245-259.

Basra, S.M., S. Iqbal, and I. Afzal. 2014. Evaluating the response of nitrogen application on growth development and yield of quinoa genotypes. *Int. J. Agric. Biol.* 16, 886-892.

Bazile, D., D. Bertero, and C. Nieto. 2014. Estado del arte de la quinua en el mundo 2013. Ed. FAO (Santiago de Chile) and CIRAD (Montpellier).

Delgado, A., J. Palacios, and C. Betancourt. 2009. Evaluación de 16 genotipos de quinua dulce (*Chenopodium quinoa* Willd.) en el municipio de Iles, Nariño (Colombia). *Agron. Colomb.* 27(2), 159-167.

Elsohaimy, S., T. Refaay, and M. Zaytoun. 2015. Physicochemical and functional properties of quinoa protein isolate. *Ann. Agric. Sci.* 60(2), 297-305. Doi: 10.1016/j.aos.2015.10.007

Escuredo, O., M.I.G. Martín, G.W. Moncada, S. Fischer, and J.M.H. Hierro. 2014. Amino acid profile of the quinoa (*Chenopodium quinoa* Willd.) using near infrared spectroscopy and chemometric techniques. *J. Cereal Sci.* 60(1), 67-74. Doi: 10.1016/j.jcs.2014.01.016

Fghire, R., F. Anaya, O. Ali, O. Benlhabib, R. Ragab, and S. Wahbi. 2015. Physiological and photosynthetic response of quinoa to drought stress. *Chil. J. Agr. Res.* 72(2), 174-183. Doi: 10.4067/S0718-58392015000200006

García, M. 2006. Principios generales de la agricultura orgánica. Fundación Universitaria Juan de Castellanos, Tunja, Colombia.

García-Parra, M., J. García-Molano, D. Melo-Ortiz, and Y. Deaquiz-Oyola. 2017. Respuesta agronómica de la quinua (*Chenopodium quinoa* Willd.) variedad dulce de Soracá a la fertilización en Ventaquemada - Boyacá. *Cultura Científica* 15, 66-77.

García-Parra, M., J. García-Molano, and D. Carvajal-Rodríguez. 2018. Evaluación del efecto de la fertilización química y orgánica en la composición bromatológica de semillas de quinua (*Chenopodium quinoa* Willd.) en Boyacá-Colombia. *Revista Invest. Agrar. Ambient.* 9(2), 99-107. Doi: 10.22490/21456453.2282

Geren, H. 2017. Effects of different nitrogen levels on the grain yield and some yield components of quinoa (*Chenopodium quinoa* Willd.) under mediterranean climatic conditions. *Tur. J. Field Crops* 20(1), 59-64. Doi: 10.17557/39586

Gómez, L. and F. Aguilar. 2016. Guía del cultivo de la quinua. Universidad Nacional Agraria la Molina. Lima, Peru.

IGAC and UPTC. 2005. Estudio general de suelos y zonificación de tierras del departamento de Boyacá. 1st ed. IGAC - UPTC, Bogota.

Jacobsen, S. 2003. The worldwide potential for quinoa (*Chenopodium quinoa* Willd.). *Food Rev. Int.* 19, 167-177. Doi: 10.1081/FRI-120018883

Jarvis, D., Y. Ho, D. Lightfoot, S. Schmöckel, B. Li, T. Borm, H. Ohyanagi, K. Mineta, C. Michell, N. Saber, N. Kharbatia, R. Rupper, A. Sharp, N. Dally, B. Boughton, Y. Woo, G. Gao, E. Schijlen, X. Guo, A. Momin, S. Negrao, S. Al-Bili, C. Gehring, U. Roessner, C. Jung, K. Murphy, S. Arold, T. Gojobori, C. Linden, E. Loo, E. Jellen, P. Maughan, and M. Tester. 2017. The genome of *Chenopodium quinoa*. *Nature* 542(7641), 307-312. Doi: 10.1038/nature21370

Jayme-Oliveira, A., W. Ribeiro, M. Gerosa, A. Camargo, and A. Jakelaitis. 2017. Amaranth, quinoa, and millet growth and development under different water regimes in the Brazilian Cerrado. *Pesqui. Agropecu. Bras.* 52(8), 561-571. Doi: 10.1590/S0100-204X2017000800001

Kakabouki, I., D. Hela, I. Roussis, P. Papastylianou, A. Sestras, and D. Bilalis. 2018. Influence of fertilization and soil tillage on nitrogen uptake and utilization efficiency of quinoa crop

- (*Chenopodium quinoa* Willd.). J. Soil. Sci. P. Nut. 18(1), 220-235. Doi: 10.4067/S0718-95162018005000901
- Liu, Z., H. Hu, H. Yu, X. Yang, H. Yang, C. Ruan, Y. Wang, and J. Tang. 2015. Relationship between leaf physiologic traits and canopy color indices during the leaf expansion period in an oak forest. *Ecosphere* 6(12), 259. Doi: 10.1890/ES14-00452.1
- Magolbo, L., E. Carmo, E. Garcia, A. Fernandes, and M. Leonel. 2015. Dry matter accumulation and mineral nutrition of aracacha in response to nitrogen fertilization. *Pesq. Agropec. Bras.* 50(8), 670-680. Doi: 10.1590/S0100-204X2015000800005
- Marschner, P. 2012. Mineral nutrition of higher plants. 3rd ed. Elsevier, USA.
- Marmouzi, I., N. Madani, Z. Charrouf, Y. Cherrah, and M. Abbas. 2015. Proximate Analysis, fatty acid and mineral composition of processed Moroccan, *Chenopodium quinoa* Willd. and antioxidant properties according to the polarity. *Phytotherapie* 13, 110-117. Doi: 10.1007/s10298-015-0931-5
- Maliro, M., V. Guwela, J. Nyaika, and K. Murphy. 2017. Preliminary studies of the performance of quinoa (*Chenopodium quinoa* Willd.) genotypes under irrigated and rainfed conditions of central Malawi. *Front. Plant Sci.* 8, 227. Doi: 10.3389/fpls.2017.00227
- Morales, A., A. Silva, J. Maldonado, and H. Silva. 2017. Transcriptional responses of Chilean Quinoa (*Chenopodium quinoa* Willd.) under water deficit conditions uncovers ABA-Independent expression patterns. *Front. Plant. Sci.* 8, 216. Doi: 10.3389/fpls.2017.00216
- Parra-Cota, F., J. Peña-Cabrales, S. Santos-Villalobos, N. Martínez-Gallardo, and J. Délano-Frier. 2014. *Burkholderia ambifaria* and *B. carobensis* promote growth and increase yield in grain Amaranth (*Amaranthus cruentus* and *A. hypocondriacus*) by improving plant nitrogen uptake. *PLoS ONE* 9(2), e88094. Doi: 10.1371/journal.pone.0088094
- Peiretti, P.G., F. Gai, and S. Tassone. 2013. Fatty acid profile and nutritive value of quinoa (*Chenopodium quinoa* Willd.) seeds and plants at different growth stages. *Anim. Feed. Sci. Tech.* 183, 56-61. Doi: 10.1016/j.anifeedsci.2013.04.012
- Raines, C. 2011. Increasing photosynthetic carbon assimilation in C₃ plants to improve crop yield: current and future strategies. *Plant Physiol.* 155, 36-42. Doi: 10.1104/pp.110.168559
- Ramzani, P., L. Shan, S. Anjum, W. Khan, H. Ronggui, M. Iqbal, Z. Virk, and S. Kausar. 2017. Improved quinoa growth, physiological response, and seed nutritional quality in three soils having different stresses by the application of acidified biochar and compost. *Plant Physiol. Biochem.* 116, 137-138. Doi: 10.1016/j.plaphy.2017.05.003
- Reguera, M., C. Conesa, A. Gil-Gómez, C. Haros, M. Pérez-Casas, V. Briones-Labarca, L. Bolaños, I. Bonilla, R. Álvarez, K. Pinto, A. Mujica, and L. Bascaña-Godoy. 2018. The impact of different agroecological conditions on the nutritional composition of quinoa seeds. *Peer J.* 6(e4442). Doi: 10.7717/peerj.4442
- Simranpreet, K., N. Kaur, and J. Gill. 2017. Effect of processing on the nutritional composition of quinoa (*Chenopodium quinoa* Willd.). *Agric. Res. J.* 54(1), 90-93. Doi: 10.5958/2395-146X.2017.00015.1
- Tabaglio, V., D. Melo, C. Ganimede, R. Boselli, and A. Vercesi. 2015. Prime esperienze di coltivazione della quinoa (*Chenopodium quinoa* Willd.). In: Pianura Padana. Convegno Nazionale Società Italiana Di Agronomia. Piacenza, Italy. Doi: 10.13140/RG.2.1.3544.4728
- Taiz, L. and E. Zeiger. 2007. Fisiología vegetal. Universitat Jaume, Castello, Spain.
- Torres, J., H. Vargas, G. Corredor, and L. Reyes. 2000. Caracterización morfoagronómica de diecinueve cultivares de quinua (*Chenopodium quinoa* Willd.) en la sabana de Bogota. *Agron. Colomb.* 17(3), 61-68.
- Valcárcel-Yamany, B. and S. Silva. 2012. Applications of quinoa (*Chenopodium quinoa* Willd.) and amaranth (*Amaranthus* spp.) and their influence in the nutritional value of cereal based foods. *Food Public Health* 2(6), 265-275. Doi: 10.5923/j.fph.20120206.12
- Vargas, D., M. Boada, L. Araca, W. Vargas, and R. Vargas. 2015. Agrobiodiversidad y economía de la quinua (*Chenopodium quinoa* Willd.) en comunidades aymaras de la cuenca del Titicaca. *Idesia* 33(4), 81-87. Doi: 10.4067/S0718-34292015000400011
- Veloza, C., G. Romero, and J. Gómez. 2016. Respuesta morfoagronómica y calidad en proteína de tres accesiones de quinua (*Chenopodium quinoa* Willd.) en la sabana norte de Bogota. *Rev. U.D.C.A Act. Div. Cient.* 19(2), 325-332.
- Winkel, T., P. Bommel, M. Chevarría-Lazo, G. Cortés, C. Del castillo, P. Gasselin, F. Lèger, J. Nina-Laura, S. Rambal, M. Tichit, J. Tourrand, J. Vacher, A. Vassas-Toral, M. Vieira-Pak, and R. Joffre. 2016. Panarchy of an indigenous agroecosystem in the globalized market: the quinoa production in the Bolivian Altiplano. *Glob. Environ. Change* 39, 195-204. Doi: 10.1016/j.gloenvcha.2016.05.007