

Ecology of rare water plant communities in lakes of north-eastern Poland

Ewa Jabłońska*, Stanisław Kłosowski

Institute of Botany, University of Warsaw, Al. Ujazdowskie 4, 00-478 Warsaw, Poland

Abstract

Habitat studies were conducted on three rare plant communities dominated by *Nuphar pumila*, *Nymphaea candida* and *Hydrilla verticillata* in lakes of north-eastern Poland. The comparison of habitat properties of these three types of phytocoenoses with those of *Nuphar lutea* common in the area under study was also performed. It was demonstrated that the plant communities studied were ecologically distinct. The habitats of the phytocoenoses of *N. pumila* differed most significantly from those of the other phytocoenoses. They often inhabited softer waters poor in Mg^{2+} , dissolved SiO_2 , but rich in total Fe, PO_4^{3-} , NO_3^- , and were associated with acidic substrates containing lower levels of Ca^{2+} and Na^+ , but greater amounts of total Fe and NO_3^- . The differences in the habitats of *H. verticillata* and *N. candida* phytocoenoses were most pronounced in the case of four properties of water: Na^+ , K^+ , Cl^- , and Mg^+ . Their values were lower in waters of the *H. verticillata* phytocoenoses. The habitats of all the three types of rare phytocoenoses differed considerably from those of *N. lutea*. The most significant differences were found between the *N. lutea* and *N. pumila* phytocoenoses and the smallest differences were between the patches of *N. lutea* and *N. candida*. The properties of water were more important in differentiating the habitats of the phytocoenoses studied than the substrate properties. Due to alkalization and increase in water hardness in the lakes studied the stands of *N. pumila* are among the most threatened. The patches of *N. candida* and *H. verticillata*, which occur in waters with a wider range of hardness and tolerating a slight increase in trophy, can still continue to persist in the lakes for a long time.

Keywords: rare plants, floating leaved and submerged phytocoenoses, properties of water and substrate, comparative analysis

Introduction

The north-eastern area of Poland (Lithuanian Lake District) is dominated by a postglacial landscape with diverse geological landforms (terminal moraines, sanders and eskers) left behind by retreating glaciers. These are accompanied by forests and numerous lakes supporting a rich vegetation [1]. There are no larger industrial plants or urban agglomerations in this area, which is less populated than the other regions of Poland. As a result the lakes remain in good condition and the vegetation has preserved to a considerable extent its original natural character. Therefore some rare plant communities dominated by such species as *Hydrilla verticillata* or *Nymphaea candida* are found to occur more frequently in this region of Poland than in other parts of the country [2-4]. Moreover, the analysed area hosts the only rich population of the glacial relict *Nuphar pumila* isolated from the main Polish area of occurrence in the Pomerania Lake District [5,6]. Data regarding the ecology of these plant communities were published in other papers

as a result of continued phytosociological and habitat studies conducted in north-eastern Poland (e.g. [4,7-11]). The investigations performed in the years 2002-2005 in 66 lakes within Lithuanian Lake District revealed new localities of *N. candida* and *H. verticillata* phytocoenoses in Poland. As in the case of the plant communities surveyed earlier they were examined with respect to their species composition, structure and habitat conditions. The numerous phytosociological data available for these three rare plant communities enabled the comparative analysis of the above types of phytocoenoses. The objective of this study, among others, was to define the ecological amplitude and the optimum growth conditions for these three rare plant communities. The present work also aimed at determining the degree to which they differed with respect to their ecology from the phytocoenoses of *N. lutea*, which are common in Poland. In addition an attempt has been made to predict the chances of survival of the phytocoenoses of *N. pumila*, *N. candida* and *H. verticillata* in lakes of the area under study.

Material and methods

The phytosociological and habitat data were collected at the height of the growing season (from the end of June to the end of August) in the years 1978-2004. Due to the distinct character of the littoral zone of lakes (e.g. [12,13]) as well as hydrochemical differences even between neighbouring patches of vegetation within the various bodies of water [14,15], the

* Corresponding author. Email: ejablonska@uw.edu.pl

This is an Open Access digital version of the article distributed under the terms of the Creative Commons Attribution 3.0 License (creativecommons.org/licenses/by/3.0/), which permits redistribution, commercial and non-commercial, provided that the article is properly cited.

investigations were carried out on particular phytocoenoses, irrespective of the lake's trophic status. Studies were conducted on 18 phytocoenoses of *N. pumila* from 5 lakes, 14 phytocoenoses of *H. verticillata* from 7 lakes, 30 phytocoenoses of *N. candida* from 11 lakes and 17 phytocoenoses of *N. lutea* from 16 lakes. From a phytosociological point of view these phytocoenoses belong to the alliance Nymphaeion within the class Potametea [16].

In each phytocoenose one phytosociological relevé (10-25 m² plots) was recorded and water depth was measured in the deepest and the shallowest point of each phytocoenose taking into account the slope of the lake bottom. The abundance of each species within the relevé area was counted according to a modified Braun-Blanquet scale in which "+" = a few scattered specimens, mean coverage 0.1%; 1 = 1-10% coverage, mean 5%; 2 = 10-25% coverage, mean 17.5%; 3 = 25-50% coverage, mean 37.5%; 4 = 50-75% coverage, mean 62.5%; 5 = 75-100% coverage, mean 87.5%. All the relevés recorded in the investigated phytocoenoses were aggregated in one synoptic phytosociological table. In addition the constancy of each species was counted according to the following scale: I = species present in 1-20% of relevés; II = 21-40%; III = 41-60%; IV = 61-80%; V = 81-100%.

In each phytocoenose one water and one substrate sample were taken for physical and chemical analyses (79 water and 79 substrate samples together). Both types of samples were collected at the same time as the relevés were performed. Water samples were taken from an intermediate depth at which a given phytocoenose occurred and transferred to two 1000 ml plastic containers. Half of the water samples were preserved by the addition of 1 ml concentrated H₂SO₄ for PO₄³⁻, NO₃⁻ and total Fe analyses. Substrate samples were taken from the rhizome-root layer with a tubular bottom sampler and transferred to plastic bags. Each sample was a mixture of three random subsamples taken within a phytocoenose. Both water and substrate samples were transported to the laboratory within several hours (3-5) and the determinations were carried out immediately. Prior the analyses the samples were filtered and stored in refrigerators at 4°C until all the analyses were performed. Altogether 14 properties of water (including water depth) and 12 properties of substrate were determined.

The following analyses were carried out in water samples: pH, total and carbonate hardness, PO₄³⁻, NO₃⁻, total Fe, dissolved SiO₂, SO₄²⁻, Cl⁻, Mg²⁺, Ca²⁺, K⁺ and Na⁺.

In the substrate samples, pH was assessed in pore water. Hydration was determined by drying substrate samples at 105°C to constant weight and organic matter content was assessed by ignition of dry samples at 550°C for 1.5 h. Concentrations of dissolved SiO₂, NO₃⁻ and Cl⁻ were determined from water extracts, total Fe, Ca²⁺, Mg²⁺, PO₄³⁻ and SO₄²⁻ from (1:1 v/v) HCl extracts (see [17]). The concentrations of K⁺ and Na⁺ in substrate were determined from dried and ignited samples. Most of the determinations were carried out by methods described by Hermanowicz et al. [18]. Determinations of total hardness were performed according to procedures described by Hermanowicz et al. [18] and Hach Company [19].

The vegetation-habitat relationships were determined by means of Canonical Correspondence Analysis (CCA) using CANOCO for Windows Version 4.0 [20]. Water and substrate data were analysed separately. Moreover the habitat conditions of the four types of phytocoenoses investigated were compared with respect to each property using quartiles. The significance of differences in water and substrate properties between the

plant communities studied were determined by the Kruskal-Wallis test and the nonparametric multiple comparisons [21].

Results

Phytosociological characteristics of the phytocoenoses studied and their range of depth in the lake littoral

The analysis of Tab. 1 showed that the phytocoenoses studied formed dense stands and were characterized by a strong dominance of one species. The patches of *H. verticillata* were highly represented by species from the class Potametea, e.g. *N. lutea*, *Myriophyllum verticillatum*, *Ceratophyllum demersum*

Tab. 1 Synoptic table of the four types of phytocoenoses studied.

Phytocoenose type	HV	NC	NP	NL
Numer of relevés	14	30	18	17
Number of lakes	7	11	5	16
Potametea				
<i>Hydrilla verticillata</i>	V ⁴⁻⁵			
<i>Nymphaea candida</i>		V ³⁻⁵		I ⁺
<i>Nuphar pumila</i>			V ³⁻⁵	
<i>Nuphar lutea</i>	III ⁺	III ⁺¹	V ⁺¹	V ⁴⁻⁵
<i>Nymphaea alba</i>				I ⁺
<i>Potamogeton natans</i>		III ⁺²	II ⁺¹	II ⁺¹
<i>Myriophyllum verticillatum</i>	III ⁺²	I ⁺¹		I ⁺
<i>Ceratophyllum demersum</i>	III ⁺¹	II ⁺²		I ⁺
<i>Stratiotes aloides</i>	II ⁺¹		II ⁺¹	I ⁺
<i>Utricularia vulgaris</i>	II ⁺	I ⁺²		I ⁻¹
<i>Myriophyllum spicatum</i>	I ¹	I ⁺	I ⁺	
<i>Potamogeton lucens</i>	I ⁺	I ⁺	II ⁺¹	I ⁺
<i>Potamogeton compressus</i>	I ⁺		I ⁺	I ⁺
<i>Potamogeton friesii</i>		I ⁺	I+	
<i>Batrachium circinatum</i>		I ⁺		I ⁺
<i>Elodea canadensis</i>		I ⁺	III ⁺²	I ⁺
<i>Hydrocharis morsus-ranae</i>		I ⁺		
<i>Potamogeton perfoliatus</i>		I ⁺		I ⁺
<i>Potamogeton crispus</i>		I ⁺		
Others				
<i>Equisetum fluviatile</i>	I ⁺	I ⁺	III ⁺²	I ⁺
<i>Sagittaria sagittifolia</i>	I+			
<i>Lemna trisulca</i>		II ⁺²	II ⁺	I ⁺
<i>Spirodela polyrhiza</i>		I ⁺³	I ⁺	I ⁺
<i>Lemna minor</i>		I ⁺¹	I ⁺	I ⁺
<i>Fontinalis antipyretica</i>		I ⁺	I ⁺	I ⁺
<i>Phragmites australis</i>		I ⁺	I ⁺	II ⁺
<i>Schoenoplectus lacustris</i>		I ⁺		I ⁺
<i>Typha angustifolia</i>		I ⁺		I ⁺
<i>Sparganium emersum</i>		I ⁺	I ⁺	
<i>Chara tomentosa</i>		I ²	I ⁺	
<i>Chara</i> sp.			I ⁺	I ⁺
<i>Typha latifolia</i>				I ⁺

I, II... V – constancy; +, 1, 2... 5 – abundance; HV – *Hydrilla verticillata*; NC – *Nymphaea candida*; NP – *Nuphar pumila*; NL – *N. lutea*.

(constancy class III), *Stratiotes aloides* and *Utricularia vulgaris* (constancy class II). In the phytocoenoses of *N. candida* there was a high contribution of floating-leaved species such as *N. lutea*, *Potamogeton natans* (constancy class III) and *C. demersum* (constancy class II). The most frequent accompanying species was *Lemna trisulca* (constancy class II). The patches with *N. pumila* were characterized by a high proportion of *N. lutea* (constancy class V). Other frequently occurring species of the class Potametea included: *Elodea canadensis* (constancy class III), *Potamogeton lucens*, *P. natans* and *S. aloides* (constancy class II). Frequent accompanying species were: *Equisetum fluviatile* (constancy class III) and *L. trisulca* (constancy class II).

The analysis of the phytocoenoses of *N. lutea* showed that in addition to *N. lutea*, which dominated in the patches, *P. natans* and, among accompanying species, *Phragmites australis* (constancy class II) also played an important role in the community structure. It should be noted that among the dominant species *N. lutea* was the only species which occurred with high constancy in all the (four) types of phytocoenoses compared in the study.

The phytocoenoses studied developed at the following depths: *H. verticillata* (0.6-3.5 m; mainly 1.5-1.75 m), *N. pumila* (0.6-1.8 m; mainly 1.1-1.3 m), *N. candida* (0.1-2.8 m; mainly 1.5-1.7 m), *N. lutea* (0.4-2.0 m; mainly 0.9-1.3 m).

Water properties

The analysis of Fig. 1 showed that among the properties of water analysed the factors associated mainly with the carbonate complex, such as total hardness, SO_4^{2-} , K^+ , carbonate hardness, Mg^{2+} , Na^+ , Cl^- , Ca^{2+} and dissolved SiO_2 , as well as total Fe and PO_4^{3-} were most important in differentiating the waters of the vegetation types studied. These properties were correlated with the first ordination axis, which clearly separated the waters of *N. pumila* (left side of the diagram) from those of *N. lutea* (right side). The habitats of *N. candida* and *H. verticillata* phytocoenoses occupied an intermediate position.

The properties related to water hardness and correlated with the first ordination axis had significantly lower values in the waters of *N. pumila* than in those of *N. candida* (except for Ca^{2+}) and *N. lutea* (Fig. 2, Tab. 2). Compared to *H. verticillata*, the waters of *N. pumila* contained lower levels of total and carbonate hardness, dissolved SiO_2 and Mg^{2+} (Tab. 2). With regard to these properties the smallest differences were detected between the waters of *N. candida* and *N. lutea*, as well as between *N. candida* and *H. verticillata* (Fig. 2, Tab. 2). The waters of *N. lutea*, which are distinctly different from those of *N. pumila*, can be distinguished from the habitat of *H. verticillata* by higher total hardness, Ca^{2+} , Na^+ , K^+ , SO_4^{2-} and Cl^- concentrations (Fig. 2, Tab. 2). A reverse trend was demonstrated for total Fe and PO_4^{3-} , which were also correlated with the first ordination axis. The waters of *N. pumila* were characterized by the highest values of these two parameters (significant differences between *N. pumila* and the other three types of phytocoenoses were found – Tab. 2). The waters of *H. verticillata* were also richer in total Fe than those of *N. lutea* (the differences between *H. verticillata* and *N. lutea* were significant – Tab. 2).

From Fig. 1 it is clear that such properties as NO_3^- and mean water depth also played an important role in differentiating the waters of the phytocoenoses studied. These properties were negatively related to each other, but were both correlated with the second ordination axis, which clearly separated the waters of *N. pumila* and *N. lutea* (the upper part of the diagram) from

those of *N. candida* and *H. verticillata* (the lower part of the diagram). The waters inhabited by *N. pumila* and *N. lutea* were rich in NO_3^- , whereas those of *N. candida* and *H. verticillata* were distinguished by lower values of this parameter (see Tab. 2 for significant differences). In the case of mean water depth, the phytocoenoses of *N. candida* and *H. verticillata* developed in deeper waters, whereas those of *N. pumila* and *N. lutea* were associated with shallower parts of the lakes. Although significant differences were found only between the *N. candida* and *N. lutea* phytocoenoses, the observed trends in water depth were clearly marked (Fig. 2).

Among the 14 properties of water analysed only pH did not significantly differentiate the waters of the phytocoenoses studied.




Fig. 1 CCA ordination diagram of 1st and 2nd axes obtained for water properties and relevés of four plant communities, dominated by: NC – *Nymphaea candida*; HV – *Hydrilla verticillata*; NP – *Nuphar pumila*; NL – *Nuphar lutea*. Eigenvalues: $\lambda_1 = 0.787$; $\lambda_2 = 0.595$; $\lambda_3 = 0.216$; $\lambda_4 = 0.108$. Monte Carlo permutation test of significance of all canonical axes: $p = 0.002$.

Substrate properties

The substrate properties which best differentiated the habitats of the plant communities studied were pH, Ca^{2+} , Mg^{2+} and Na^+ , which were correlated with the first ordination axis (Fig. 3). As in the case of water properties, the axis clearly separated the substrates of *N. pumila* from those of *N. lutea*, whereas the habitats of *N. candida* and *H. verticillata* occupied an intermediate position. Total Fe, which was also correlated with the first ordination axis but negatively related to the five properties mentioned above (Fig. 3), appeared to play a significant role in differentiating the substrates of the plant communities studied as well.

The lowest values of pH, Ca^{2+} , Na^+ and Mg^{2+} were detected in the substrates of *N. pumila* (Fig. 4; significant differences between *N. pumila* and the other types of phytocoenoses were found regarding pH, Ca^{2+} and Na^+ , and between *N. pumila* and *N. lutea* with regard to Mg^{2+} – Tab. 2). High values of pH, Mg^{2+} , Na^+ and Ca^{2+} were recorded in the case of substrates of *N. lutea* (Fig. 4), which differed significantly from those of *N.*




Fig. 2 Differentiation of the four communities studied in relation to water properties. Black boxes show 25–75% interquartile ranges of values and white boxes show the medians. NC – plant communities dominated by *Nymphaea candida*; HV – by *Hydrilla verticillata*, NP – by *Nuphar pumila*, NL – by *Nuphar lutea*.

pumila and (except for Na⁺) *N. candida* (Tab. 2). It should be noted that no significant differences were detected between the substrates inhabited by *N. candida* and those of *H. verticillata*, whereas only slight differences were noted between the substrates of *H. verticillata* and *N. lutea* (Mg²⁺ content). As in the case of water, the substrates of *N. pumila* were distinguished from those of the other types of phytocoenoses by significantly higher values of total Fe (Fig. 3, Fig. 4, Tab. 2).

Fig. 3 shows that such properties as PO₄³⁻, Cl⁻ and, to a smaller degree, dissolved SiO₂ also played an important role in differentiating the substrates of the phytocoenoses studied. These properties were correlated with the second ordination axis, which clearly separated the substrates inhabited by *N. pumila* and *N. lutea* from those of *N. candida*

and *H. verticillata*. The substrates of *N. pumila* and *N. lutea* were generally poorer in PO₄³⁻ but richer in Cl⁻ (significance of differences – see Tab. 2). It is interesting to note that the highest content of NO₃⁻ was detected in the substrates of *N. pumila* (the differences between *N. pumila* and the other types of phytocoenoses were significant). Such properties as organic matter, K⁺ and SO₄²⁻ did not significantly differentiate the substrates of the plant communities studied.

Discussion

The comparative analysis showed that in north-eastern Poland the three rare water plant communities examined in

Tab. 2 Significance of differences in water and substrate properties between the four types of phytocoenoses studied.

Properties of water	NP-NC	NP-HV	NP-NL	NC-HV	NC-NL	HV-NL
Depth	ns	ns	ns	ns	+	ns
pH	ns	ns	ns	ns	ns	ns
Ca ²⁺	ns	ns	+	ns	+	+
Mg ²⁺	+	+	+	+	ns	ns
Total hardness	+	+	+	ns	ns	+
Carbonate hardness	+	+	+	ns	ns	ns
Na ⁺	+	ns	+	+	ns	+
K ⁺	+	ns	+	+	ns	+
Cl ⁻	+	ns	+	+	+	+
SO ₄ ²⁻	+	ns	+	ns	ns	+
SiO ₂ dissolved	+	+	+	ns	ns	ns
Total Fe	+	+	+	ns	ns	+
PO ₄ ³⁻	+	+	+	ns	ns	ns
NO ₃ ⁻	+	+	ns	ns	+	ns

Properties of substrate	NP-NC	NP-HV	NP-NL	NC-HV	NC-NL	HV-NL
Organic matter	ns	ns	ns	ns	ns	ns
K ⁺	ns	ns	ns	ns	ns	ns
SO ₄ ²⁻	ns	ns	ns	ns	ns	ns
SiO ₂ dissolved	+	ns	ns	ns	ns	ns
Cl ⁻	+	+	ns	ns	ns	+
Ca ²⁺	+	+	+	ns	+	ns
Mg ²⁺	ns	ns	+	ns	+	+
Na ⁺	+	+	+	ns	ns	ns
Total Fe	+	+	+	ns	ns	ns
pH	+	+	+	ns	+	ns
NO ₃ ⁻	+	+	+	ns	ns	ns
PO ₄ ³⁻	ns	ns	ns	ns	+	+

Symbols of phytocoenoses – see Tab. 1. “+” – significant at $p < 0.05$; ns: non-significant.

this study were distinct with respect to water and substrate chemistry preferences. The phytocoenoses of *N. pumila* were found to be ecologically most distinct among the phytocoenoses compared (acidic substrates, rich in total Fe and NO₃⁻ and soft waters, containing greater amounts of total Fe, PO₄³⁻ and NO₃⁻).

The relationship of *N. pumila* to soft waters was indicated by various authors (e.g. [8,9,11,22-24]). The specific character of the substrates (mostly organic and frequently peat bottoms) inhabited massively by this species was also emphasized [8,9,11,23,25]. Moreover the findings of some authors [9,26] showed that the phytocoenoses formed by *N. pumila* were ecologically distinct from those of *N. lutea* and *N. candida*. The results of this study demonstrated the distinct character of each of the habitats of *N. pumila* and *H. verticillata*.

The habitats of the phytocoenoses dominated by *H. verticillata* were also distinct from those of *N. lutea*, especially with regard to properties of water (lower values of most of the parameters analysed). Their substrates contained lower levels of Mg²⁺ and Cl⁻, and greater amounts of PO₄³⁻ compared to those of *N. lutea*, which may be ascribed to a higher proportion of well decomposed organic matter in the substrates of

H. verticillata.

The ecological distinctness of the *H. verticillata* phytocoenoses in relation to the patches of *Elodea canadensis* was already indicated earlier [10]. Moreover recent study showed the distinct character of the habitats of the phytocoenoses with *H. verticillata* in relation to some submerged plant communities of the alliance Potamion [27], especially with regard to waters which were poorer in Na⁺, K⁺ and Cl⁻. The data obtained by other authors revealed that *H. verticillata* had a wide ecological amplitude with respect to properties of water [4,22,28] and the species was very tolerant to organic pollution [29]. The persistence of *H. verticillata* in north-eastern Poland in dense, often large phytocoenoses and the establishment of new stands of the species may indicate that *H. verticillata* finds optimal growth conditions in this region of Poland.

The presented data referring to all the known localities of *N. candida* in north-eastern Poland confirmed the ecological distinctness of the phytocoenoses dominated by this species in relation to the phytocoenoses of *N. pumila* [9] and *N. lutea* [9,30,31]. It is interesting to note that the habitats of *N. candida* are very similar to those of *H. verticillata*, especially with regard to substrate properties (no significant differences between these two communities were found). These two types of phytocoenoses inhabit more frequently organic substrates than other types of phytocoenoses. These findings confirm the results of several other studies concerning *N. candida* [31-33]. Earlier data indicated that *N. candida* phytocoenoses were associated mostly with mesotrophic waters [9,34]. The supplemented wider study conducted in north-eastern Poland shows that at present the phytocoenoses of *N. candida* inhabit more fertile eutrophic waters, as evidenced by a wide range of PO₄³⁻ concentration in water. It appears that both *N. candida*




Fig. 3 CCA ordination diagram of 1st and 2nd axes obtained for substrate properties and relevés of four plant communities, dominated by: NC – *Nymphaea candida*; HV – *Hydrilla verticillata*; NP – *Nuphar pumila*; NL – *Nuphar lutea*. Eigenvalues: $\lambda_1 = 0.628$; $\lambda_2 = 0.472$; $\lambda_3 = 0.153$; $\lambda_4 = 0.029$. Monte Carlo permutation test of significance of all canonical axes: $p = 0.002$.

and *H. verticillata* attain their optimum development in lakes of north-eastern Poland.

The following question should also be considered: What might happen to the three types of phytocoenoses studied in the future? Recent studies conducted in several tens of lakes within north-eastern Poland (2002-2005) indicate that twenty-five years after the first habitat investigations (1978-1980) there has been a slight increase in the trophic level of the lakes [27]. In addition alkalization proceeds gradually and an increase in water hardness is observed. It appears, therefore, that the phytocoenoses of *N. pumila* are among the most threatened communities in the lakes. The increase in water trophy does not seem to affect these phytocoenoses as *N. pumila* tolerates a wide range of trophic conditions [8], which was also indicated by some other authors [23,25]. However, changes in water hardness appear to play an important role. The phytocoenoses of *N. pumila* are associated mostly with soft waters. The data obtained hitherto from Poland [8,9,35] indicate that the stands of this species in north-eastern Poland are established outside of their ecological optimum, which are very soft waters, i.e. they are rather closer to the tolerance limits of the species for water hardness. It may be assumed, therefore, that the

phytocoenoses of *N. pumila* will retreat from the lakes even when there is a slight increase in water hardness. By contrast, the *N. candida* and *H. verticillata* phytocoenoses, which inhabit waters with a wider range of hardness (slightly soft to moderately hard waters) and tolerate a moderate increase in trophic level, will probably continue to persist in lakes of north-eastern Poland for a long time provided that they are not subject to stronger anthropogenic impact.

References

1. Tomaszewicz H, Kłosowski S. Roślinność wodna i szuwarowa jezior Pojezierza Sejneńskiego. Monogr Bot. 1985;67:69-141.
2. Tomaszewicz H. Fitocenozy z udziałem *Hydrilla verticillata* (L. fil.) Casp. na Pojezierzu Suwalskim i ich stanowisko syntaksonomiczne. Fragn Flor Geobot. 1976;22:347-350.
3. Czapik R, Tomaszewicz H. Przesiąkora okółkowa *Hydrilla verticillata* kwitnie na Suwalszczyźnie. Chronimy Przyr Ojcz. 1979;35(4):26-36.
4. Pietsch W. Zur Bioindikation *Najas marina* L. s.l. – und




Fig. 4 Differentiation of the four communities studied in relation to substrate properties. Black boxes show 25-75% interquartile ranges of values and white boxes show the medians. NC – plant communities dominated by *Nymphaea candida*; HV – by *Hydrilla verticillata*; NP – by *Nuphar pumila*; NL – by *Nuphar lutea*.

- Hydrilla verticillata* (L. fil.) Royle-reicher Gewässer Mitteleuropas. Feddes Repert. 1981;92:125-174.
5. Podbielkowski Z, Tomaszewicz H. Rzadkie zbiorowiska roślinne Pojezierza Suwalskiego. Rocznik Białostocki. 1981;15:193-209.
 6. Kłosowski S. *Nuphar pumila* (Timm.) DC. Grąźel drobny. In: Kaźmierczakowa R, Zarzycki K, editors. Polska czerwona księga roślin. Paprotniki i rośliny kwiatowe. Kraków: W. Szafer Institute of Botany, Polish Academy of Sciences; 2001. p. 116-118.
 7. Kłosowski S, Tomaszewicz H. Habitat conditions of the *Nymphaeetum candidae* Miljan 1958 and *Nupharo-Nymphaeetum albae* Tomaszewicz 1977 dominated by *Nymphaea alba*. Acta Soc Bot Pol. 1989;58:613-624.
 8. Kłosowski S, Tomaszewicz H. Standortverhältnisse des *Nupharetum pumili* Oberdorfer 1957 in der Suwałki-Seenplatte. Arch Hydrobiol. 1990;117:365-382.
 9. Kłosowski S, Tomaszewicz H. Standortverhältnisse der Gesellschaften mit dominant einzelner Nymphaeaceen in Nordost-Polen. Tuexenia. 1993;13:75-90.
 10. Kłosowski S, Tomaszewicz H. Zur Sozologie und Ökologie des *Hydrilietum verticillatae* Tomaszewicz 1979 und des *Elodeetum canadensis* (Pign. 1953) Pass. 1964 in Nordost-Polen. Tuexenia. 1997;17:125-136.
 11. Kłosowski S, Krężelewska I, Tomaszewicz H. Habitat conditions of the phytocoenoses of *Nupharetum pumili* in Poland. Fragm Flor Geobot. 1996;41:707-715.
 12. Pieczyńska E. Ecology of the eulittoral zone of lakes. Ekol Pol. 1972;20:637-732.
 13. Planter M. Physical and chemical conditions in the helophytes zone of lakes. Arch Hydrobiol. 1973;20:1-7.
 14. Zachwieja J. Wahania dobowe temperatury, O₂, CO₂, pH i alkaliczności w litoralu jeziora Mamry. Pol Arch Hydrobiol. 1965;13:5-27.
 15. Kłosowski S. Temporal and spatial variation of habitat conditions in the zonation of littoral plant communities. Aquat Bot. 1992;43(2):199-208. [http://dx.doi.org/10.1016/0304-3770\(92\)90043-I](http://dx.doi.org/10.1016/0304-3770(92)90043-I)
 16. Matuszkiewicz W. Przewodnik do oznaczania zbiorowisk roślinnych Polski. 3rd ed. Warszawa: Polish Scientific Publishers PWN; 2001.
 17. Kłosowski S. The relationships between environmental factors and the submerged Potamogeton associations in lakes of north-eastern Poland. Hydrobiologia. 2006;560(1):15-29. <http://dx.doi.org/10.1007/s10750-005-1141-1>
 18. Hermanowicz W, Dożańska, Dojlido JR, Koziorowski B. Fizyczno-chemiczne badanie wody i scieków. Warszawa: Arkady; 1999.
 19. Hach Company. Hach water analysis handbook. Loveland CO: Hach Company; 1992.
 20. ter Braak CJF, Šmilauer P. CANOCO reference manual and user's guide to Canoco for Windows: software for canonical community ordination (version 4). Ithaca NY: Microcomputer Power; 1998.
 21. Zar JH. Biostatistical analysis. Englewood Cliffs NJ: Prentice Hall; 1984.
 22. Kadono Y. Occurrence of aquatic macrophytes in relation to pH, alkalinity, Ca²⁺, Cl⁻ and conductivity. Jap J Ecol. 1982;32:39-44.
 23. Roweck H. Ökologische Untersuchungen an Teichrosen. Arch Hydrobiol. 1988;81 suppl:103-358.
 24. Doll R. Die Pflanzengesellschaften der stehenden Gewässer in Mecklenburg-Vorpommern Teil I. 3. Potamogetonetea Tx. et Prsg. 42 – Laichkrautgesellschaften. Feddes Repert. 1991;102(3-4):217-317. <http://dx.doi.org/10.1002/fedr.19911020310>
 25. Kozłowski G, Eggenberg S. Vorkommen der Kleinen Teichrose *Nuphar pumila* und des Hybrids *N. × intermedia* in der Schweiz. Bot Helv. 2005;115:125-136.
 26. Mäkelä S, Huitu E, Arvola L. Spatial patterns in aquatic vegetation composition and environmental covariates along chains of lakes in the Kokemäenjoki watershed (S. Finland). Aquat Bot. 2004;80(4):253-269. <http://dx.doi.org/10.1016/j.aquabot.2004.08.006>
 27. Kłosowski S, Tomaszewicz GH, Tomaszewicz H. The expansion and decline of charophyte communities in lakes within the Sejny Lake District (north-eastern Poland) and changes in water chemistry. Limnologica. 2006;36(4):234-240. <http://dx.doi.org/10.1016/j.limno.2006.07.001>
 28. Kunii H, Kunii K, Aso K, Sakata K. Records of aquatic macrophyte flora and environmental factors from the irrigation ponds around Lake Shinji, Shimane, Japan. Mem Fac Sci Shimane Univ. 1991;25:85-96.
 29. Cook CDK, Lüönd R. A revision of the genus *Hydrilla* (Hydrocharitaceae). Aquat Bot. 1982;13(0):485-504. [http://dx.doi.org/10.1016/0304-3770\(82\)90074-2](http://dx.doi.org/10.1016/0304-3770(82)90074-2)
 30. Pietsch W. Makrophytische Indikatoren für ökochemische Beschaffenheit der Gewässer. In: Breitig G, Tumpling W, editors. Ausgewählte Methoden der Wasseruntersuchung. Jena: Gustav Fischer Verlag; 1982. p. 67-88. (vol 2).
 31. van der Velde G, Cysters CPC, de Lyon MJH. The distribution of four nymphaeid species in the Netherlands in relation to selected abiotic factors. Proceedings EWRS/AAB 7th Symposium on Aquatic Weeds; Loughborough. 1986. p. 363-368.
 32. de Lyon MJH, Roelofs JGM. Waterplanten in relatie tot waterkwaliteit en bodemgesteldheid. Nijmegen: Katholieke Universiteit, Laboratorium voor Aquatische Oecologie; 1986. (pt 1).
 33. de Lyon MJH, Roelofs JGM. Waterplanten in relatie tot waterkwaliteit en bodemgesteldheid. Nijmegen: Katholieke Universiteit, Laboratorium voor Aquatische Oecologie; 1986. (pt 2).
 34. Szańkowski M, Kłosowski S. Habitat conditions of nymphaeid associations in Poland. Hydrobiologia. 1999;415:177-185. <http://dx.doi.org/10.1023/A:1003837520267>
 35. Kłosowski S. Ökologische Differenzierung innerhalb von Wasserpflanzen Gesellschaften am Beispiel des *Myriophylletum alterniflori*, des *Nupharetum pumili* und des *Polygonetum natantis*. Fragm Flor Geobot. 1994;39:571-587.