

Leray-Serre Spectral Sequence for Quasi-Fibrations

Vicente Muñoz

Departamento de Matemáticas, Facultad de Ciencias

Universidad Autónoma de Madrid

28049 Madrid, Spain

e-mail address: vicente.munoz@uam.es

URL address: <http://www.adi.uam.es/~vmunoz>

ABSTRACT. Let $f : X \rightarrow Y$ be a map between two spaces for which Y is stratified such that f is a fibration over each open stratum. We find some spectral sequences to compute the homology of X in terms of the homology of Y and that of the fibers. We apply this to give a Lefschetz theorem for the degeneracy loci of a morphism between holomorphic bundles on a complex manifold.

1 Homotopy and Homology

To understand the topology of a space X , one of the primary issues is the computation of the homotopy groups $\pi_n(X)$. Even in the simplest examples of spaces, homotopy groups turn out to be very difficult to compute. On the other hand, simplicial homology $H_n(X)$ is a good alternative being easy to compute due to the existence of a Mayer-Vietoris principle. This allows to divide the space X and "glue" the homology of the pieces. The good news

2000 *Mathematics Subject Classification.* 55N10, 32Q55.

Key words and phrases. singular homology, homotopy, spectral sequence, Lefschetz theorem, degeneracy loci.

Partially supported by The European Contract Human Potential Programme, Research Training Network HPRN-CT-2000-00101.

is that Hurewicz theorem relates both up to some extent, which may help us to compute homotopy groups.

Let us first review the classical notions about homotopy from [Sp]. Consider a topological space X with a fixed base point $\ast \in X$. Then $\pi_n(X)$ consists of equivalence classes of maps $\alpha : S^n \rightarrow X$ from the n -sphere sending a base point $x_0 \in S^n$ to \ast , modulo the equivalence relation \simeq given by $\alpha_0 \simeq \alpha_1$ if there is a homotopy $H : S^n \times [0, 1] \rightarrow X$ with $H(x, 0) = \alpha_0(x)$, $H(x, 1) = \alpha_1(x)$, for all $x \in S^n$ and $H(x_0, t) = \ast$, for all $t \in [0, 1]$. A space X is said to be n -connected if $\pi_i(X) = 0$ for $i \leq n$. For the special case $n = 0$, we have that $\pi_0(X) = 0$ if X is arc-wise connected. Also $\pi_1(X)$ is the fundamental group of X , so that X is 1-connected means that it is simply-connected.

There is a relative version of the homotopy groups: if $A \subset X$ is a subspace containing the base point, then $\pi_n(X, A)$ consists of equivalence classes of maps $\alpha : (D^n, \partial D^n) \rightarrow (X, A)$, this meaning that α is a map from the disk D^n to X sending the boundary to A . The equivalence relation is through homotopies that send the boundaries into A .

For notions in relation with homology one can look at [Ma]. Let X be a topological space. Let $I = [0, 1]$ be the unit interval. A n -cube is a continuous map $T : I^n \rightarrow X$, and it is degenerate if it is independent of one of its variables. The complex of singular chains in X is defined as the free abelian group $C_n(X)$ generated by the n -cubes, modulo the degenerate ones. The boundary operator ∂ associates to every n -cube the $(n-1)$ -chain which consists in the sum of its faces with an appropriate sign given by its orientation. The homology $H_*(X)$ of X is given by the homology of $(C_*(X), \partial)$.

For the relative version, let $A \subset X$. Then the complex $C_*(X, A)$ is the quotient complex $C_*(X)/C_*(A)$ with the induced boundary operator. The homology of this complex is the relative homology $H_*(X, A)$.

There is a morphism, the Hurewicz homomorphism,

$$\pi_n(X, A) \longrightarrow H_n(X, A),$$

which sends the class of the map $\alpha : (D^n, \partial D^n) \rightarrow (X, A)$ to the cycle $\alpha \in C_n(X, A)$ using the homeomorphism $I^n \cong D^n$ and since $\partial\alpha \in C_n(A)$. For the following theorem see [Sp, §7].

Hurewicz theorem. *Let (X, A) be a pair of 1-connected spaces. Then $\pi_i(X, A) = 0$ for $i \leq n-1$ implies that $\pi_n(X, A) \cong H_n(X, A)$. In plain terms, all the homotopy and homology groups are zero up to the first non-zero ones, which are isomorphic.*

This means that (X, A) is n -connected, i.e., $\pi_i(A) \cong \pi_i(X)$, for $i < n$, and $\pi_n(A) \rightarrow \pi_n(X)$. If the spaces A and X are CW-complexes then, up to homotopy equivalence, X is constructed out of A by attaching cells of dimension $n + 1$ and over.

2 Spectral Sequence for a Filtration

Suppose that X is a space and $A \subset X$, then we have the long exact sequence of the pair

$$\cdots \rightarrow H_n(A) \rightarrow H_n(X) \rightarrow H_n(X, A) \xrightarrow{\partial_*} H_{n-1}(A) \rightarrow \cdots$$

This comes from the exact sequence of chain complexes $C_*(A) \rightarrow C_*(X) \rightarrow C_*(X, A)$. The connecting homomorphism ∂_* is defined as follows. Let $[z] \in H_n(X, A)$ with a representative $z \in C_n(X, A)$. Lift z to $\tilde{z} \in C_n(X)$, then $\partial\tilde{z} \in C_n(A)$ and set $\partial_*[z] = [\partial\tilde{z}]$.

We may think of this case as a two terms filtration $X_0 = A \subset X_1 = X$. What happens when we have a longer filtration $X_0 \subset X_1 \subset \cdots \subset X_d = X$? Then $C_*(X)$ is a filtered complex with filtration given by $C_*(X_q) \subset C_*(X)$. This gives rise to a *spectral sequence*. A spectral sequence is a sequence of double complexes (E_{pq}^r, d^r) with differentials $d^r : E_{pq}^r \rightarrow E_{p+r-1, q-r}^r$ such that the homology of the r -th term is the following one $H(E^r, d^r) = E^{r+1}$. We say that it converges if for each fixed (p, q) the terms E_{pq}^r stabilizes for r large enough. The limiting one is denoted as E_{pq}^∞ . For specific material on spectral sequences, the reader is recommended the nice book [Mc] or [BT, §14].

In our case we generate a spectral sequence with

$$E_{pq}^0 = C_{p+q}(X_q)/C_{p+q}(X_{q-1}) = C_{p+q}(X_q, X_{q-1}).$$

This has an induced differential d^0 which is the standard boundary in chains. The homology is

$$E_{pq}^1 = H_{p+q}(X_q, X_{q-1}).$$


The general machinery on spectral sequences tells us that the differential

$$d^1 : H_{p+q}(X_q, X_{q-1}) \rightarrow H_{p+q-1}(X_{q-1}, X_{q-2})$$

is as follows: let $[z] \in H_{p+q}(X_q, X_{q-1})$. Select a representative $z \in C_{p+q}(X_q, X_{q-1})$ and lift it to $\tilde{z} \in C_{p+q}(X_q)$. Then consider the image of $\partial\tilde{z} \in C_{p+q-1}(X_{q-1})$ in $C_{p+q-1}(X_{q-1}, X_{q-2})$, and set $d_1^1[z] = [\partial\tilde{z}] \in H_{p+q-1}(X_{q-1}, X_{q-2})$. In particular, it follows that $d^1[z] = 0$ means that there exists $\tilde{z}_1 \in C_{p+q}(X_{q-1})$ such that $\partial\tilde{z} + \partial\tilde{z}_1 \in C_*(X_{q-2})$.

In general, a class $[z]$ survives to E^r , i.e., $d^1[z] = 0, \dots, d^{r-1}[z] = 0$ if there are $\tilde{z}_i \in$

$C_{p+q}(X_{q-1})$ such that $\partial(\tilde{z} + \tilde{z}_1 + \cdots + \tilde{z}_{r-1})$ belongs to $C_{p+q-1}(X_{q-r})$. The latter element defines $d^r[z] \in E^r$. Moreover, the class $[z]$ is zero in E^r if there are $\tilde{w}_i \in C_{p+q+1}(X_{q+i})$ with $\partial(\tilde{w}_1 + \tilde{w}_2 + \cdots + \tilde{w}_r) - \tilde{z} \in C_*(X_{q-1})$.



The E^∞ -term is the homology $H_*(X)$ in the following sense. There is a filtration of the homology $H_*(X)$ given by the subspaces $F_q H_*(X) = \text{im}(H_*(X_q) \rightarrow H_*(X))$. Then

$$E_{pq}^\infty = \frac{F_q H_{p+q}(X)}{F_{q-1} H_{p+q}(X)}.$$

If we work with homology over a field, e.g. the reals, we have $\bigoplus_{p+q=n} E_{pq}^\infty \otimes \mathbb{R} \cong H_n(X) \otimes \mathbb{R}$ as vector spaces.

Now we are going to look to the case of a pair (X, B) . For any $A \subset X$, we have an exact sequence

$$\cdots \rightarrow H_n(A, A \cap B) \rightarrow H_n(X, B) \rightarrow H_n(X, A \cup B) \rightarrow H_{n-1}(A, A \cap B) \rightarrow \cdots$$

This comes from the exact sequence of complexes

$$\frac{C_*(A)}{C_*(A \cap B)} \hookrightarrow \frac{C_*(X)}{C_*(B)} \twoheadrightarrow \frac{C_*(X)}{C_*(A) + C_*(B)}$$

For proving this we need that $C_*(A) + C_*(B) \subset C_*(A \cup B)$ induces isomorphism in homology [Ma, page 151]. This is termed as $\{A, B\}$ is an excisive couple, i.e., that the Mayer-Vietoris holds for $A \cup B$, A and B . This is true when the interiors of A and B cover $A \cup B$ in the relative topology of $A \cup B$. The proof consists on a process of subdivision of the cycles in $A \cup B$. But also it is true when A and B are CW-subcomplexes of a CW-complex X , since then there are open sets $U \supset A$, $V \supset B$ in $A \cap B$ such that U retracts to A , V retracts to B and $U \cap V$ retracts to $A \cap B$.

Theorem (spectral sequence for a filtration). *Let (X, B) be a pair of CW-complexes, $X_0 \subset X_1 \subset \dots \subset X_d = X$ a filtration by CW-subcomplexes. Then there is a spectral sequence with $E_{pq}^1 = H_{p+q}(X_q, X_{q-1} \cup (B \cap X_q))$ converging to $H_n(X, B)$.*

We look at the filtration given by

$$\frac{C_*(X_q)}{C_*(B \cap X_q)} \subset \frac{C_*(X)}{C_*(B)}.$$

The E^0 term is the cokernel in

$$\frac{C_*(X_{q-1})}{C_*(B \cap X_{q-1})} \hookrightarrow \frac{C_*(X_q)}{C_*(B \cap X_q)} \twoheadrightarrow \frac{C_*(X_q)}{C_*(X_{q-1}) + C_*(B \cap X_q)}.$$

Under the assumptions, the various couples are excisive, so we get a spectral sequence as before.

3 Spectral Sequences for Quasi-Fibrations

Let us now turn our attention to a different issue. Very typically spectral sequences are used for a fibration $f: E \rightarrow B$ with fiber F . Suppose that B is 1-connected. Then the homology of the different fibers are identified in a canonical way. Therefore $H_*(F)$ is a system of (constant!) coefficients over the base B . We have the following standard relationship between the homology of the base B and fiber F and that of the total space E (see [Mc] and [BT]).

Leray-Serre spectral sequence. *Suppose that B is 1-connected. There is a spectral sequence whose second term is $E_{pq}^2 = H_q(B; H_p(F))$ and converging to $H_n(E)$.*

We prove this result for CW-complexes. Take a CW-decomposition for B and consider the skeleta $B_n = B^{(n)}$. This is a filtration of the space B . (It may be an infinite filtration $B = \bigcup_{n \geq 1} B_n$, but the theory works because B has the weak topology with respect to skeleta). Now for any group G of coefficients, the homology $H_*(B; G)$ can be computed as the homology of the $H_*(B_n, B_{n-1}) \otimes G$. This is true since the spectral sequence $E_{pq}^1 = H_{p+q}(B_n, B_{n-1}; G)$ is only non-zero for $n = p + q = q$, i.e., for $p = 0$. Therefore the only non-trivial differential is $d^1: E_{pq}^1 \rightarrow E_{p, q-1}^1$. All the other ones have to vanish and hence $H_*(B; G) = H_*(E^1, d^1)$.

Consider now the following filtration $E_n = f^{-1}(B_n)$ of the total space E and the corresponding spectral sequence for the filtration. The fibration over each n -cell in (E_n, E_{n-1}) is

trivial. This means that the complexes $C_*(E_n, E_{n-1})$ and $C_*(B_n, B_{n-1}) \otimes C_*(F)$ are almost the same, in the sense that their homologies coincide, $H_*(E_n, E_{n-1}) = H_*(B_n, B_{n-1}) \otimes H_*(F)$, which is the E^1 -term of the spectral sequence of the filtration. The E^2 -term is the homology $H_*(B; H_*(F))$. Note that the first non-zero differential will be a map $d^r : H_q(B; H_p(F)) \rightarrow H_{q-r}(B; H_{p+r-1}(F))$.

Instead it is more common to have a map $f : X \rightarrow Y$ which is not a fibration but can be decomposed into pieces where it actually is. We want to extend the Leray-Serre spectral sequence to this case, relating the homology of X with that of Y .

Definition. A map $f : X \rightarrow Y$ between two spaces is called a *quasi-fibration* if there is a filtration Y_q of Y by closed subspaces such that denoting by $X_q = f^{-1}(Y_q)$ the corresponding filtration for X ,

1. the restriction of f to $X_q - X_{q-1}$, $f_q : X_q - X_{q-1} \rightarrow Y_q - Y_{q-1}$ is a fibration with fiber F_q , for every q .
2. For each q , there is an open neighborhood U of Y_{q-1} in Y_q that retracts to Y_{q-1} , such that $f^{-1}(U)$ is a neighborhood of X_{q-1} in X_q that retracts to X_{q-1} .

Before going on further, let us say some words on the hypothesis on f . The condition of U being a neighborhood of Y_{q-1} in Y_q retracting to Y_{q-1} is technically referred to as NDR=Neighbourhood Deformation Retract. It occurs very often, for instance when Y_{q-1} is a CW-subcomplex of a CW-complex Y_q (e.g. a submanifold, even singular, in a differentiable manifold).

If we are dealing with CW-complexes (the filtration given by CW-subcomplexes) and f is cellular, then condition (ii) above is satisfied assuming f is proper: Take a retracting neighborhood U of Y_{q-1} in Y_q . We consider the neighborhood $f^{-1}(U)$ of X_{q-1} and take a retracting neighborhood $V \subset f^{-1}(U)$ of X_{q-1} in X_q . Then since f is proper, f is closed, so there exists U' (in fact, $U' \subset Y - f(X - V)$) with $U' \subset U$ and $f^{-1}(U') \subset V$. Moreover we may assume (since X_q is a CW-complex) that U also retracts to \bar{U}' . Let $H : U \times [0, 1] \rightarrow U$ be the map with $H(y, 0) = y$, $H(y', t) = y'$ for $y' \in \bar{U}'$ and $H(y, 1) \in \bar{U}'$ for $y \in U$. Easily we may also assume that $H(y, t) \in U - Y_{q-1}$ for any $y \in U - Y_{q-1}$, $t \in [0, 1]$.

By the fibration property, we lift the retraction to $\tilde{H} : (f^{-1}(U) - X_{q-1}) \times [0, 1] \rightarrow f^{-1}(U) - X_{q-1}$ with $\tilde{H}(x, 0) = x$, and $\tilde{H}(x, 1) \in f^{-1}(\bar{U}')$. Now take a continuous function $u : U \rightarrow [0, 1]$ which is 0 in a neighborhood $W \subset U'$ of X_{q-1} and 1 in $f^{-1}(U - U')$. Then $K(x, t) = \tilde{H}(x, u(x)t)$ is a retraction of $f^{-1}(U) - X_{q-1}$ to $f^{-1}(\bar{U}') - X_{q-1}$ which is the

identity in $f^{-1}(W) - X_{q-1}$. Therefore it may be extended with $K(x, t) = x$ for $x \in X_{q-1}$. Follow this retraction with the retraction form V to X_{q-1} to get a retraction from $f^{-1}(U)$ to X_{q-1} as required.

Theorem. *Let $f : X \rightarrow Y$ be a quasi-fibration with all of the Y_q being 1-connected. Then for a fixed integer q , there is a spectral sequence whose E^2 term is $H_*(Y_q, Y_{q-1}; H_*(F_q))$ and converging to $H_*(X_q, X_{q-1})$.*

Under the assumptions we have

$$H_*(X_q, X_{q-1}) = H_*(X_q, f^{-1}(U)) = H_*(X_q - X_{q-1}, f^{-1}(U - Y_{q-1})).$$

the last equality by excision. Also $H_*(Y_q, Y_{q-1}) = H_*(Y_q, U) = H_*(Y_q - Y_{q-1}, U - Y_{q-1})$. Since

$$(X_q - X_{q-1}, f^{-1}(U - Y_{q-1})) \rightarrow (Y_q - Y_{q-1}, U - Y_{q-1})$$

is a fibration with fiber F_q we have the result from the Leray-Serre spectral sequence.

For pairs we have a similar result which we state in the context of CW-complexes.

Theorem. *Let $f : X \rightarrow Y$ be a proper map between CW-complexes and let (X, B) be a pair of CW-complexes. Let Y_q be a filtration of Y by CW-subcomplexes and let $X_q = f^{-1}(Y_q)$, $B_q = f^{-1}(Y_q) \cap B$ be the corresponding filtrations of X and B . Suppose that $(X_q - X_{q-1}, B_q - B_{q-1}) \rightarrow (Y_q - Y_{q-1})$ is a fibration (of a pair of spaces) with fiber (F_q, G_q) . Then for fixed q , there is a spectral sequence with $E^2 = H_*(Y_q, Y_{q-1}; H_*(F_q, G_q))$ and converging to $E^\infty = H_*(X_q, X_{q-1} \cup B_q)$.*

4 The Fundamental Group on a Quasi-Fibration

The Leray-Serre spectral sequence (at least in the simplified version we have stated it in section 3) leaves aside an important issue of the behaviour of the topology of spaces under a quasi-fibration, namely the question relative to the fundamental group. Let us deal with this by hand.

Theorem. *Let $f : X \rightarrow Y$ be a quasi-fibration between 0-connected spaces, such that the fibers F_q are connected. Then*

$$f_* : \pi_1(X) \rightarrow \pi_1(Y)$$

is surjective.

To see this, take a loop $\gamma : [0, 1] \rightarrow Y$. Consider a sub-interval $[0, t_1]$ such that $\gamma([0, t_1])$ is included in $Y_q - Y_{q-1}$ and $\gamma(t_1) \in Y_{q-1}$. This allows to write $\gamma = \gamma_1 * \gamma_2$ as a juxtaposition of $\gamma_1 = \gamma|_{[0, t_1]}$ and $\gamma_2 = \gamma|_{[t_1, 1]}$.

Now consider a neighborhood U of Y_{q-1} in Y_q that retracts to Y_{q-1} such that $f^{-1}(U)$ retracts to X_{q-1} . Choose some $\varepsilon > 0$ with $\gamma_1([t_1 - \varepsilon, t_1]) \subset U$. Then we use the fibration property for $X_q - X_{q-1} \rightarrow Y_q - Y_{q-1}$ to lift $\gamma_1|_{[0, t_1 - \varepsilon]}$ to a path $\tilde{\gamma}_1$. With the retraction, we may join $\tilde{\gamma}_1(t_1 - \varepsilon)$ with a point in X_{q-1} by a path $\tilde{\delta}$ in $f^{-1}(U)$. Let $\delta = f \circ \tilde{\delta}$ be the induced path. The path $\delta^{-1} * \gamma_1|_{[t_1 - \varepsilon, t_1]}$ is in U with end-points in Y_{q-1} so it is homotopic (via the retraction) to a path α in Y_{q-1} . This means that $\delta * \alpha \simeq \gamma_1|_{[t_1 - \varepsilon, t_1]}$. So we decompose

$$\gamma = ((\gamma_1|_{[0, t_1 - \varepsilon]}) * \delta) * (\alpha * \gamma_2).$$

The first loop is lifted to a path $\tilde{\gamma}_1 * \tilde{\delta}$ with end-point in X_{q-1} .

We perform the above construction simultaneously in every portion of γ in Y_q . Specifically, suppose $\gamma([0, 1]) \subset Y_q$. Write $\gamma = \gamma_1 * \dots * \gamma_r$ with $\gamma_i = \gamma|_{[t_{i-1}, t_i]}$ satisfying that either $\gamma_i([t_{i-1}, t_i]) \subset Y_{q-1}$ or $\gamma_i([t_{i-1}, t_i]) \subset Y_q - Y_{q-1}$ and $\gamma_i(\{t_{i-1}, t_i\}) \subset Y_{q-1}$ (except for possibly $\gamma_1(0)$ and $\gamma_r(1)$). Let $I \subset \{1, \dots, r\}$ be the set of indices of those γ_i satisfying the second condition.

The method above allows to change γ by $\gamma \simeq \gamma'_1 * \dots * \gamma'_r$ such that γ'_i is liftable to $\tilde{\gamma}'_i$ for $i \in I$, and γ'_i is in Y_{q-1} for $i \notin I$. Since γ'_i is included in Y_{q-1} , for $i \notin I$, by induction on q we may suppose that they are liftable to $\tilde{\gamma}'_i$. If the end points of consecutive $\tilde{\gamma}'_i$ do not match then we use that the fibers are arc-wise connected to join them.

Therefore there is a loop $\tilde{\gamma} = \tilde{\gamma}'_1 * \dots * \tilde{\gamma}'_r$ such that $f \circ \tilde{\gamma} \simeq \gamma$, as required to have surjectivity of f_* .

Obviously the condition on the fibers F_q being 0-connected is necessary. Just think of the projection

$$f : S^1 = \{(x, y) | x^2 + y^2 = 1\} \rightarrow [-1, 1] \\ (x, y) \mapsto x$$

which is a quasi-fibration, but not surjective in the fundamental groups.

5 Degeneracy Loci

An important point of a mathematical theory is to have applications which prove its usefulness. Our application touches the world of algebraic geometry. Start with a complex manifold M of (complex) dimension n . This means that M is covered by charts which are open sets of \mathbb{C}^n and the change of charts are bi-holomorphisms. A holomorphic vector bundle $E \rightarrow M$ of rank e is a complex vector bundle which admits local trivializations where the transition functions are holomorphic [We].

For two holomorphic vector bundles E and F over M , of ranks e and f respectively, let $\phi: E \rightarrow F$ be a holomorphic morphism. This means that ϕ is holomorphic as a section of $\text{Hom}(E, F)$. The r -degeneracy locus of ϕ is the subset

$$D_r(\phi) = \{x \in M \mid \text{rk } \phi(x) \leq r\}.$$

This is a complex submanifold of M but it is not smooth in general. It is usually said that $D_r(\phi)$ has a determinantal structure [ACGH], which forces $D_r(\phi)$ to be singular along $D_{r-1}(\phi)$.

The way to understand the degeneracy loci is the following. Construct the grassmannian bundle $G = \text{Gr}(e-r, E)$ over M . This consists of putting the grassmannian $\text{Gr}(e-r, E_x)$ of $(e-r)$ -subspaces of the fiber E_x over every $x \in M$ in a differentiable varying way. The canonical projection of this fibration to M will be denoted by π . There is a universal bundle U over G . Every point $V \in G$ is actually an $(e-r)$ -dimensional subspace $V \subset E_x$ for $x = \pi(V)$. So the bundle U has fiber over $V \in G$ the space V itself. Since $V \subset E_{\pi(V)}$, we get $U \subset \pi^*E$. Composing this inclusion with the map $\pi^*(\phi): \pi^*E \rightarrow \pi^*F$ we get a map $\phi_r: U \rightarrow \pi^*F$. Now ϕ_r is zero at a point $V \in G$ if and only if $\phi(V) = 0$. This means that at $x = \pi(V)$ there is an $(e-r)$ -dimensional subspace of E_x in the kernel of $\phi(x)$ and therefore $\text{rk}(\phi(x)) \leq r$. So denoting by $Z(\phi_r) \subset G$ the zero set of ϕ_r , we have

$$\pi(Z(\phi_r)) = D_r(\phi).$$

Moreover this map is one to one over $D_r(\phi) - D_{r-1}(\phi)$. We have an obvious filtration of $D_r(\phi)$ given by those $D_{r-i}(\phi)$ with $i \geq 0$. Over $D_{r-i}(\phi) - D_{r-i-1}(\phi)$ the fiber of the map $\pi: Z(\phi_r) \rightarrow D_r(\phi)$ is $\text{Gr}(e-r, e-r+i)$. So

$$\pi: Z(\phi_r) \rightarrow D_r(\phi)$$

is a quasi-fibration.

When the section ϕ_r of $\text{Hom}(U, \pi^*F)$ is transversal, $Z(\phi_r)$ are smooth subvarieties of

G , so that they can be considered as a desingularization of $D_r(\phi)$. Moreover $\dim G = \dim M + 2r(e - r)$ and $\text{rank}_{\mathbb{C}}(U^* \otimes \pi^* F) = f(e - r)$, so that the dimension of $Z(\phi_r)$ is $2n - 2(e - r)(f - r)$. If this is negative then $D_r(\phi)$ is empty. Finally let

$$\rho(r) = n - (e - r)(f - r).$$

Note that it must be $r \leq e$ and $r \leq f$ since otherwise we have $D_r(\phi) = M$.

We pose ourselves a question: what can be said about the topology of the degeneracy loci $D_r(\phi)$?

6 A Lefschetz Theorem

Let M be a complex manifold and let $E \rightarrow M$ be a rank e holomorphic vector bundle. For a holomorphic section s of E , let $W = Z(s)$ be the zero set of s . The topology of W can be controlled under an extra assumption on E .

We say that E is a positive vector bundle [We, page 223][Gr, Chapter 0] if there exists an hermitian metric h_E in the fibers of E such that the curvature Θ associated to the hermitian connection induced by the holomorphic structure satisfies that $\sqrt{-1}\Theta$ is a positive 2-form: $\langle s, \sqrt{-1}\Theta(u, Ju)s \rangle > 0$ for any non-zero $u \in T_p M$ and non-zero $s \in E_p$, where Ju is the vector obtained by multiplying u by i in the complex space $T_p M \cong \mathbb{C}^n$. To go into details of the definition of curvature and consequences of positivity see [We]. When E is of rank 1 this produces a Kähler form for M .

Lefschetz theorem on hyperplane sections. *Let E be a rank e positive vector bundle over a compact complex manifold M and let s be a holomorphic section of E . Let $W = Z(s)$ be the zero set of s and $\rho = n - e$. If the section s is transverse then W has dimension 2ρ . In any case, the pair (M, W) is ρ -connected.*

The result is originally given in [AF][Bo]. We also have a Lefschetz theorem for degeneracy loci given in [MP] (see [De] for the homology version built up on considerations of connectedness of degeneracy loci in [FL]).

Lefschetz theorem for degeneracy loci. *Let ϕ be a morphism between the holomorphic vector bundles E and F , of ranks e and f respectively, over a compact complex manifold M , such that $\text{Hom}(E, F)$ is positive. Let $W = Z(\phi_r) \subset G = \text{Gr}(e - r, E)$ and $\rho = n - (e - r)(f - r)$. Then (G, W) is ρ -connected.*

The theorem does not follow from the previous Lefschetz theorem, since the positivity of $\text{Hom}(E, F)$ does not imply that $\text{Hom}(\mathcal{U}, \pi^*F)$ is positive. Actually the proof of the theorem in [MP] goes through the way of proving that $\text{Hom}(\mathcal{U}, \pi^*F)$ keeps "part of" the positivity of $\text{Hom}(E, F)$, namely the positivity along the section ϕ_r .

Our application is the following information on the topology of $D_r(\phi)$, using the map $\pi: G \rightarrow M$.

Theorem. *Let ϕ be a morphism between the holomorphic vector bundles E and F over a compact complex manifold M , such that $\text{Hom}(E, F)$ is positive. Suppose that $\rho(r+k) \geq 0$, for some $k \geq 0$. Let $\varepsilon = 0$ if $\rho(r+k) > 0$ and $\varepsilon = 1$ if $\rho(r+k) = 0$. Then the pair $(M, D_r(\phi))$ is $(2k+1-\varepsilon)$ -connected.*

Proof. We prove this by induction on k . For $k = 0$ it is very easy: if $\rho(r) = 0$ then $(G, Z(\phi_r))$ is 0-connected, i.e., every connected component of G has at least a point of $Z(\phi_r)$. Since the connected components of G are in 1-1 correspondence with those of M and $Z(\phi_r)$ surjects onto $D_r(\phi)$, the same happens to $(M, D_r(\phi))$. If $\rho(r) > 0$ then $(G, Z(\phi_r))$ is 1-connected. The commutative diagram

$$\begin{array}{ccc} \pi_1(Z(\phi_r)) & \rightarrow & \pi_1(G) \\ \downarrow & & \parallel \\ \pi_1(D_r(\phi)) & \rightarrow & \pi_1(M) \end{array} \quad (1)$$

implies that the arrow in the bottom row is a surjection. Therefore $(M, D_r(\phi))$ is also 1-connected.

Now let us turn to the case $k > 0$. We may suppose that M is connected from now on, working on every connected component of M if necessary. If $\rho(r-1) \geq 0$ then $\rho(r) \geq (e-r) + (f-r) + 1 \geq 2$, unless $e = f = r$ in which case $D_r(\phi) = M$ and there is nothing to prove. If $\rho(r) \geq 2$ then $(G, Z(\phi_r))$ is 2-connected. Then the top row in diagram (1) is an isomorphism. By the result in section 4, $\pi_1(Z(\phi_r)) \rightarrow \pi_1(D_r(\phi))$ is a surjection. This implies that the bottom row in (1) must be an isomorphism. So the fundamental groups of $Z = Z(\phi_r)$, G , M and $D_r = D_r(\phi)$ all coincide. Let now $p: \widetilde{M} \rightarrow M$ be the universal covering space. Also $\tilde{p}: p^*G \rightarrow G$ is a universal covering space, and $\widetilde{Z} = \tilde{p}^{-1}(Z)$, $\widetilde{D}_r = \tilde{p}^{-1}(D_r)$ are the universal covering spaces of Z and D respectively. Pulling back all our construction by p we see that to prove that $(M, D_r(\phi))$ is n -connected is equivalent to prove that $(\widetilde{M}, \widetilde{D}_r(\phi))$ is n -connected.

The above considerations imply that we can work on \widetilde{M} . To simplify notation, we denote \widetilde{M} by M , i.e., we assume to start with that M is simply-connected and keep the

notations without tildes. Under this condition, Hurewicz theorem in section 1 says that (M, D_r) is $(2k+1-\varepsilon)$ -connected if we manage to prove that

$$H_i(M, D_r) = 0, \quad i \leq 2k+1-\varepsilon.$$

By Lefschetz theorem (G, Z) is $\rho(r)$ -connected. Now $\rho(r-k) \geq 0$ implies that $\rho(r) = (\varepsilon-r)k + (f-r)k + k^2 \geq 2k+1$ (again if either $\varepsilon=r$ or $f=r$ then $D_r(\phi) = M$ and there is nothing to prove). So $H_n(G, Z) = 0$ for $n \leq 2k+1$.

Let us stratify M with $M \supset D_r \supset D_{r-1} \supset \dots \supset D_{r-k}$ and let G be stratified with $G_{r-i} = \pi^{-1}(D_{r-i})$ and $Z_{r-i} = Z \cap G_{r-i}$. With this notations in place, we have that

$$(G_{r-i} - G_{r-i-1}, Z_{r-i} - Z_{r-i-1}) \rightarrow D_{r-i} - D_{r-i-1}$$

is a fibration with fiber $(\text{Gr}(e-r, e), \text{Gr}(e-r, e-r+i))$. Let $i > 0$. The last theorem in section 3 implies that there is a spectral sequence with $E^2 = H_*(D_{r-i}, D_{r-i-1}) \otimes H_*(\text{Gr}(e-r, e), \text{Gr}(e-r, e-r+i))$ converging to $E^\infty = H_*(G_{r-i}, G_{r-i-1} \cup Z_{r-i})$. (We can take the homology of the grassmannian out of the 'coefficient group' since it is free.) Now the pair $(\text{Gr}(e-r, e), \text{Gr}(e-r, e-r+i))$ is $(2i+1)$ -connected. By induction hypothesis, the pair (M, D_{r-i}) is $(2(k-i)+1-\varepsilon)$ -connected. Therefore by the long exact sequence of the pair, $H_n(D_{r-i}, D_{r-i-1}) = 0$ for $n \leq (2(k-i)-\varepsilon)$. Hence $E_{pq}^2 = 0$ for $p+q \leq 2(k-i)-\varepsilon+(2i+1)+1$. This implies that $H_n(G_{r-i}, G_{r-i-1} \cup Z_{r-i}) = 0$ for $n \leq 2k+2-\varepsilon$.

The spectral sequence for the filtration $G \supset G_r \supset G_{r-1} \supset \dots \supset G_{r-k}$ has

$$\begin{aligned} E_{p, r+1}^2 &= H_{p+r+1}(G, G_r), \\ E_{p, r-i}^2 &= H_{p+q}(G_{r-i}, G_{r-i-1} \cup Z), \quad i \geq 0, \end{aligned}$$

and converges to $E^\infty = H_n(G, Z)$. By the above considerations $E_{pq}^2 = 0$ for $q < r$ and $p+q \leq 2k+2-\varepsilon$. Looking at the limit, $H_n(G, Z) = 0$ for $n \leq 2k+1$. We have that it must be $d_1 = \partial_* : H_n(G, G_r) \rightarrow H_{n-1}(G_r, G_{r-1} \cup Z)$ an isomorphism, for $n \leq 2k+2$.

q			
r+1	•	•	•
r	⋮	⋮	⋮
	•	•	•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
			•
			⋮
	</		

Now there are two spectral sequences: $E_{ij}^2 = H_j(M, D_r) \otimes H_i(\text{Gr}(e - r, e))$ converging to the $H_n(G, G_r)$, and $\hat{E}_{ij}^2 = H_{j-1}(D_r, D_{r-1}) \otimes H_i(\text{Gr}(e - r, e), pt)$ converging to $H_{n-1}(G_r, G_{r-1} \cup Z)$, where pt is the base point. Here we have 'moved' the second spectral sequence one unit to the top so that $f = \partial_*$ induces a map of bidegree $(0, 0)$ between the two spectral sequences, i.e., a map $f^r : E^r \rightarrow \hat{E}^r$ for every step, such that $f^{r+1} = H(f^r)$. In the limit, $f^\infty : E^\infty \rightarrow \hat{E}^\infty$ must be an isomorphism.

Look at $f_{ij}^2 : E_{ij}^2 \rightarrow \hat{E}_{ij}^2$. For $i + j \leq 2k + 2 - \varepsilon$ it is surjective, since $H_j(M, D_r) \rightarrow H_{j-1}(D_r, D_{r-1})$ is surjective for $j \leq 2k - \varepsilon$. Also f_{ij}^2 is injective for $i > 0$, $j \leq 2k - 1 - \varepsilon$, since $H_j(M, D_r) \rightarrow H_{j-1}(D_r, D_{r-1})$ is injective for $j \leq 2k - 1 - \varepsilon$. Now let us prove that $f^r : E_{ij}^r \rightarrow \hat{E}_{ij}^r$ is isomorphism for $i + j \leq 2k + 1 - \varepsilon$, $i > 0$ and an epimorphism for $i + j \leq 2k + 2 - \varepsilon$, by induction on r . We consider the diagram

$$\begin{array}{ccccc} E_{i-r+1, j+r}^r & \xrightarrow{d_r} & E_{ij}^r & \xrightarrow{d_r} & E_{i+r-1, j-r}^r \\ f_{i-r+1, j+r}^r \downarrow & & f_{ij}^r \downarrow & & f_{i+r-1, j-r}^r \downarrow \\ E_{i-r+1, j+r}^{r+1} & \xrightarrow{d_r} & E_{ij}^{r+1} & \xrightarrow{d_r} & E_{i+r-1, j-r}^{r+1} \end{array}$$

For $i = 0$ we have $\hat{E}_{0j}^r = 0$, so it must be $d_r : E_{0j}^r \rightarrow E_{r-1, j-r}^r$ the zero map for $j \leq 2k + 2 - \varepsilon$. Hence $E_{0j}^{r+1} \cong E_{0j}^r$.

For $i + j < 2k + 1 - \varepsilon$ and $i > 0$, $i \neq r - 1$, the vertical arrows are isomorphisms, so f_{ij}^{r+1} is an isomorphism. For $i = r - 1$ the result also holds using that $d_r : E_{0j}^r \rightarrow E_{r-1, j-r}^r$ is the zero map. For $i + j = 2k + 1 - \varepsilon$, $i > 0$, the first vertical arrow is an epimorphism, the second and third are isomorphisms, so f_{ij}^{r+1} is an isomorphism. Finally for $i + j = 2k + 2 - \varepsilon$, f_{ij}^r is an epimorphism and $f_{i+r-1, j-r}^r$ is an isomorphism, hence f_{ij}^{r+1} is an epimorphism.

Looking at the E^∞ stage, $f_{0j}^\infty : E_{0j}^\infty \rightarrow \hat{E}_{0j}^\infty = 0$ is an isomorphism for $j \leq 2k + 1 - \varepsilon$. This means that $E_{0j}^\infty = E_{0j}^2 = H_j(M, D_r) = 0$ for $j \leq 2k + 1 - \varepsilon$ as desired. \square

References

- [ACGH] ARBARELLO, E., CORNALBA, M., GRIFFITHS, P.A. AND HARRIS, J., *Geometry of algebraic curves*, Grundlehren der Mathematischen Wissenschaften, Vol. 267, Springer-Verlag, New York, 1985.
- [AF] ANDREOTTI, A. AND FRAENKEL, T., *The Lefschetz theorem on hyperplane sections*, *Annals Math.* (2) **69** (1959), 713-717.
- [Bo] BOTT, R., *On a theorem of Lefschetz*, *Michigan Math. J.* **6** (1959), 211-216.

- [BT] BOTT, R. AND TU, L.W., *Differential forms in algebraic topology*, Graduate Texts in Maths, Vol. 82, Springer-Verlag, 1982.
- [De] DEBARRE, O., *Lefschetz theorems for degeneracy loci*, Bull. Soc. Math. France **128** (2000), 283-308.
- [GM] GORESKEY, M. AND MACPHERSON, R., *Stratified Morse theory*.
- [Gr] GRIFFITHS, P.A., *Hermitian differential geometry, Chern classes and positive vector bundles*, in Global Analysis, Princeton University Press, Princeton, N.J., 1969.
- [FL] FULTON, W. AND LAZARSFELD, R., *On the connectedness of degeneracy loci and special divisors*, Acta Math. **146** (1981), 271-283.
- [Ma] MASSEY, W.S., *Algebraic Topology: an Introduction*, Graduate Texts in Maths, Vol. 56, Springer-Verlag, 1977.
- [Mc] MCCLEARY, J., *A user's guide to spectral sequences*, Second edition. Cambridge Studies in Advanced Mathematics, Vol. 58. Cambridge University Press, Cambridge, 2001.
- [MP] MUÑOZ, V. AND PRESAS, F., *Semipositive bundles and Brill-Noether theory*, math.DG/0107226.
- [Sp] SPANIER, E.H., *Algebraic topology*, McGraw-Hill Book Co., 1966.
- [We] WELLS JR., R.O., *Differential Analysis on Complex Manifolds*, Springer Verlag, Graduate Texts in Maths, Vol. 65, 1980.