

Erratum to: Lipase and invertase activities in midgut and salivary glands of *Chilo suppressalis* (Walker) (Lepidoptera, Pyralidae), rice striped stem borer [5: 180-189, 2008]

A Zibae, AR Bandani, S Ramzi

Plant Protection Department, Faculty of Agriculture, University of Tehran, Karaj 31584, Iran

In the above article **Table 2** was reproduced incorrectly, the table and caption should have appeared as below:

Table 2 Relative activity of *C. suppressalis* invertase towards different compounds

Compounds	Concentration (mmol/l)	Relative activity (midgut)	Relative activity (salivary gland)
Control	-	100	100
NaCl	5	83.33	152.63*
	10	63.88*	121.05*
	20	41.66*	94.7*
	40	10*	52.63*
CaCl ₂	5	25.83*	78.94*
	10	66.66*	131.57*
	20	113.88	157.89*
	40	125*	194.73*
KCl	5	161.11*	273.68*
	10	105.55	252.63*
	20	69.44*	194.73*
	40	50*	89.47*
MgCl ₂	5	97.22	168.42*
	10	61.11*	152.63*
	20	44.44*	84.21*

	40	21.38*	27.36*
EDTA	0.5	144.44*	168.42*
	1	116.66	131.57*
	2	97.22*	105.28
	4	52.77*	73.68*
SDS	2	111.11	173.68*
	4	91.66	136.84*
	6	61.11*	121.05
	8	38.88*	73.68*
Urea	1000	83.33*	173.68*
	2000	63.88*	147.36*
	4000	50*	115.78
	5000	33.33*	73.68*
	6000	25*	47.36*
Plant extract	10 %	82*	91*
	15 %	64*	74*
	25%	21*	52*

* $P < 0.05$ vs control

Corresponding author:
 Ali Reza Bandani
 Plant protection Department
 College of agriculture and natural resources
 University of Tehran, Karaj, 31584 Iran
 E-mail: abandani@ut.ac.ir