Letter to the Editor

Sand Fly Fever with Different Names

Ali Mehrabi-Tavana

Health Management Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

(Received 16 July 2016; accepted 15 Jan 2017)

Dear Editor-in-Chief

With regard to a paper published in Journal of Arthropod-Borne Diseases;

A Case Report entitled "Sand fly Fever with Skin Lesions: A Case Series from Turkey" was published in 2016 (Fatih Temocin, et al. 2016). Different names such as "Phlebotomus fever", "mosquito fever", three-day fever or "Papatasi fever" called the disease of Sand fly fever which all are true except "mosquito fever" (Tavana 2001, 2015). Mosquito is not a general name in English and is used just for Culicidae. Blackflies and biting midges are used for Simuliidae and Ceratopogonidae, respectively. Therefore, is not related to sand fly fever (Saidi et al. 1977, Izri et al. 2008). In my opinion it should be corrected. Sand fly fever is an arthropod-borne viral disease, starts with acute onset of high fever, and lasts for two to four days, sometimes much longer.

Photophobia and muscle ache are more common in the patients. This disease is more prevalent in Europe, Asia, and Persian Gulf region (Konstantinou et al. 2007, Tavana 2007). It is important to be aware and take care of this disease in endemic and no-endemic area among travelers returning from endemic areas.

References

- Fatih Temocin, Tugba Sari, Necla Tulek (2016) Sand fly Fever with Skin Lesions: A Case Series from Turkey. J Arthropod-Borne Dis. 10(4): 608–612.
- Tavana AM (2001) The seroepidemiological studies of sand fly fever in Iran during imposed war. Iran J Public Health. 30: 145–146.
- Tavana AM (2007) Minire view on sand fly fever. J Entomol. 4: 401–403.
- Tavana AM (2015) Sand fly fever in the world. Ann Trop Med Public Health. 8: 83–87.
- Saidi S, Tesh RB, Javadian E, Sahabi Z, Nadim A (1977) Studies on the epidemiology of sand fly fever in Iran. II. The prevalence of human and animal infection with five Phlebotomus fever virus serotypes in Isfahan Province. Am J Trop Med Hyg. 26: 288–293.
- Konstantinou GN, Papa A, Antoniadis A (2007) Sand fly-fever outbreak in Cyprus: are phleboviruses still a health problem? Travel Med Infect Dis. 5: 239–242.
- Izri A, Temmam S, Moureau G, Hamrioui B, deLamballerie X, Charrel RN (2008) Sand fly fever Sicilian virus, Algeria. Emerg Infect Dis. 14: 795–797.

http://jad.tums.ac.ir Published Online: March 14, 2017