

JOURNAL OF HORTICULTURAL SCIENCES

Volume 16

June 2021

Issue 1



Society for Promotion of Horticulture

ICAR - Indian Institute of Horticultural Research, Bengaluru - 560 089





JOURNAL OF HORTICULTURAL SCIENCES

(Founded in 2005 by the Society for Promotion of Horticulture, Bengaluru, India)

Email : chiefeditor.jhs@gmail.com Webpage : <https://jhs.iihr.res.in/index.php/jhs>

Editor-in-Chief

Dr. S. Sriram

Editors

Dr. K. Himabindu

Dr. G. Senthilkumaran

Dr. Tejaswini Prakash

Dr. M. Manamohan

Dr. Anil Kumar Nair

Dr. J. Satisha

Dr. P. Venkata Rami Reddy

Dr. I.N. Doreyappa Gowda

Dr. R.H. Laxman

Dr. G.C. Sathisha

Editorial Advisory Board

International Editorial Advisory Board

Dr. Nanthi S. Bolan, Australia

Dr. Rod Drew, Australia

Dr. J. Mithila, USA

Dr. Claus Helmut Franz Orth, South Africa

Dr. IlanParan, Israel

Dr. Gi-Cheol Song, Republic of Korea

Dr. Jill Stanley, New Zealand

Dr. PalithaWeerakkody, Sri Lanka

National Editorial Advisory Board

Dr. S.D. Shikhamany

Dr. V.A. Parthasarathy

Dr. K.V. Peter

Dr. Sisir K. Mitra

Dr. S.K. Tikoo

Dr. SeetharamaAnnadama

Dr. A. Krishnamoorthy

Dr. Leela Sahijram

SOCIETY FOR PROMOTION OF HORTICULTURE (REGD.)

Email : sphiihr2005@gmail.com Website : sph.iihr.res.in

Executive Council 2021

President : Dr. M.R. Dinesh

Vice Presidents : Dr. C. Aswath

Dr. K. Madhavi Reddy

General Secretary : Dr. M.V. Dhananjaya

Editor-in- Chief : Dr. S. Sriram

Treasurer : Dr. E. Sreenivasa Rao

Joint Secretaries : Dr. T.H. Singh

Dr. Laxmi Narayan Mahawar

Members : Dr. B.N. Srinivasa Murthy

Dr. K. Manivannan

Dr. Jayoti Majumder

Dr. Rajendra Singh Rathore

This Journal is abstracted in CABI, Current Contents, AGRIS, Indian Science Abstracts, Scopus, DOAJ and Redalyc. It is a participant of AmelICA.

Request for membership subscriptions along with cheque/DD drawn in favour of **Society for Promotion of Horticulture, Bengaluru** may be sent to General Secretary, Society for Promotion of Horticulture, Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bengaluru - 560 089, India. All members except student members and subscribers get all publications of SPH free of cost. Any correspondence other than editorial may be addressed to General Secretary, Society for Promotion of Horticulture, Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bengaluru - 560 089, India.

Sponsorship for programmes related to promotion of horticulture is accepted.

For information please contact
General Secretary, SPH, IIHR,
Bengaluru - 560089, India.

Coverpage Courtesy :

Top : **Lavanya et al.**, P. 67

Bottom : **Waman et al.**, P. 74

SUBSCRIPTION RATES

Patron	₹ 20,000
Life member	₹ 5,000
Annual Member	₹ 1,000 / US \$ 100 (US \$ 50 for SAARC countries)
Student Member	₹ 500
Student Life Memeber	₹ 3,000
Annual Subscription	₹ 1,500 / US \$ 100 (US \$ 60 for SAARC countries) (for institutions)
Enrolment Fee	₹ 200 / US \$ 5 (Additional for all types of Membership)

NAAS rating of this journal is 5.08. JHS is now available online.
Authors have to submit manuscripts using the link : <https://jhs.iihr.res.in/index.php/jhs>

Technical Assistance : **Dr. Sridhar Gutam, Thippeswamy S. and Pramida A.**

CONTENTS

In this Issue

i-ii

Review

Moringa (*Moringa oleifera* L.): An underutilized and traditionally valued tree holding remarkable potential 1-13

Jattan M., Kumari N., Raj Kumar, Kumar A., Rani B., Phogat D.S., Kumar, S. and Kumar, P.

Original Research in Papers

Characterization and evaluation of mountain sweet thorn (*Flacourtia montana* J. Grah) collections 14-25

Tripathi P.C., Ganeshan S., Radhika V. and Shetti D.L.

Optimization of methodology for the extraction of polyphenolic compounds with antioxidant potential and α -glucosidase inhibitory activity from jamun (*Syzygium cumini* L.) seeds 26-35

Arivalagan M., Priyanka D.R. and Rekha A.

Genetic variability studies in amaranthus (*Amaranthus* spp.) 36-44

Agadi A.H., Kolakar S., Lakshmana D., Nadukeri S. and Hanumanthappa M.

Morpho-physiological parameters associated with chlorosis resistance to iron deficiency and their effect on yield and related attributes in potato (*Solanum tuberosum* L.) 45-52

Challam C., Dutt S., Sharma J., Raveendran M. and Sudhakar D.

Responses of different Okra (*Abelmoschus esculentus*) cultivars to water deficit conditions 53-63

Ayub Q., Khan S.M., Hussain I., Naveed K., Ali S., Mehmood A., Khan M.J., Haq N.U., Shehzad Q.

Induced variability for yield and its attributing traits in cluster bean [*Cyamopsis tetragonoloba* (L.) Taub] through gamma irradiation 64-68

Lavanya H.N., Mishra S., Sood M., Aghora T.S., Anjanappa M., Rao V.K. and Reddy A.B.

***In vitro* multiplication protocol for *Curcuma mangga* : Studies on carbon, cytokinin source and explant size** 69-76

Waman A.A., Bohra P., Karthika Devi R. and Pixy J.



- Effect of fungicide and essential oils amended wax coating on quality and shelf life of sweet orange (*Citrus sinensis* Osbeck)** 77-90
Bhandari M., Bhandari N. and Dhital M.
- Post-harvest quality and quantification of betalains, phenolic compounds and antioxidant activity in fruits of three cultivars of prickly pear (*Opuntia ficus-indica* L. Mill)** 91-102
Gonzalez F.P.H., Saucedo V.C., Guerra R.D., Suarez E.J., Soto H.R.M. Lopez J.A., Garcia C.E. and Hernandez R.G.
- Soil microbial community dynamics as influenced by integrated nutrient management practices in sweet basil (*Ocimum basilicum* L.) cultivation** 103-113
Baraa AL-Mansour and D. Kalaivanan
- Effect of spectral manipulation and seasonal variations on cut foliage production and quality of *Philodendron* (*Philodendron* 'Xanadu')** 114-120
Sujatha A. Nair, Laxman R.H. and Sangama

Short Communications

- Studies on mutagenic sensitivity of seeds of pummelo (*Citrus maxima* Merr.)** 121-124
Sankaran M., Kalaivanan D. and Sunil Gowda D.C.
- Isolation and characterization of microsatellite markers from *Garcinia indica* and cross species amplification** 125-129
Ravishankar K.V., Vasudeva R., Hemanth B., Nischita P., Sthapit B.R., Parthasarathy V.A. and Rao V.R.

In this issue...

*The farmers involved in the production of horticultural produce have defied the challenges due to second wave of the **COVID pandemic**. Especially, the information that India's horticulture production is expected to increase by 2.93 per cent to a record level of 329.86 million tonnes (MT) in 2020-21, according to the second advance estimate of horticulture production released by the Ministry of Agriculture is very much motivating. Though many hurdles are faced by the farmers due to the prevailing **Covid Pandemic** with respect to timely marketing, storage and other issues, the horticultural production has maintained the steady growth. Wish that in other countries also the horticulture production enhances to provide the nutritional security. Journal of Horticultural Sciences takes pride in sharing the recent research developments in different disciplines of Horticulture.*

*Many native vegetable crops have nutraceutical values. One of them is Moringa or commonly called as drumstick. **Jattan et al.** have reviewed the traditional values, requirement for crop cultivation, crop improvement, value addition, nutraceutical and pharmaceutical values with respect to this crop.*

*Many under-utilized fruit crops are brought to cultivation again with scientific interventions. One such crop is mountain sweet corn (*Flacourtia montana*). The fruits have high potential for processing into jam, jelly, wine, etc. and this plant has good medicinal value. **Tripathi et al.** evaluated and characterized the mountain sweet corn accessions from Western Ghats and identified a suitable line with higher yield, regular bearing nature and less thorniness. **Ravishankar et al.** isolated and characterized the microsatellite markers from the under-utilized tree species *Garcinia indica*. In their study, 3725 microsatellites were identified and primers were designed for 1374 microsatellites. The SSR developed will be useful in studying genetic diversity, mapping and fingerprinting of *Garcinia indica* and related species.*

*Among the leafy vegetables, *Amaranthus* ranks first in production. **Agadi et al.** estimated the nature and extent of genetic variability among twenty *Amaranthus* genotypes. They found that Arka Arunima, Chikmagalur local, IC-551486, IC-551494 and IC-551466 recorded high foliage yield per plot and could be utilized in further breeding programme. **Challam et al.** studied the morpho-physiological parameters associated with resistance to iron deficiency induced chlorosis in potato and their effect on yield attributed. They found that genotype CP-3443) was found tolerant to Fe deficiency induced chlorosis. **Ayub et al.** evaluated responses of different okra (*Abelmoschus esculentus*) cultivars to water deficit conditions in Pakistan. They concluded that drought caused significant variation on physical and biochemical attributes of okra whereas the cultivar 'Sabz Pari' showed resistance towards the water stress.*

*Gamma ray is an effective mutagen which creates useful variability in crops where the natural variation is very meagre and creation of variability by conventional methods is cumbersome. **Lavanya et al.** studied the induced variability in cluster bean due to gamma irradiation and found that the traits like*



plant height, pod length, pod width, pulp to seed ratio showed sufficient variability. **Sankaran et al.** also employed the gamma irradiation to generate variability in pummelo. They found that 60 Gy gamma dose can effectively be used for raising the mutant populations to identify a desirable mutation in pummelo

Soil microbiome plays important role in crop production. There is need to pay attention on the nutrition management practices. They should encourage the soil microbe population that will indirectly help the plant health. **Al-Mosour and Kalaivanan** have demonstrated that integrated nutrient management can maximize soil microbial community dynamics which is considered as driving force behind regulating soil processes that support sustainable sweet basil cultivation.

While attempting to evaluate the spectral manipulations on cultivation of cut foliage crops, **Nair et al.** found that coloured shade nets did not influence vase life of the cut foliage in *Philodendron* and observed that cultivation of *Philodendron* 'Xanadu' under white shade resulted in maximum cut foliage yield and quality.

Mango ginger is an underutilized rhizomatous species that has been valued in the tropical Asian countries as a source of vegetable, spice, salad, medicine and essential oil. Huge quantity of seed rhizomes is required to promote this crop in larger area. **Waman et al.** developed an in vitro protocol for the multiplication of mango ginger.

Jamun seed is a rich source of polyphenolic compounds with antioxidant potential and alpha - glucosidase inhibitory activity. **Arivalagan et al.** have optimized the methodology for the extraction of such polyphenols from jamun seeds. This will be of much in nutraceutical industry. Similarly, **González et al.** studied the post-harvest quality and quantification of betalains, phenolic compounds and antioxidant activity in fruits of three cultivars of prickly pear in Mexico. They observed that there was high correlation between antioxidant activity and phenolic compounds. The methodologies developed by them will be useful tool for the quantification of bioactive compounds fruit tissues.

Post-harvest disease management is an important aspect in delivering the harvested produce safely to the end-user. **Bhandari et al.** have recorded that application of essential oils with wax improved shelf life of sweet oranges in Nepal and this treatment enhanced juice retention, firmness, titratable acidity, vitamin C and disease reduction.

S. Sriram
Editor in Chief

Original Research Paper

Optimization of methodology for the extraction of polyphenolic compounds with antioxidant potential and α -glucosidase inhibitory activity from Jamun (*Syzygium cumini* L.) seeds

Arivalagan M.^{1*}, Priyanka D.R.¹ and Rekha A.²

¹Division of Basic Sciences, ²Division of Fruit Crops
ICAR-Indian Institute of Horticultural Research (IIHR), Bengaluru - 560 089, Karnataka, India.

*Corresponding author E. mail: arivalagan2100@gmail.com

ABSTRACT

Jamun (*Syzygium cumini* L.) seed is one of the rich sources of polyphenolic compounds with antioxidant potential and α -glucosidase inhibitory activity. A study was conducted to optimize the methodology for the extraction of polyphenolic compounds (total phenolic and flavonoid contents) with antioxidant potential and α -glucosidase inhibitory activity from jamun seed powder. The study showed that the nature of solvent and extraction conditions had a significant effect on total phenolic content (TPC), total flavonoid content (TFC), antioxidant potential, and α -glucosidase inhibitory activity. The TPC varied between 6.0 (mg/g jamun seed powder) for the acetone extract to 119.2 (mg/g) for 80% aqueous acetone extract, and TFC varied between 1.06 mg/g for the acetone to 10.81 mg/g for the 80% aqueous methanol. From the study, it was apparent that an aqueous form of acetone (acetone: water 80:20, v/v) is a better solvent system for extraction of polyphenolic compounds with high antioxidant potential and α -glucosidase inhibitory activity. Ultrasonication for 60 min increased the efficiency of phenolic extraction.

Keywords: α -glucosidase inhibitory activity, Antioxidant potential; Flavonoids, Jamun seed powder, Natural antioxidants and Polyphenols.

INTRODUCTION

Jamun (*Syzygium cumini* L.) is one of the most important indigenous fruits belongs to the family Myrtaceae. It is a treasure of wide range of secondary metabolites with numerous health benefits. The fruit pulp is sweet and the seeds are pungent and tangy. Both pulp and seeds are extensively used in traditional medicine against various ailments such as diabetes, diarrhea, and ringworm (Benherlal and Arumughan, 2007). While jamun fruits are a rich source of anthocyanins, their seeds are high in polyphenolic compounds. Oxalic acid, tannic acid, gallic acid, and some of the alkaloids are the major secondary metabolites responsible for the astringency taste of seed as well as pulp (Hameed *et al.*, 2020). Most of the secondary metabolites, especially polyphenolic compounds present in the jamun pulp and seeds are reported to possess free radical scavenging potential (Ayyanar and Subash-babu, 2012) as well as anti-inflammatory activities (De Bona *et al.*, 2016; Hossain *et al.*, 2016).

Earlier studies have suggested that some of the polyphenolic compounds present in jamun fruit pulp and seeds have the potential to inhibit the activities of the enzymes, alpha-amylase and intestinal alpha-glucosidase which are responsible for the digestion of dietary starch (Mahmood, 2016). These enzymes have been a target of many drugs developed for the treatment of type-II diabetes (Sim *et al.*, 2008). Since, diabetes is a major problem affecting the human population worldwide and synthetic drugs are risky due to associated side effects, plant-based secondary metabolites especially polyphenolic compounds are the ideal candidate to address the problems of hyperglycemia and diabetes in the human population. Investigation made earlier has reported that the presence of polyphenolic compounds in jamun seeds has appreciably good antioxidant potential and hypoglycemic effect (Aqil *et al.*, 2012). Thus, the extraction of polyphenolic compounds from jamun seeds and its usage in the medication for regulation



of blood glucose level has attracted many researchers. Polyphenolic compounds possess diverse nature in terms of physical and chemical properties and some of them exist in the free as well as complex forms with carbohydrates and proteins. Due to this nature, the extent of the solubility of these polyphenolic compounds varies in different solvents (Khoddami *et al.*, 2013). Mostly water and other polar solvents such as ethanol, methanol, acetone, and diethyl ether have been extensively used for the extraction of polyphenolic compounds from plant sources (Arivalagan *et al.*, 2018). Since, the levels of active principle in the plant extract depends on the type of solvent used and the extraction methods being adopted (Sun and Ho, 2005; Turkmen *et al.*, 2006; Hayouni *et al.*, 2007), the present investigation was undertaken to optimize the suitable solvent system and extraction conditions for the extraction of polyphenolic compounds with antioxidant potential and alpha-glucosidase inhibitory activity from jamun seeds, as it is rich in polyphenolic compounds.

MATERIALS AND METHODS

Chemicals

DPPH (1,1-diphenyl-2-picrylhydrazyl), Trolox (6-hydroxy-2,5,7,8-tetramethyl chroman-2- carboxylic acid), TPTZ (2,4,6-tris-2,4,6-tripyridyl-2-triazine), ABTS (2,2-azino-di-(3-ethylbenzothiazolone-6-sulphonic acid) diammonium salt), potassium persulfate, and neocuproine (2,9-dimethyl-1,10-phenanthroline) were purchased from Sigma-Aldrich Co St. Louis, MO, United States of America. Analytical grade Ethanol, methanol, acetone, acetic acid (glacial), sodium acetate, hydrochloric acid (conc.), ferric chloride, ammonium acetate, copper (II) chloride, Folin–Ciocalteu’s phenol reagent, aluminium chloride, sodium nitrite, and sodium carbonate were purchased from Merck KGaA, Darmstadt, Germany.

Sample preparation

The experiment was conducted at ICAR- Indian Institute of Horticultural Research (IIHR), Bengaluru, Karnataka, India during 2020. Matured and fully ripened fruit samples were randomly collected from jamun germplasm (IC-0621954, 8 years old tree with semi-spreading type with broad leaves) and brought to the laboratory in the icebox. After removal of pulp, the seeds were dried in a mechanical tray dryer

(Make: M/s Servewell Instruments Pvt. Ltd, Karnataka, India) at 50°C until the constant dry weight was achieved. The seed coat was then removed and the kernels were powdered finely using a ball mill (Mixer Mill MMOL400, RETSCH GmbH, Germany) to get uniform particle size and stored in a vacuum desiccator for further analysis.

Solvent system and extraction procedure

About 100 mg jamun seed powder was mixed with 5 mL of selected solvent in 15 mL centrifuge tube and extracted. For extraction seven solvent systems *viz.* water, methanol, aqueous methanol (methanol: water 80:20 v/v), ethanol, aqueous ethanol (ethanol: water 80:20 v/v), acetone, and aqueous acetone (acetone: water 80:20 v/v) were used. The extraction procedure was carried out at two different conditions (room temperature [25±2°C] and ultra-sonic bath set at 50±2°C), each with 30 min and 60 min separately. Extracts were centrifuged (at 5°C for 10 min) and the supernatant was collected in amber reagent bottles. The residue was re-extracted twice using the same procedure as mentioned above. Filtrates collected from all three successive extractions were pooled and dried under a nitrogen atmosphere. The dried samples were dissolved in 5 mL of water, centrifuged, and then the clear supernatant collected was used for the analysis.

Determination of total phenolic and flavonoid content

Determination of total phenolic content (TPC) in the extract was done by Folin-Ciocalteu (FC) assay as described by Singleton *et al.* (1999) using gallic acid as a standard, and TPC was expressed as mg gallic acid equivalent (GAE), per g dry weight of jamun seed powder. The total flavonoid content (TFC) was determined according to Zhishen *et al.* (1999). Quercetin was used as a standard and the results were expressed as mg of quercetin equivalent (QE) per g jamun seed powder.

Determination of antioxidant potential

The DPPH and ABTS radical scavenging activities (IC₅₀) of the different solvent extracts of jamun seed powder were determined by the methods of Brand-Williams *et al.* (1995) and Arnao *et al.* (2001), respectively. IC₅₀ value signifies the concentration of test samples to scavenge 50% of the DPPH and ABTS•⁺ radical and expressed as mM TE/g jamun

seed powder. The FRAP (Ferric Reducing Antioxidant Power) assay was done according to Benzie & Strain (1996) and CUPRAC (Cupric ion reducing antioxidant capacity) of the extract was determined according to the method of Apak *et al.*, (2004). In both methods, Trolox served as a positive control and results were expressed as mM TE/g dry jamun seed powder.

Determination of alpha-glucosidase inhibitory activity

Individual sample extracts of varying concentrations were made up to 1 mL with phosphate buffer (0.1M, pH 6.9) and 100 μ L of alpha-glucosidase enzyme (25 μ g of enzyme) was added. The contents were incubated at 25°C for 5 min. After pre-incubation, about 100 μ L of p-nitrophenyl- α -D-glucopyranoside (250 μ g/mL) added, contents mixed well and the reaction mixture incubated for 20 min at 25°C. The reaction was terminated by the addition of 1.0 mL of 1 N Na₂CO₃. After 10 min, the yellow color developed was measured at 405 nm and the α -glucosidase inhibition % (IC %) was calculated using the following equation:

$$IC\% = [(A_{control} - A_{sample}) / A_{control}] \times 100;$$

Where, A_{control} is the absorbance of the blank control (containing both enzyme and substrate except jamun seed extract): A_{sample} is the absorbance of the test sample.

A graph was plotted with concentration along x axis and IC% along y axis, and IC₅₀ values were calculated. IC₅₀ value signifies the concentration of tested samples to inhibit 50% of the α -glucosidase enzyme.

Statistical analysis

All the experiments were conducted on triplicate samples. The effect of different extraction solvents, time, and conditions on TPC, TFC, antioxidant potential, and alpha-glucosidase inhibitory activity was evaluated by ANOVA using SAS (SAS, 2011). Pearson's linear correlation was performed to measure the correlation and strength of the relationship between TPC, TFC, antioxidant potential, and alpha-glucosidase inhibitory activity of different solvent extracts.

RESULTS AND DISCUSSION

Effect of solvent type with varying polarity, condition, and extraction time on the extraction of total phenolics and total flavonoids from jamun seeds

Total phenolics (TPC), flavonoids (TFC), antioxidant potential, and alpha-glucosidase inhibitory activity of different solvent extracts of jamun seeds are given in Table 1.

Total phenolic content (TPC)

The TPC varied between 6.0 (mg/g) jamun seed powder for acetone to 119.2 (mg/g) for 80% aqueous acetone. Significant variation was observed for the yield of total phenolic content with respect to solvent type, extraction condition, and time. The phenolic extraction ability was found significantly higher for aqueous solvents compared to their absolute forms. Absolute forms of methanol, ethanol, and acetone extracted less amount of TPC compared to their aqueous counterparts irrespective of extraction time and conditions. Among the solvents with absolute form, extractability of phenolics was found more in methanol (69.8 mg/g) followed by ethanol (42.3 mg), while acetone extracted significantly less TPC (19.3 mg). Water alone extracted a significantly high amount of TPC (78.1 mg) compared to solvents with absolute form. The addition of water to the solvents significantly increased the extraction ability of the solvents. Among the aqueous forms of solvents, aqueous acetone extracted higher amount of TPC (111.9 mg) followed by aqueous methanol (89.8 mg) and aqueous ethanol (83.6 mg). About 5 to 10-fold increase in TPC content was observed for aqueous acetone with different conditions and extraction time compared to its absolute form. From the results, it is clear that the solvents with varying polarity had a significant effect on extractability, and aqueous acetone was superior among the solvent systems studied.

Earlier studies also reported similar findings with high extractability of phenolics by aqueous acetone with varying proportions starting from 50 %, 70%, and 80% (Zhao *et al.*, 2006; Sulaiman *et al.*, 2011, Wijekoon *et al.*, 2011). In the present study, two extraction conditions were employed, one at room temperature [25±2°C] and the other using ultrasonication maintained at 50±2°C. Ultrasonication extracted significantly high amount of TPC compared to room temperature irrespective of solvent type. Similarly, in most cases, 60 min extraction time significantly increased the extraction ability of solvents compared to 30 min extraction time. From the results, it is clear that the aqueous form of acetone with ultrasonication for 60 min is the ideal condition for the extraction of more TPC from jamun seed powder.

Table 1: Total phenolics, flavonoids, antioxidant potential and alpha-glucosidase inhibitory activity of different solvent extracts of jamun seeds

Solvents	Condition	Time (min)	TPC	TFC	FRAP	CUPRAC	DPPH	ABTS	α -glucosidase inhibitory activity*
Water	RT	30	53.0	4.00	0.51	2.06	0.62	0.58	285.2
		60	53.6	3.93	0.52	1.92	0.62	0.92	206.0
	Sonication	30	67.7	5.65	0.61	2.54	0.86	0.86	222.2
		60	78.1	5.47	0.83	2.74	0.89	1.06	197.0
Methanol	RT	30	50.2	6.68	0.55	2.00	0.62	0.55	120.9
		60	57.1	6.82	0.54	2.09	0.61	0.77	57.7
	Sonication	30	68.8	8.41	0.65	2.24	0.76	0.93	194.2
		60	69.8	8.13	0.75	2.64	0.77	0.96	111.2
Aqueous methanol (80% v/v)	RT	30	74.7	9.20	0.89	3.09	0.90	0.98	58.4
		60	76.7	8.69	0.80	2.90	0.92	0.98	90.1
	Sonication	30	89.8	10.81	0.72	3.56	0.99	1.17	107.5
		60	72.4	9.09	0.62	2.38	0.70	1.19	86.6
Ethanol	RT	30	16.5	2.59	0.18	0.62	0.21	0.12	313.7
		60	17.8	3.20	0.16	0.50	0.23	0.22	206.8
	Sonication	30	31.2	3.86	0.34	1.13	0.35	0.34	215.7
		60	42.3	5.31	0.44	1.50	0.52	0.58	173.8
Aqueous ethanol (80% v/v)	RT	30	63.8	7.13	0.64	2.59	0.80	1.02	93.1
		60	65.9	6.61	0.66	2.53	0.83	0.90	83.8
	Sonication	30	79.0	8.48	0.78	3.05	0.94	1.06	120.6
		60	83.6	8.33	0.94	3.16	0.94	1.08	78.8
Acetone	RT	30	7.0	1.50	0.10	0.30	0.06	0.06	972.7
		60	6.0	1.06	0.07	0.16	0.10	0.28	771.0
	Sonication	30	14.0	1.94	0.16	0.53	0.12	0.28	536.9
		60	19.3	3.98	0.21	0.58	0.20	0.35	268.7
Aqueous acetone (80% v/v)	RT	30	83.7	8.60	0.93	3.54	1.08	1.24	81.6
		60	94.4	7.44	0.95	3.70	1.11	1.12	58.0
	Sonication	30	97.2	8.98	0.88	3.85	1.08	1.35	54.0
		60	111.19	9.01	1.08	4.23	1.17	1.42	47.0
Interaction effect with CD									
Solvent		2.29	0.48	0.03	0.09	0.05	0.08	51.7	
Condition		1.23	0.26	0.02	0.05	0.03	0.04	27.7	
Time		1.23	NS	0.02	NS	NS	0.04	27.7	
Solvent * Condition		3.24	NS	0.05	0.13	0.07	0.11	73.2	
Solvent * Time		1.73	0.36	0.02	0.07	0.04	0.06	NS	
Condition * Time		NS	NS	0.05	0.13	NS	NS	NS	
Solvent * Condition * Time		4.58	0.96	0.07	0.18	0.10	NS	103.5	

Three independent experiments were performed and data are presented as mean per gram of dried jamun seed powder;

RT- Room temperature [$25 \pm 2^\circ\text{C}$]; TPC- Total phenolic content; TPC was expressed as GAE - Gallic acids equivalent (GAE); TFC- Total flavonoid content; TFC was expressed as QE - Quercetin equivalent.

DPPH and ABTS values are expressed as mM trolox equivalent/ g dry jamun seed powder, obtained from a trolox solution having a free radical scavenging activity (IC₅₀) equivalent to that of sample.

FRAP and CUPRAC values are expressed as mM trolox equivalent / g dry jamun seed powder, obtained from a trolox solution having reducing power equivalent to that of sample. CD- Critical Difference at 5% level of significance;

* α -glucosidase inhibitory activity was expressed as the amount of sample (in μg) required to inhibit 50% of the enzyme alpha-glucosidase (25 μg) activity (IC₅₀) in the presence of substrate p-nitro phenyl glucopyranoside (25 μg)

Total flavonoid content (TFC)

The TFC varied between 1.06 mg/g for acetone to 10.81 mg/g for 80% aqueous methanol. Significant variation was observed for the yield of TFC in terms of solvent type and extraction condition. Among the absolute solvents used, methanol was found to be a better solvent for TFC extraction (6.68-6.82 mg at room temperature and 8.13-8.41mg under ultra-sonication) followed by ethanol. Absolute acetone was found to be poor solvent that could extract only 1.06 to 1.50 mg of TFC at room temperature and 1.94-3.98 mg TFC under ultra-sonication. The addition of water to the absolute solvents significantly increased the TFC extraction ability. Among the aqueous forms, 80% methanol extracted the maximum amount of flavonoids (10.81 mg) followed by 80% acetone (9.01 mg). The increase in extraction ability was found high for acetone followed by ethanol and methanol. The extraction condition significantly increased the TFC, and ultra-sonication extracted a high amount of TFC compared to room temperature, and the effect was high for acetone. Extraction time did not vary significantly among the condition and solvent system used for extraction of flavonoids, except for acetone and methanol. In case of acetone, 60 min extraction time significantly increased the TFC content under ultra-sonication (1.94 mg to 3.98 mg), but in case of methanol, 60 min extraction time significantly reduced the TFC content from 9.20 to 8.69 mg and 10.8 to 9.09 mg for absolute methanol and 80% methanol, respectively). From the study, it was found the aqueous form of methanol with ultrasonication for 30 min is the ideal condition for extraction of more TFC from jamun seed powder.

Effect of solvent type with varying polarity, conditions and time on antioxidant potential and alpha-glucosidase inhibitory activity.

Antioxidant potential

Jamun seeds contain a wide array of polyphenolic compounds with different characteristics, thus employing a single method to evaluate the total antioxidant potential may not be appropriate to accurately measure the antioxidant potential. Thus,

in the present study four complementary methods *viz.* DPPH and ABTS radical scavenging activity, FRAP, and CUPRAC reducing power – based on the single electron transfer mechanism were tested to evaluate the antioxidant activity due to their simplicity, stability, and accuracy. The results obtained in the study were expressed as mM TE per g jamun seed powder (Table1). Significant differences were observed between various solvent systems and extraction conditions for antioxidant potential measured by various methods. The antioxidant potential measured by FRAP and ABTS method showed significant variation with respect to extraction time. Among the antioxidant methods employed, CUPRAC method measured higher antioxidant potential, followed by ABTS and DPPH. Correlation studies clearly showed that the total phenolics and flavonoids contents have a significant positive correlation with antioxidant potential [0.987**, 0.991**, 0.987**, 0.967** for TPC with FRAP, CUPRAC, DPPH and ABTS; and 0.884**, 0.891**, 0.881**, 0.882**for TFC with FRAP, CUPRAC, DPPH and ABTS, respectively, at the 0.01 level ($P < 0.01$)] (Fig. 1, 2). Since the antioxidant potential of the plant samples is directly related to the total polyphenolic content, the polarity of the solvents had a similar response to the total antioxidant potential. Among the studied solvents with different polarities, absolute methanol extract showed higher antioxidant potential measured by all methods followed by ethanol. Absolute acetone extract showed lesser antioxidant potential. Antioxidant potential due addition of water to the solvents had a similar effect observed for TPC. Aqueous solvent extracts had significantly higher amount of antioxidant potential compared to their counterpart with absolute forms.

Samples extracted using ultra-sonication had significantly higher amount of antioxidant potential compared to the samples extracted under room temperature. The time of extraction did not vary significantly for antioxidant potential measured by CUPRAC and DPPH, while FRAP and ABTS significantly varied with the time duration of extraction. In most cases, radical scavenging values obtained for ABTS radical scavenging




Fig. 1: Relationship between TPC and antioxidant potential.

**Correlation is significant at the 0.01 level ($P < 0.01$).




Fig. 2: Relationship between TFC and antioxidant potential.

**Correlation is significant at the 0.01 level ($P < 0.01$).

activity were significantly higher than DPPH radical scavenging activity. This could be because of the ABTS radical's solubility in hydrophilic and lipophilic solvents, and its activity in a wide range of pH (Cano *et al.*, 1998). In case of reducing power, CUPRAC method resulted in higher reducing power compared to FRAP, and it is mainly due to the pH of the reaction medium. In FRAP method, the reaction is carried out in acidic pH (3.6) whereas CUPRAC assay is carried out in neutral pH (7.0). Acidic pH may cause a reduction in reducing power due to protonation of antioxidant compounds, while in neutral pH and basic conditions, antioxidant potential increases due to proton dissociation (Apak *et al.*, 2004).

α -Glucosidase inhibitory activity

The α -glucosidase inhibitory action of the different solvent extract of jamun seed powder was assessed and the results are expressed as the amount of sample (in μg) required to inhibit 50% of the enzyme alpha-glucosidase (25 μg) activity (IC₅₀) in the presence of substrate p-nitrophenyl glucopyranoside (25 μg) (Table 1). Lesser the amount required to inhibit the α -glucosidase enzyme indicated the better inhibitory activity. As stated earlier, the α -glucosidase inhibitory activity is mainly due to the presence of polyphenolic compounds, the solvent system which extracted higher polyphenolic compounds showed better α -glucosidase inhibitory activity. Total phenolic content and flavonoid content negatively correlated [$r=-0.754^{**}$ and -0.784^{**} , at the 0.01 level ($P<0.01$), respectively] with the amount of jamun seed powder required to inhibit the α -glucosidase enzyme (Fig. 3, 4). Solvent types, extraction condition, and time duration for extraction significantly affected the α -glucosidase inhibitory activity of the extract. Among the absolute solvents, the amount of jamun seed powder extract required to inhibit the enzyme is lesser for methanol followed by ethanol irrespective of extraction condition and time duration of extraction. Significantly very high amount of jamun seed powder is required to inhibit the enzyme when

extracted with absolute acetone. Among the aqueous solvents, 80 % acetone showed better inhibitory activity compared to 80% methanol and 80% ethanol. Solvent with extraction condition and extraction condition with time duration did not vary significantly for α -glucosidase inhibitory activity, but compared to 30 min, 60 min extraction duration had better α -glucosidase inhibitory activity especially for 80 % acetone extract. From this assay, it was found the aqueous form of acetone is a better solvent for extraction of more α -glucosidase inhibitory compounds under ultrasonication for 60 min. Presence of significant amount of antioxidant compounds such as phenolic acids, and flavonoids attributed to jamun seed's medicinal values such as anti-diabetic properties (Ayya *et al.*, 2015). Laboratory studies revealed that the antidiabetic effect of jamun seeds are mainly due to their inhibitory activity on the major enzymes *viz* alpha-amylase and alpha-glucosidase which involves in hydrolysis of carbohydrate (Omar *et al.*, 2012). Kim *et al.*, (2016) stated that the phenolics can impact carbohydrate metabolism and help maintain blood glucose homeostasis through multiple pathways. Results from the present study corroborated the earlier reports that the increased amount of phenolics compounds with antioxidant potential can inhibit more amount of α -glucosidase, which helps in maintaining blood glucose homeostasis.

Polyphenolic compounds derived from plant sources gaining more interest among researchers, food manufacturers, and consumers due to their numerous health-benefits. Jamun seed powder is rich in polyphenolic compounds with both antioxidant and α -glucosidase inhibitory activity. The present study showed that the nature of solvent and extraction condition had a significant effect on TPC, TFC, antioxidant, and α -glucosidase inhibitory activity. The aqueous form of acetone (acetone: water 80:20, v/v) is the better solvent system for extraction of polyphenolic compounds with antioxidant and α -glucosidase inhibitory activity. Ultrasonication for 60 min facilitated higher extraction of phenolics.




Fig. 3: Relationship between total phenolic content and α-glucosidase inhibitory activity.

**Correlation is significant at the 0.01 level ($P < 0.01$).




Fig. 4: Relationship between total flavonoid content and α-glucosidase inhibitory activity.

**Correlation is significant at the 0.01 level ($P < 0.01$).

ACKNOWLEDGEMENT

This work was financially supported by ICAR, India (Project No. HORTIIHRCIL2015 (010) (02)). We are

grateful to the Director, ICAR-IIHR, Bengaluru, Karnataka, for constant support and encouragement in conducting this study.

REFERENCES

- Apak, R., Guclu, K., Ozyurek, M. and Karademir, S.E. 2004. Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *J. Agric. Food Chem.*, **52**: 7970-7981.
- Aqil, F., Gupta, A., Munagala, R., Jeyabalan, J., Kausar, H., Sharma, R. J., Singh, I. P. and Gupta, R.C. 2012. Antioxidant and antiproliferative activities of anthocyanin/ellagitannin-enriched extracts from *Syzygium cumini* L. (Jamun, the Indian Blackberry). *Nutr. Cancer.*, **64**(3): 428-438.
- Arivalagan, M., Roy, T. K., Yasmeen, A. M., Pavithra, K. C., Jwala, P. N., Shivasankara, K. S., Manikantan, M. R., Hebbar, K. B. and Kanade, S. R. 2018. Extraction of phenolic compounds with antioxidant potential from coconut (*Cocos nucifera* L.) testa and identification of phenolic acids and flavonoids using UPLC coupled with TQD-MS/MS. *LWT - Food Sci Technol.*, **92**: 116-126.
- Arnao, M.B., Cano, A. and Acosta, M. 2001. The hydrophilic and lipophilic contribution to total antioxidant activity. *Food Chem.*, **73**: 239-244.
- Ayya, N., Nalwade, V and Khan, T.N. 2015. Effect of jamun (*Syzygium cumini* L.) seed powder supplementation on blood glucose level of type-II diabetic subject. *Food Sci. Res. J.*, **6**(2): 353-356.
- Ayyanar, M. and Subash-babu, P. 2012. *Syzygium cumini* (L.) Skeels: a review of its phytochemical constituents and traditional uses. *Asian Pac. J. Trop. Biomed.*, **2**:240-246
- Benherlal, P. and Arumughan, C. 2007. Chemical composition and in vitro antioxidant studies on *Syzygium cumini* fruit. *J. Sci. Food Agric.*, **87**:2560-2569
- Benzie, I.E.F. and Strain J.J. 1996. The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': The FRAP assay. *Anal. Biochem.*, **239**: 70-76.
- Brand-Williams, W., Cuvelier, M.E., and Berset, C. 1995. Use of free radical method to evaluate antioxidant activity. *LWT - Food Sci Technol.*, **28**: 25-30.
- Cano, A., Hernandez-Ruiz, J., Garcia-Canovas, F., Acosta, M. and Arnao, M.B. 1998. An endpoint method for estimation of the total antioxidant activity in plant material. *Phytochem. Anal.*, **9**: 196-202.
- De Bona, K.S., Bonfanti, G., Bitencourt, P.E., da Silva, T.P., Borges, R.M., Boligon, A., Pigatto, A., Athayde, M.L., Moretto, M.B. 2016. Protective effect of gallic acid and *Syzygium cumini* extract against oxidative stress-induced cellular injury in human lymphocytes. *Drug Chem. Toxicol.*, **39**(3):256-263
- Hameed F., Gupta N., Rahman R., Anjum N. and Nayik G.A. 2020. Jamun. In: Nayik G.A., Gull A. (eds) Antioxidants in Fruits: Properties and Health Benefits. Springer, Singapore. https://doi.org/10.1007/978-981-15-7285-2_32
- Hayouni, A., Abedrabba, M., Bouix, M. and Hamdi, M. 2007. The effects of solvents and extraction method on the phenolic contents and biological activities *in vitro* of Tunisian *Quercus coccifera* L. and *Juniperus phoenicea* L. fruit extracts. *Food Chem.*, **105**: 1126-1134.
- Hossain, H., Rahman, S.E., Akbar, P.N., Khan, T.A., Rahman, M., Jahan, I.R. 2016. HPLC profiling, antioxidant and in vivo anti-inflammatory activity of the ethanol extract of *Syzygium jambos* available in Bangladesh. *BMC Res. Notes*, **9**:191-198.
- Kim, Y., Keogh, J. B., and Clifton, P.M. 2016. Polyphenols and glycemic control. *Nutrients*, **8**, E17.
- Khoddami, A., Meredith A.W. and Thomas H.R. 2013. Techniques for Analysis of Plant Phenolic Compounds. *Molecules*, **18**, 2328-2375

- Mahmood, N. 2016. A review of α -amylase inhibitors on weight loss and glycemetic control in pathological state such as obesity and diabetes. *Comp. Clin. Pathol.*, **25**: 1253–1264. <https://doi.org/10.1007/s00580-014-1967-x>
- Omar, R., Li, L., Yuan, T., and Seeram NP. 2012. α Glucosidase Inhibitory hydrolyzable tannins from *Eugenia jambolana* seeds. *J. Nat. Prod.*, **75**, 1505"1509
- SAS. 2011. Statistical Analysis Software System, Version 9.3. SAS Institute, Cary, NC, USA.
- Sim, L. Quezada-Calvillo, R. Sterchi, E.E. Nichols, B.L. Rose D.R. 2008. Human intestinal maltase–glucoamylase: crystal structure of the N-terminal catalytic subunit and basis of inhibition and substrate specificity *J. Mol. Biol.*, **375**(3): 782-792.
- Singleton, V.L. Orthofer, R., and Lamuela-Raventos R.M. 1999. Analysis of total phenols and otherOxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods Enzymol.*, **299**:265–275.
- Sulaiman, S.F., Sajak, A.A.B., Supriatno, K.L.O. and Seow, E.M. 2011. Effect of solvents in extracting polyphenols and antioxidants of selected raw vegetables. *J. Food Compos. Anal.*, **24**: 506–515.
- Sun, T. and Ho, H. 2005. Antioxidant activities of buckwheat extracts. *Food Chem.*, **90**: 743–749.
- Turkmen, N., Sari, F. and Velioglu, Y.S. 2006. Effects of extraction solvents on concentration and antioxidant activity of black and black mate tea polyphenols determined by ferrous tartrate and Folin-Ciocalteu methods. *Food Chem.*, **99**: 835–841.
- Wijekoon, M.M.J.O., Bhat, R. and Karim, A.A. 2011. Effect of extraction solvents on the phenolic compounds and antioxidant activities of bungakantan (*Etlingera elatior* Jack.) inflorescence. *J. Food Compos. Anal.*, **24**: 615–619
- Zhao, H., Dong, J., Lu, J., Chen, J., Li, Y., Shan, L., Lin, Y., Fan, W. and Gu, G. 2006. Effect of extraction solvent mixtures on antioxidant activity evaluation and their extraction capacity and selectivity for free phenolic compounds in barley (*Hordeum vulgare* L.). *J. Agric. Food Chem.*, **54**: 7277–7286.
- Zhishen, J., Mengcheng, T. and Jianming, W. 1999. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food chem.*, **64**: 555-559.

(Received on 9.04.2021, Revised on 25.05.2021, Accepted on 26.05.2021)