

JOURNAL OF HORTICULTURAL SCIENCES

Volume 16

December 2021

Issue 2





JOURNAL OF HORTICULTURAL SCIENCES

Volume 16

Issue 2

June 2021

CONTENTS

In this Issue

i-ii

Review

- Phytoremediation of indoor air pollutants: Harnessing the potential of plants beyond aesthetics** 131-143
Shalini Jhanji and U.K.Dhatt

Research Articles

- Response of fruit yield and quality to foliar application of micro-nutrients in lemon [*Citrus limon* (L.) Burm.] cv. Assam lemon** 144-151
Sheikh K.H.A., Singh B., Haokip S.W., Shankar K., Debbarma R.
- Studies on high density planting and nutrient requirement of banana in different states of India** 152-163
Debnath Sanjit Bauri F.K., Swain S., Patel A.N., Patel A.R., Shaikh N.B., Bhalerao V.P., Baruah K., Manju P.R., Suma A., Menon R., Gutam S. and P. Patil
- Mineral nutrient composition in leaf and root tissues of fifteen polyembryonic mango genotypes grown under varying levels of salinity** 164-176
Nimbolkar P.K., Kurian R.M., Varalakshmi L.R., Upreti K.K., Laxman R.H. and D. Kalaivanan
- Optimization of GA3 concentration for improved bunch and berry quality in grape cv. Crimson Seedless (*Vitis vinifera* L)** 177-184
Satisha J., Kumar Sampath P. and Upreti K.K.
- RGAP molecular marker for resistance against yellow mosaic disease in ridge gourd [*Luffa acutangula* (L.) Roxb.]** 185-192
Kaur M., Varalakshmi B., Kumar M., Lakshmana Reddy D.C., Mahesha B. and Pitchaimuthu M.
- Genetic divergence study in bitter melon (*Momordica charantia* L.)** 193-198
Nithinkumar K.R., Kumar J.S.A., Varalakshmi B, Mushrif S.K., Ramachandra R.K. , Prashanth S.J.
- Combining ability studies to develop superior hybrids in bell pepper (*Capsicum annuum* var. *grossum* L.)** 199-205
Varsha V., Smaranika Mishra, Lingaiah H.B., Venugopalan R., Rao K.V. Kattedgoudar J. and Madhavi Reddy K.
- SSR marker development in *Abelmoschus esculentus* (L.) Moench using transcriptome sequencing and genetic diversity studies** 206-214
Gayathri M., Pitchaimuthu M. and K.V. Ravishankar



Generation mean analysis of important yield traits in Bitter gourd (<i>Momordica charantia</i>)	215-221
Swamini Bhoi, Varalakshmi B., Rao E.S., Pitchaimuthu M. and Hima Bindu K.	
Influence of phenophase based irrigation and fertigation schedule on vegetative performance of chrysanthemum (<i>Dendranthema grandiflora</i> Tzelev.) var. Marigold	222-233
Vijayakumar S., Sujatha A. Nair, Nair A.K., Laxman R.H. and Kalaivanan D.	
Performance evaluation of double type tuberose IHR-4 (IC-0633777) for flower yield, quality and biotic stress response	234-240
Bharathi T.U., Meenakshi Srinivas, Umamaheswari R. and Sonavane, P.	
Anti-fungal activity of <i>Trichoderma atroviride</i> against <i>Fusarium oxysporum</i> f. sp. <i>Lycopersici</i> causing wilt disease of tomato	241-250
Yogalakshmi S., Thiruvudainambi S., Kalpana K., Thamizh Vendan R. and Oviya R.	
Seed transmission of bean common mosaic virus-blackeye cowpea mosaic strain (BCMV-BICM) threaten cowpea seed health in the Ashanti and Brong-Ahafo regions of Ghana	251-260
Adams F.K., Kumar P.L., Kwoseh C., Ogunsanya P., Akromah R. and Tetteh R.	
Effect of container size and types on the root phenotypic characters of <i>Capsicum</i>	261-270
Raviteja M.S.V., Laxman R.H., Rashmi K., Kannan S., Namratha M.R. and Madhavi Reddy K.	
Physio-morphological and mechanical properties of chillies for mechanical harvesting	271-279
Yella Swami C., Senthil Kumaran G., Naik R.K., Reddy B.S. and Rathina Kumari A.C.	
Assessment of soil and water quality status of rose growing areas of Rajasthan and Uttar Pradesh in India	280-286
Varalakshmi LR., Tejaswini P., Rajendiran S. and K.K. Upreti	
Qualitative and organoleptic evaluation of immature cashew kernels under storage	287-291
Sharon Jacob and Sobhana A.	
Physical quality of coffee bean (<i>Coffea arabica</i> L.) as affected by harvesting and drying methods	292-300
Chala T., Lamessa K. and Jalata Z	
Vegetative vigour, yield and field tolerance to leaf rust in four F1 hybrids of coffee (<i>Coffea arabica</i> L.) in India	301-308
Divya K. Das, Shivanna M.B. and Prakash N.S.	
Limonene extraction from the zest of <i>Citrus sinensis</i>, <i>Citrus limon</i>, <i>Vitis vinifera</i> and evaluation of its antimicrobial activity	309-314
Wani A.K., Singh R., Mir T.G. and Akhtar N.	
Event Report	315-318
National Horticultural Fair 2021 - A Success Story	
Dhananjaya M.V., Upreti K.K. and Dinesh M.R.	
Subject index	319-321
Author index	322-323

Original Research Paper

**Influence of phenophase based irrigation and fertigation schedule
on vegetative performance of chrysanthemum
(*Dendranthema grandiflora* Tzelev.) var. Marigold**

Vijayakumar S.¹, Sujatha Anil Nair¹, Nair A.K.², Laxman R.H.³ and Kalaivanan D.⁴

¹Division of Flower and Medicinal Crops, ²Division of Vegetable crops, ³Division of Basic sciences and
⁴Division of Natural Resources, ICAR – Indian institute of Horticultural Research, Bangalore, Karnataka

*Corresponding author Email : agrivijay483@gmail.com

ABSTRACT

The vegetative performance of chrysanthemum var. Marigold with respect to phenophase based irrigation and fertigation schedule was evaluated. In the vegetative phase, the maximum plant height (62.44 cm), number of secondary branches per plant (42.65), number of primary branches per plant (10.85), leaf area (3793.81 cm²) was recorded in the treatment combination. Whereas, the maximum average plant spread (47.98 cm) was in I₁F₄, number of leaves per plant (217.76) was in I₃F₁. Scheduling irrigation regime I₃-(0.8 ER each at vegetative, bud and flowering phases) in combination with weekly application of (F₄) 75:112.5:75 kg NPK/ha in three splits 40:20:20 % NPK (vegetative phase), 30:40:40 % NPK (bud phase) 30:40:40% NPK (flowering phase) through fertigation recorded maximum loose flower yield (26.27 t/ha) and this can be correlated with increased values for most of the vegetative parameters that directly influence the yield of the crop. Hence the above was observed best treatment over other treatment combinations with respect to vegetative parameters of chrysanthemum var. Marigold.

Key words: Chrysanthemum var. Marigold, fertigation, irrigation, phenophase and vegetative performance.

INTRODUCTION

Chrysanthemum (*Dendranthema grandiflora* Tzelev.) is one of the important commercial flower crops in India as well as in the world. It is native of the Northern hemisphere, chiefly Europe and Asia. It belongs to family Asteraceae and is commonly called as the “Queen of the East”. Its flowers are valued for its long keeping quality, wide array of colours and different forms, which make it suitable for use in floral bouquets, flower arrangements and decorations. Chrysanthemum is the second most important flower crop after rose in India. The area under flower crops is 339000 ha with an overall production of 19.91 lakh tonnes. The leading chrysanthemum growing state is Karnataka with an area of 5453 ha and production of 59.54 thousand tonnes of loose flowers in 2017-18 after Tamil Nadu. Water and fertilizer are the two vital inputs for crop production. Apart from the economic considerations, it is also well known that the injudicious use of water and fertilizer can have

far reaching deleterious implications on the environment. Therefore, the need arises for technological options, which will help in sustaining the precious resources and maximizing crop production without any pernicious impact on the environment. Optimum plant nutrition is very essential in plant growth and development, if it is not in sufficient amount then it reduces the vigor of the plant and affects yield of flower crops by producing small leaves, light green or off-color foliage, fewer branches and poor flowering (Melvin and James, 2001). Excessive application of nutrients can cause adverse effects on plant growth, increase the potential for environmental contamination through leaching and waste of resources. Method of nutrient application to plants is also a key issue to get the optimum potential of the crop. Fertigation helps in reducing the wastage of nutrients through enhanced use efficiency of fertilizer besides providing flexibility in timing of



fertilizer application in relation to crop demand based on phenological stages of growth (Papadopoulos, 1992). It also determines quantity of nutrients, timing of application and most important component of water distribution (Ahmad and Khan, 2017). The amount of nutrient and water requirement of a plant varies according to its phenophase and dispensation of water and nutrients can be scheduled accordingly. The fertigation scheduling should be based on plant, soil-air, plant water relations and growth stage of plant (Sankaranarayanan, 2007).

It is essential to work out an economically feasible and technologically efficient fertigation scheduling for optimum use of water and nutrients for enhanced water productivity with reference to different growth and developmental stages. Hence, it is important to evaluate under phenophase based irrigation and fertigation treatments for improving vegetative performance of chrysanthemum var. Marigold under open field condition.

MATERIAL AND METHODS

The present investigation conducted during two seasons *i.e.* 2018 & 2019, at the Division of Flowers and Medicinal Crops, ICAR-Indian Institute of Horticultural Research (ICAR-IIHR), Bengaluru. The experimental site is situated in eastern dry zone of Karnataka state at 13° 7' north latitude, 77° 29' east longitudes and at an altitude of 890 meters above the mean sea level. The experiment was laid out in split plot design with fifteen treatment combinations along with three replications. The treatment consists of three main plot treatments at phenophases of vegetative phase *i.e.* I₁ – (0.8, 1.0 and 1.2 ER at vegetative, bud and flowering phases, respectively), I₂ - (0.6, 0.8 and 1.0 ER at vegetative, bud and flowering phases, respectively) and I₃- (0.8 ER each at vegetative, bud and flowering phases) and five sub plot treatments (F₁: 33.3:33.3:33.3 % NPK (vegetative phase), 33.3:33.3:33.3 % NPK (Bud phase) 33.3:33.3:33.3 % NPK (Flowering phase) @ 100:150:100 Kg NPK/ha (RDF), F₂: 40:20:20 % NPK (vegetative phase), 30:40:40 % NPK (Bud phase) 30:40:40% NPK (Flowering phase) @ 100:150:100 Kg NPK/ha (RDF), F₃: 33.3:33.3:33.3 % NPK (vegetative phase), 33.3:33.3:33.3 % NPK (Bud phase) 33.3:33.3:33.3 % NPK (Flowering phase @ 75:112.5:75 Kg NPK/ha (75% RDF), F₄: 40:20:20 % NPK (vegetative

phase), 30:40:40 % NPK (Bud phase) 30:40:40% NPK (Flowering phase) @ 75:112.5:75 Kg NPK/ha (75% RDF), F₅: Soil application of recommended dose of fertilizer (100:150:100 Kg NPK/ha) and F₁-F₄: 25% of fertilizer dose *i.e.* 100:150:100 and 75:112.5:75 kg NPK/ha was applied as basal dose. The previous day open pan evaporimeter observation was considered for scheduling the irrigation as per the treatment. The

Evaporation replenishment (ER) =

$$\frac{\text{Bed Size (m}^2\text{)} \times \text{pan evaporation rate (mm)}}{\text{Discharge capacity of drip per minute (ml)}}$$

irrigation schedule was calculated by using following formula.

The organic manure *i.e.* farmyard manure (20 t/ha) and basal application (Urea, DAP and MOP) was applied as per the treatments as earlier to transplanting. Transplanting was followed with a spacing of 60 cm × 45 cm. The dose of fertilizers was applied based on treatments through fertigation in the form of water-soluble fertilizers (Urea, MAP and SOP). The fertigation was given at weekly intervals from thirty days after transplanting to 120 days.

RESULTS AND DISCUSSION

The vegetative parameters *viz.*, plant height (cm), number of primary and secondary branches per plant, average plant spread (cm) at flowering and leaf area (cm²) as influenced by phenophase based different irrigation and fertigation regimes are discussed below.

The plant height (cm) of chrysanthemum was significantly influenced by different levels of phenophase based irrigation and fertigation. Among interactions effects the maximum plant height (61.19 cm) was recorded in I₃F₄ and it was on par with I₂F₄ (59.19 cm) and I₂F₃ (59.10 cm) whereas, the minimum (41.10 cm) was recorded in the treatment combination I₂F₂ during the first year. The maximum plant height (65.30 cm), was recorded in I₃F₁ and it was on par with the treatments, I₁F₄ (64.50 cm), I₂F₄ (64.43 cm) and I₃F₄ (63.68 cm) whereas, the minimum (44.60 cm) was recorded in I₁F₂ during the second year. In pooled interaction, the maximum plant height (62.44 cm) was recorded in I₃F₄ and it was on par with the treatment I₂F₄ (61.81 cm) and the minimum (46.91 cm) was recorded in I₁F₂ (Table 1 & 2) (Fig.1).

Table 1. Influence of phenophase based irrigation and fertigation scheduling on plant height (cm) and number of primary branches of chrysanthemum var. Marigold

Treatments	Plant height (cm)			Number of primary branches per plant		
	I year	II year	Pooled mean	I year	II year	Pooled mean
I ₁	51.42	54.48	52.95	9.64	9.96	9.80
I ₂	52.32	56.30	54.31	9.74	9.14	9.44
I ₃	48.88	58.97	53.92	9.43	9.20	9.32
SE. d	0.65	0.40	0.38	0.03	0.08	0.05
CD (P=0.05)	1.83	1.11	1.07	0.10	0.23	0.14
F ₁	51.70	57.50	54.60	8.71	9.30	9.00
F ₂	44.14	52.40	48.27	9.77	9.50	9.63
F ₃	55.33	54.76	55.04	10.13	9.57	9.85
F ₄	58.83	64.20	61.52	10.61	10.83	10.72
F ₅	44.36	54.05	49.21	8.80	7.96	8.38
SE. d	0.66	0.58	0.40	0.11	0.11	0.08
CD (P=0.05)	1.14	1.20	0.83	0.22	0.23	0.17

Table 2. Interaction effect of phenophase based irrigation and fertigation scheduling on plant height (cm) of chrysanthemum var. Marigold

Treatments	I year					II year					Pooled Mean							
	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean
I ₁	51.90	49.21	56.79	56.10	43.09	51.42	53.71	44.60	55.60	64.50	54.00	54.48	52.81	46.91	56.20	60.30	48.55	52.95
I ₂	55.10	41.10	59.10	59.19	47.10	52.32	53.50	56.70	51.27	64.43	55.60	56.30	54.30	48.90	55.19	61.81	51.35	54.31
I ₃	48.10	42.10	50.10	61.19	42.90	48.88	65.30	55.90	57.40	63.68	52.56	58.97	56.70	49.00	53.75	62.44	47.73	53.92
Mean	51.70	44.14	55.33	58.83	44.36		57.50	52.40	54.76	64.20	54.05		54.60	48.27	55.04	61.52	49.21	
SE. d	CD (P=0.05)					SE. d	CD (P=0.05)					SE. d	CD (P=0.05)					
I	0.65		1.83			0.40					1.11							1.07
F	0.66		1.14			0.58					1.20							0.83
I at F	1.22		2.77			0.99					2.15							1.66
F at I	1.14		2.37			1.01					2.09							1.43

Table 3. Interaction effect of phenophase based irrigation and fertigation scheduling on number of primary branches per plant of chrysanthemum var. Marigold

Treatments	I year						II year						Pooled Mean					
	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean
I ₁	8.79	9.20	10.80	10.50	8.90	9.64	10.60	9.80	10.20	11.20	8.00	9.96	9.70	9.50	10.50	10.85	8.45	9.80
I ₂	8.93	10.40	10.00	10.77	8.60	9.74	8.10	9.60	9.60	10.19	8.19	9.14	8.52	10.00	9.80	10.48	8.40	9.44
I ₃	8.40	9.70	9.60	10.56	8.89	9.43	9.20	9.10	8.90	11.10	7.70	9.20	8.80	9.40	9.25	10.83	8.30	9.32
Mean	8.71	9.77	10.13	10.61	8.80		9.30	9.50	9.57	10.83	7.96		9.00	9.63	9.85	10.72	8.38	
	SE. d						SE. d						SE. d					
	CD (P=0.05)						CD (P=0.05)						CD (P=0.05)					
I	0.10						0.08						0.14					
F	0.22						0.11						0.17					
I at F	0.36						0.19						0.30					
F at I	0.39						0.19						0.30					




Fig. 1. Influence of phenophase based irrigation and fertigation scheduling on plant height (cm)

The irrigation treatment I_3 - (0.8 ER each at vegetative, bud and flowering phases) in combination with F_4 fertigation at 40:20:20 % NPK (vegetative phase), 30:40:40 % NPK (bud phase) 30:40:40% NPK (flowering phase) @ 75:112.5:75 kg NPK/ha² recorded the maximum plant height (62.44 cm) in chrysanthemum var. Marigold. The increase in plant height with irrigation at I_3 might be due to adequate moisture provided in the soil throughout the crop period. Adequate soil moisture resulted in greater development of meristematic tissues leading to higher rate of photosynthesis and assimilation in the plant system in marigold (Chawla, 2008).

In the fertigation treatment F_4 , higher proportion of nitrogen fertilizer at vegetative phase might have increased the plant height because of the synergistic interaction of nitrogen with available endogenous auxin resulting in enhanced cell wall plasticity and increased cell elongation thus resulting in increase in the height of the plant. Further, during the bud and flowering phases, the sustained growth of the plant might have been the result of optimum application of nitrogen. The results from the present investigation could hence be attributed to the frequent and constant application of optimum levels of fertilizers at appropriate intervals at crop phenophases, which increases the available nutrient status in the root rhizosphere at constant levels during all the phases thus increasing the uptake of nutrients rapidly, and further influencing the growth of the plant. Similar observations were earlier reported by Mamata *et al.* (2017) in marigold, Parya *et al.* (2017) in gerbera, Priyanka *et al.* (2017) in gladiolus and Satapathy *et al.* (2016) in marigold, Jamil *et al.* (2016), Zawadziska and Janicka (2007) in amaryllis and viola respectively.

The treatment I_1F_4 was on par with I_3F_4 for maximum (10.83), number of primary branches per plant (Table 1 & 3) and the maximum number of secondary branches per plant (42.65) was recorded in the treatment combination I_3F_4 and it was on par with I_1F_4 (41.44) and the minimum (17.75) was recorded in I_1F_5 . The treatment I_3F_4 recorded the maximum number of secondary branches per plant (42.65) in chrysanthemum var. Marigold. This increase in number branches might be mainly due to the increased irrigation scheduled favoring longer availability of soil moisture which leads to better growth and development of vegetative part of the plant. The greater availability

of nutrient at optimum proportions at critical growth stages in the present fertigation treatment might have resulted in production of more number of branches per plant as observed by Siraj Ali (1998) in bird-of-paradise. Polara *et al.* (2015) recorded similar results in African marigold. These findings are in conformation with the earlier results of Jawaharlal and Ganesh (2020) in chrysanthemum and Nagaraju *et al.* (2003) in rose (Table 4 & 5).

The average plant spread was significantly influenced and showed linear increase with irrigation regime and with optimum dosage of water-soluble fertilizers through fertigation. Among interactions effect the maximum average plant spread (53.23 cm) was recorded in the treatment combination I_1F_4 followed by the treatment I_1F_3 (45.76 cm) and the minimum (31.60 cm) was recorded in the treatment combination of I_1F_5 during the first year. The maximum average plant spread (49.33 cm) was recorded in the treatment combination I_3F_1 followed by I_2F_3 (44.87 cm) and the minimum (30.80 cm) was recorded in the treatment combination I_1F_2 during the second year. In pooled interaction, the maximum average plant spread (47.98 cm) was recorded in the treatment combination I_1F_4 followed by the treatment I_1F_3 (43.61 cm) and the minimum (32.23 cm) was recorded in the treatment combination of I_3F_2 (Table 4 & 6).

It was recorded that irrigation regime I_1 - (0.8, 1.0 and 1.2 ER at vegetative, bud and flowering phases, respectively) in combination with fertigation at 40:20:20 % NPK (vegetative phase), 30:40:40 % NPK (bud phase) 30:40:40% NPK (flowering phase) @ 75:112.5:75 kg NPK/ha registered maximum average plant spread (47.98 cm). This result clearly showed that higher amount of nitrogen supplied at vegetative phase along with higher soil moisture levels leads to increased vegetative growth of chrysanthemum var. Marigold. According to Paul *et al.* (1996) the plant spread could be attributed to the frequent application of fertilizers with constant supply of nutrients, at regular intervals for better growth which would have resulted in reduced nutrient losses by leaching and efficient use of nutrients through fertigation compared to soil application. This is in accordance with the findings of Deshmukh and Wavhal (1998) in china aster and Ahirwal *et al.* (2012) in African marigold.

The maximum number of leaves (235.03) was recorded in the treatment combination I_1F_4 and it was

Table 4. Influence of phenophase based irrigation and fertigation scheduling on vegetative parameters of chrysanthemum var. Marigold

Treatments	Number of secondary branches per plant			Average plant spread (cm)			Number of leaves per plant		
	I year	II year	Pooled mean	I year	II year	Pooled mean	I year	II year	Pooled mean
I ₁	29.18	30.65	29.91	42.32	39.67	41.00	221.93	136.89	179.40
I ₂	32.42	27.27	29.85	36.71	41.04	38.88	220.26	141.43	180.82
I ₃	30.97	29.29	30.13	35.69	40.48	37.43	218.84	156.34	187.59
SE. d	0.78	0.61	0.06	0.34	0.13	0.58	1.23	1.99	3.20
CD (P=0.05)	1.41	1.20	0.12	0.95	0.26	1.62	2.60	4.02	6.98
F ₁	30.06	34.06	32.06	37.13	43.87	40.50	224.07	159.64	191.86
F ₂	26.32	23.44	24.88	38.26	35.26	36.76	220.74	136.21	178.47
F ₃	31.50	27.30	29.40	40.54	40.92	40.73	214.97	143.18	178.57
F ₄	42.12	39.86	40.99	42.98	41.81	42.39	225.88	154.83	190.36
F ₅	24.30	20.68	22.49	34.30	38.45	36.37	217.97	130.57	173.77
SE. d	0.89	0.60	0.55	0.53	0.96	0.79	0.26	0.27	0.05
CD (P=0.05)	1.54	1.19	1.02	1.10	2.03	1.64	0.45	0.55	0.10

Table 5. Interaction effect of phenophase based irrigation and fertigation scheduling on number of secondary branches per plant of chrysanthemum var. Marigold

Treatments	I year						II year						Pooled Mean					
	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean
I ₁	26.82	26.59	31.78	42.04	18.69	29.18	37.24	28.42	29.93	40.84	16.81	30.65	32.03	27.50	30.85	41.44	17.75	29.91
I ₂	36.48	27.05	31.99	39.91	26.67	32.42	29.15	29.17	17.28	37.84	22.91	27.27	32.82	28.11	24.64	38.88	24.79	29.85
I ₃	26.88	25.31	30.72	44.40	27.55	30.97	35.78	12.74	34.70	40.89	22.33	29.29	31.33	19.02	32.71	42.65	24.94	30.13
Mean	30.06	26.32	31.50	42.12	24.30		34.06	23.44	27.30	39.86	20.68		32.06	24.88	29.40	40.99	22.49	
	SE. d						SE. d						SE. d					
	CD (P=0.05)						CD (P=0.05)						CD (P=0.05)					
I	0.78				1.41		0.61				1.20		0.06					0.17
F	0.89				1.54		0.60				1.19		0.55					1.02
I at F	1.34				2.68		1.01				2.07		0.89					1.76
F at I	1.33				2.66		1.00				2.06		0.90					1.75

Table 6. Interaction effect of phenophase based irrigation and fertigation scheduling on average plant spread (cm) of chrysanthemum var. Marigold

Treatments	I year						II year						Pooled Mean					
	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean
I ₁	38.40	42.63	45.76	53.23	31.60	42.32	40.46	30.80	41.46	42.73	42.90	39.67	39.43	36.71	43.61	47.98	37.25	41.00
I ₂	37.50	35.23	38.73	37.20	34.90	36.71	41.82	41.47	44.87	39.23	37.83	41.04	39.66	38.35	41.80	38.21	36.36	38.88
I ₃	35.50	30.93	37.13	38.50	36.40	35.69	49.33	33.53	36.43	43.47	33.10	40.48	42.42	32.23	36.78	40.99	34.75	37.43
Mean	37.13	38.26	40.54	42.98	34.30		43.87	35.27	40.92	41.81	38.45		40.50	36.76	40.73	42.39	36.38	
	SE. d						SE. d						SE. d					
	CD (P=0.05)						CD (P=0.05)						CD (P=0.05)					
I	0.34				0.95		0.13				0.26		0.58					1.62
F	0.53				1.10		0.96				2.03		0.79					1.64
I at F	0.89				1.93		2.53				5.59		1.36					2.99
F at I	0.92				1.90		2.54				5.24		1.37					2.84

on par with I_1F_1 (229.61) and the minimum number of leaves per plant (205.01) were recorded in I_1F_5 during the first year. The maximum number of leaves per plant (215.50) was recorded in the treatment combination I_3F_1 and it was on par with I_2F_2 (192.21), I_1F_3 (171.61) and I_3F_4 (175.90) whereas, the minimum (89.61) was recorded in I_1F_2 during the second year. In pooled interaction the maximum number of leaves per plant (217.76) were recorded in the treatment combination I_3F_1 and it was on par with I_2F_2 (208.41),

I_1F_3 (195.96), I_1F_4 (197.22) and I_3F_4 (198.75) whereas, the minimum (154.61) was recorded in I_1F_2 (Table 4 & 7).

The treatment I_3F_4 registered maximum number of leaves per plant and maximum leaf area (2404.74 cm²) was recorded in I_1F_4 and it was on par with I_3F_4 (2352.18 cm²) and the lowest (1308.31 cm²) was recorded in I_3F_1 during the vegetative phase (Tables 8 & 9) (Fig. 2a, 2b & 2c). In the present study, the increase in number of leaves and leaf area could be

Fig. 2. Influence of phenophase based irrigation and fertigation scheduling on leaf area (cm²) at vegetative phase




Fig. 2.a. Influence of phenophase based irrigation and fertigation scheduling on leaf area (cm²) at vegetative phase during first year




Fig. 2.b. Influence of phenophase based irrigation and fertigation scheduling on leaf area (cm²) at vegetative phase during second year




Fig. 2.c. Pooled influence of phenophase based irrigation and fertigation scheduling on leaf area (cm²) at vegetative phase

Table 7. Interaction effect of phenophase based irrigation and fertigation scheduling on number of leaves per plant of chrysanthemum var. Marigold

Treatments	Leaf area (cm ²) at vegetative phase																	
	I year					II year					Pooled Mean							
	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean
I ₁	229.61	219.60	220.31	235.03	205.01	221.93	124.21	89.61	171.61	159.40	139.60	136.89	176.91	154.61	195.96	197.22	172.31	179.40
I ₂	222.60	224.61	208.60	221.00	224.30	220.26	139.21	192.21	104.21	129.20	142.30	141.43	180.91	208.41	156.41	175.10	183.30	180.82
I ₃	220.01	218.00	213.00	221.60	221.60	218.84	215.50	126.80	153.71	175.90	109.80	156.34	217.76	172.40	183.36	198.75	165.70	187.59
Mean	224.07	220.74	214.97	225.88	217.97		159.64	136.21	143.18	154.83	130.57		191.86	178.47	178.57	190.36	173.77	
	SE. d					CD (P=0.05)	SE. d					CD (P=0.05)	SE. d					CD (P=0.05)
I	1.23					2.60	1.99					4.02	3.20					6.98
F	0.26					0.45	0.27					0.55	0.05					0.10
I at F	4.27					9.08	23.26					52.41	12.39					27.79
F at I	4.58					9.45	22.28					45.98	11.99					24.76

Table 8. Influence of phenophase based irrigation and fertigation scheduling on leaf area (cm²) at vegetative phase of chrysanthemum var. Marigold

Treatments	I year	II year	Pooled mean
I ₁	1640.80	2373.74	2007.27
I ₂	1286.42	2362.97	1824.68
I ₃	1345.88	2575.28	1960.58
SE. d	8.89	45.52	5.30
CD (P=0.05)	24.68	101.36	10.50
F ₁	1102.67	2319.68	1711.18
F ₂	1334.65	2408.43	1871.54
F ₃	1569.73	2388.75	1979.24
F ₄	1778.79	2851.51	2315.15
F ₅	1336.00	2218.27	1777.14
SE. d	20.81	26.44	135.35
CD (P=0.05)	42.96	59.02	279.33

Table 9. Interaction effect of phenophase based irrigation and fertigation scheduling on leaf area (cm²) at vegetative phase of chrysanthemum var. Marigold

Treat-ments	I year						II year						Pooled Mean					
	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean	F ₁	F ₂	F ₃	F ₄	F ₅	Mean
I ₁	1430.34	1698.00	1794.19	1896.34	1385.15	1640.80	2792.27	2198.03	1857.80	2913.13	2107.47	2373.74	2111.31	1948.02	1826.00	2404.74	1746.31	2007.27
I ₂	897.36	1154.90	1145.80	1986.50	1247.54	1286.42	2530.47	2118.10	2794.03	2390.57	1981.67	2362.97	1713.92	1636.50	1969.92	2188.54	1614.61	1824.68
I ₃	980.32	1151.04	1769.20	1453.53	1375.31	1345.88	1636.30	2909.17	2514.43	3250.83	2565.70	2575.28	1308.31	2030.11	2141.82	2352.18	1970.51	1960.58
Mean	1102.67	1334.65	1569.73	1778.79	1336.00		2319.68	2408.43	2388.75	2851.51	2218.27		1711.18	1871.54	1979.24	2315.15	1777.14	
	SE.d						SE.d						SE.d					
	CD (P=0.05)						CD (P=0.05)						CD (P=0.05)					
I	8.89						45.52						5.30					
F	20.81						26.44						135.35					
I at F	33.45						37.86						23.49					
F at I	36.05						35.78						23.51					

attributed to Application of higher proportion of nitrogen fertilizer and optimum irrigation regimes at vegetative phase might have increased the number of leaves and leaf area. It may be due to the fact that the vegetative growth increased with nitrogen application and hence nitrogen is an essential part of nucleic acid, which plays a vital role in promoting vegetative growth. The present results were also in line with the reports of Maharnor *et al.* (2011) and Polara *et al.* (2014) in African marigold, Karam *et al.* (2007) in sunflower and Jaleel *et al.* (2009) in *Catharanthus*. Rawat and Mathpal (1984), Paul *et al.* (1996) and Khan *et al.* (1996) in various crops.

CONCLUSION

In the vegetative phase of chrysanthemum var. Marigold, the irrigation treatment I₃-(0.8 ER each at

vegetative, bud and flowering phases) in combination with fertigation treatment F₄ at 40:20:20 % NPK (vegetative phase), 30:40:40 % NPK (bud phase) 30:40:40% NPK (flowering phase) @ 75:112.5:75 kg NPK/ha was found adequate to cater the demand of water as well as nutrient requirement for vegetative phase of chrysanthemum var. Marigold. This can be correlated with the maximum loose flower yield (26.27 t/ha) registered by the same treatment. Further better plant growth as recorded during the investigation is indicative of better uptake of nutrients which in turn are involved in basic reaction of photosynthesis and in synthesis of metabolites required for plant growth with above irrigation and fertigation schedule. Hence it is concluded that the above treatment combination I₃F₄ was registered as the best treatment to improve the vegetative growth of chrysanthemum var. Marigold.

REFERENCES

- Ahirwar, M.K., Ahirwar, K. and Megha, Sukla. 2012. Effect of plant densities, nitrogen and phosphorus levels on growth, yield and quality of African marigold. *Annals of Plant and Soil Research*, **14(2)**:153- 155.
- Ahmad, A., and Khan, S. 2017. Water and energy scarcity for agriculture: Is irrigation modernization the answer. *Irrigation and Drainage*, **66**: 34-44.
- Chawla, S.L. 2008. Effect of irrigation regimes and mulching on vegetative growth, quality and yield of flowers of African marigold. Thesis, Doctor of Philosophy in Horticulture, Maharana Pratap University of Agriculture and Technology, Udaipur.
- Deshmukh, A.S., Shinde, P.P. and Jadhav, S.B. 1996. Fertigation under drip irrigation for sugarcane. **In:** All India Seminar on Modern Irrigation Techniques, Proceedings (June). pp. 217-219.
- Deshmukh, R. and Wahal, N. 1998. Effect of iron on growth and flowering of aster. *Journal of Maharashtra Agricultural University*, **23(2)**: 99-101.
- Hatwar, G.P. Gondane, S. V. Urkude, S.M. and Gahukar, O.V. 2003. Effect of micronutrients on growth and yield of chilli. *Soil and Crops*, **13**:123-1254.
- Jaleel, C.A., P. Manivannan, B. Sankar, A. Kishorekumar, R. Gopi, R. Somasundaram and R. Panneerselvam. 2007. Induction of drought stress tolerance by ketoconazole in *Catharanthus roseus* is mediated by enhanced antioxidant potentials and secondary metabolite accumulation. *Colloids Surf. B: Biointerfaces*, **60**: 201-206
- Jamil, M. K. J., Ahman, M. M. I. R., & Ossain, M. M. O. H. 2016. Response of N, P and K On The Growth and Flowering of Hippeastrum (*Hippeastrum hybridum* Hort.). *Journal of Agriculture*, **41(1)**, 91–101.
- Jawaharlal, M and Ganesh, S. 2020. Studies on the effect of fertigation in greenhouse chrysanthemum. *Journal of Pharmacognosy and Phytochemistry*, Sp.9(2): 254-259.
- Karam, F., Lahoud, R., Masaad, R., Kabalan, R., Breidi, J., Chalita, C., Roupahel, Y. 2007. Evapotranspiration, seed yield and water use efficiency of drip irrigated sunflower under full and deficit irrigation conditions. *Agricultural Water Management* **90(3)**, 213– 223.
- Kaushik, H., Singh, J.P., Mohan, B., Rajbeer and Nathiram. 2013. Effect of inorganic fertilize (nitrogen) and bio-fertilizer (*Azospirillum*) on growth and flowering in African marigold (*Tagetes erecta* L.) cv. Pusa Narangi Gainda. *International Journal of Agricultural Sciences*, **9(1)**:189-192.
- Khan, M.M., Sujatha, K., Krishna, R., Manohar, M., Kariyanna, A. A., Farooqui and Shivshankar.

1996. Fertigation studies in Horticultural crops In: Proc. All India Seminar on MIT, Bangalore, Ed Khan, M. M., pp. 178-186.
- Maharnor, S.I., Chopde, N., Thakre, S. and Raut, P.D. 2011. Effect of nitrogen and pinching on growth and yield of African marigold. *Asian Journal of Horticulture*, **6(1)**: 43-45.
- Mamta Kumawat, S.K., Khandelwal, M.R., Choudhary, P.K., Kumawat, G. Sharma and Paru Panwar. 2017. Effect of integrated nutrient management on growth, flowering and yield of African marigold (*Tagetes erecta* L.). *International Journal of Current Microbiology and Applied Sciences*, **6(8)**: 60-65.
- Marchner, H. 1983. Introduction to the mineral nutrition of plants. *Handbook Plant Physiology*, **154**: 31 - 38.
- Marschner, H. and Cakmak, I. 1986. Mechanism of phosphorus-induced zinc deficiency in cotton. II. Evidence for impaired shoot control of phosphorus uptake and translocation under zinc deficiency. *Physiologia Plantarum*, **68**: 491-496.
- Melvin, R.K. James, J.K. 2001. Fertilizing shade and ornamental trees, MSU Extension Forestry Bulletin, Bozeman.
- Nagaraju, C.G., Reddy, T.V. and Madaiah, D. 2003. Effect of plant density, irrigation and oil cakes on growth, production and quality of field grown rose cultivar Landora. *Journal of Ornamental Horticulture*, **6(3)**: 172-179.
- Pafli, G. 1965. Relations between abundant N supply and amino acid concentration on leaves of rice plants. *Plant and Soil*, **23**: 275- 284.
- Papadopoulos, I. 1992. Phosphorus fertigation of trickle-irrigated potato. *Fertilizer Research*, **31**: 9-13.
- Parya, C. 2017. Effect of integrated plant nutrient system for gerbera flower production under protected cultivation. *Journal of Applied Horticulture*, **19(2)**, 139-142.
- Paul, J., Joseph, J. and Kabeer, A. 1996. Fertilizer irrigation - an overview. In proceedings of All India Seminar on MIT, Bangalore, 196-204.
- Paul, J., Joseph, J. and Kabeer, A. 1996. Fertilizer irrigation - an overview. In proceedings of All India Seminar on MIT, Bangalore, 196-204.
- Polara, N.D., Gajipara, N.N. and Barad, A.V. 2014. Effect of nitrogen and phosphorus on nutrient content and uptake in different varieties of African marigold (*Tagetes erecta* L.). *International Quarterly Journal of Life Sciences*, **9(1)**:115-119.
- Priyanka Tirkey, Lagamanna R Kullur and VM Prasad. 2017. Effect of organic and Inorganic source of N.P.K on growth and yield parameters of gladiolus (*Gladiolus grandiflorus*) cv. Jester. *Journal of Pharmacognosy and Phytochemistry*, **6(5)**: 1004-1006.
- Rawat, P.S. and Mathpal, K.N. 1984. Effect of micronutrients on yield and sugar metabolism of some of the vegetables under Kumaon hill conditions. *Scientific Culture*, **50**: 243-244.
- Sankaranarayanan. 2007. Integrated water management system for better fibre quality and high production. TMC Annual Report, 2006-2007.
- Satapathy, S.P., Toppo, R., Dishri, M., and Mohanty, C.R. 2016. Impact of integrated nutrient management (INM) on flowering and corm production in gladiolus. *Biometrics & Biostatistics International Journal*, **4(7)**: 296 298.
- Singatkar, S.S., Swant, R.B., Ranpise, S.A. and Wavhal, K.N. 1995. Effects of different levels of N, P and K on growth and flower production of gaillardia. *Journal of Maharashtra Agriculture University*, **20(3)**: 392-394.
- Siraj Ali, M.S. 1998. Effect of singral and nitrophoska fertilizers on growth, flowering and mineral composition of bird-of-paradise (*Strelitzia reginae* Ait.) plants. *Indian Journal of Horticulture*, **55(3)**: 257-262.
- Terangpi, H. and Paswan, L. 2003. Effect of NPK on growth and flowering of Gerbera. *Journal of Ornamental Horticulture*. **6(1)**: 71-72.
- Zawadziska, A., & Janicka, D. 2007. Effects Of Compost Media On Growth and Flowering Of Parviflorous Garden Pansy (*Viola Wittrockiana* Gams. *Acta Agrobotanica*, **60(2)**: 161-166.

(Received on 05.09.2021, Revised on 23.09.2021 and Accepted on 15.01.2022)