

JOURNAL OF HORTICULTURAL SCIENCES

Volume 15

December 2020

Number 2



Conserving Honey Bees with Forage Plant Mexican Creeper - *Antigonon leptopus*



Society for Promotion of Horticulture
ICAR - Indian Institute of Horticultural Research, Bengaluru - 560 089



JOURNAL OF HORTICULTURAL SCIENCES

(Founded in 2005 by the Society for Promotion of Horticulture, Bengaluru, India)

Email : chiefeditor.jhs@gmail.com Webpage : <https://jhs.iihr.res.in/index.php/jhs>

Editor-in-Chief

Dr. S. Sriram

Editors

Dr. K. Himabindu

Dr. G. Senthilkumaran

Dr. Tejaswini Prakash

Dr. M. Manamohan

Dr. Anil Kumar Nair

Dr. J. Satisha

Dr. P. Venkata Rami Reddy

Dr. I.M. Doreyappa Gowda

Dr. R.H. Laxman

Dr. G.C. Sathisha

Editorial Advisory Board

International Editorial Advisory Board

Dr. Nanthi S. Bolan, Australia

Dr. Rod Drew, Australia

Dr. J. Mithila, USA

Dr. Claus Helmut Franz Orth, South Africa

Dr. Ilan Paran, Israel

Dr. Gi-Cheol Song, Republic of Korea

Dr. Jill Stanley, New Zealand

Dr. Palitha Weerakkody, Sri Lanka

National Editorial Advisory Board

Dr. S. D. Shikhamany

Dr. V. A. Parthasarathy

Dr. K. V. Peter

Dr. Sisir K. Mitra

Dr. S.K. Tikoo

Dr. Seetharam Annadana

Dr. A. Krishnamoorthy

Dr. Leela Sahijram

SOCIETY FOR PROMOTION OF HORTICULTURE (REGD.)

Email : sphiihr2005@gmail.com Website : www.sphindia.org

Executive Council - 2020

President : Dr. M.R. Dinesh

Vice Presidents : Dr. G. S. Prakash
Dr. T.N. Shivananda

General Secretary : Dr. C. Aswath

Editor-in-Chief : Dr. S. Sriram

Treasurer : Dr. D.V. Sudhakar Rao

Joint Secretaries : Dr. P.C. Tripathi
Dr. T.H. Singh

Members : Dr. T.S. Aghora
Dr. K.S. Shivashankara
Dr. Prakash Patil
Dr. H. S. Oberoi
Dr. C.K. Narayana
Dr. B. Narayanaswamy
Dr. B. Hemla Naik
Dr. L.N. Mahawer
Dr. Sanjay Kumar Singh
Dr. S.K. Mitra
Dr. S. Hazarika
Dr. Gobind Acharya

This Journal is abstracted in CABI, Current Contents, AGRIS, Indian Science Abstracts, Scopus, DOAJ and Redalyc. It is a participant of AmelICA.

Request for membership subscriptions along with cheque/DD drawn in favour of **Society for Promotion of Horticulture, Bengaluru** may be sent to General Secretary, Society for Promotion of Horticulture, Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bengaluru - 560 089, India. All members except student members and subscribers get all publications of SPH free of cost. Any correspondence other than editorial may be addressed to General Secretary, Society for Promotion of Horticulture, Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bengaluru - 560 089, India.

Advertising space in the journal is available. For information and rates, please contact General Secretary, SPH, IIHR, Bengaluru - 560 089, India. Advertising material

should cater to the interest of researchers, subscribers, etc. who are involved in promotion of horticulture. Publication of advertisement is not an endorsement or approval, expressed or implied by the SPH or the editors of any service, product or claim made by the manufacturer.

Coverpage Courtesy : **Rami Reddy P.V.**, P.No. 225

SUBSCRIPTION RATES

Patron	₹ 20,000
Life member	₹ 5,000
Annual Member	₹ 1,000 / US \$ 100 (US \$ 50 for SAARC countries)
Student Member	₹ 500
Student Life Member	₹ 3,000
Annual Subscription	₹ 1,500 / US \$ 100 (US \$ 60 for SAARC countries) (for institutions)
Enrolment Fee	₹ 200 / US \$ 5 (Additional for all types of Membership)

NAAS rating of this journal is 3.43. JHS is now available online.

Authors have to submit manuscripts using the link : <https://jhs.iihr.res.in/index.php/jhs>

Technical Assistance : **Dr. Sridhar Gutam, Thippeswamy S. and Pramida A.**

JOURNAL OF HORTICULTURAL SCIENCES

Volume 15

Number 2

December 2020

CONTENTS

In this Issue

i-ii

Review

- Biodiversity of tropical fruits and their conservation in India** 107-126
Sankaran M. and Dinesh M.R.
- An overview of canopy management in cashew (*Anacardium occidentale* L.)** 127-135
Adiga D.J., Veena G.L., Thondaiman V. and Babli M.

Original Research in Papers

- Phenotypic variability for horticultural and fruit quality attributes in plastic house grown tomato** 136-146
Adeniji O.T., Tenebe A.V., Ishaka A., Jandong E., Adamu J.T., Adekoya M., Zamzam M.A. and Aremu C.A
- Development and evaluation of novel gladiolus hybrid selections IHRG-7 (IC620379) and IHRG-11 (IC620380) for flower quality and *Fusarium* wilt resistance** 147-152
Rao T.M., Janakiram T., Negi S.S., Aswath C., Dhananjaya M.V., Kumar R. and Ramachandran N.
- Evaluation of potassium salt of phosphonic acid in Nagpur mandarin with special reference to *Phytophthora* management** 153-160
Ingle Y.V., Paithankar D.H., Sadawarte A.K. and Bhonde S.R.
- Genetic analysis in mango (*Mangifera indica* L.) based on fruit characteristics of 400 genotypes** 161-172
Sankaran M., Dinesh M.R., Gowda D.C.S. and Venugopalan R.
- Standardization of nitrogen application for potted *Chrysanthemum morifolium* cv. kikiobiory** 173-176
Tanya Thakur
- Influence of inorganic nutrients on growth, flowering and quality of *Dendrobium* cv. Singapore white** 177-182
Sujatha A. Nair, Sankar V., Muralidhara, B.M., Awcharae C.M. and Singh D.R.
- Palynological investigations in *Jasminum* spp.** 183-190
Ganga M., Lakshmi J., Manivannan N. and Rajamani K.



- Effect of putrescine and benzyl adenine on growth, flowering and post-harvest keeping quality parameters in chrysanthemum (*Chrysanthemum morifolium ramat*)** 191-196
Taranjit Singh and Madhu Bala
- Studies on bioavailability of iron from fe-fortified commercial edible mushroom *Hypsizygusulmarius* and standardization of its delivery system for human nutrition** 197-206
Pandey M., Gowda N.K.S., Satisha G.C., Azeez S., Chandrashekara C., Zamil M. and Roy T.K.
- Amino acid profile of eighteen isolate of different edible macrofungal species** 207-220
Azeez S., Pandey M., Jasmin M.R., Rachitha R., Satisha G.C., Roy T.K.
Chandrashekara C. and Shivashankara K.S.

Short Communications

- A promising new tamarind selection-lakshamana : Linking biodiversity with livelihood** 221-224
Kanupriya C., Karunakaran G. and Singh P.
- Mexican creeper, *Antigonon leptopus* Hook. and Arn : An effective bee forage plant to conserve honey bee** 225-228
Rami Reddy P.V.
- First report on honeydew excretion by the melon thrips, *Thrips palmi* karny (Thysanoptera : Thripidae) and its biochemical analysis** 229-232
Aravintharaj R., Asokan R. and Roy T.K.
- Influence of potting mixture on growth and economics of stone graft of mango cv. alphonso** 233-237
Lad O.A., Kulkarni M.M., Ragaji S.G., Gavankar M.S., Burondkar M.M., Gokhale N.B.
Pawar C.D., Khandekar R.G., Kshirsagar P.J. and Desai V.S.

In this issue...

Hearty New Year Greetings from our Editorial Team to all the readers of JHS!

As the world is slowly coming out of glitches of pandemic, there is no other better way than celebrating 2021 as Year of Fruits and Vegetables as announced by United Nations Assembly to welcome the new year and recognize the importance of nutrition for better health. Fruits and Vegetables ensure the Nutritional Security to humankind. They play key role in addressing the malnutrition that is a major concern. We are proud that JHS creatins awareness of importance of fruits and vegetables by publishing the recent developments in research with respect to these crops.

*Diversity of fruit crops and genetic resources available with respect to fruit crops are important for developing better fruit crop varieties. **Sankaran and Dinesh** have reviewed the “Biodiveristy of Fruit Crops in India” in a very comprehensive way. There is diversity in Jasmine species. **Ganga et al.** carried out the palynological investigations and recorded the variability in pollen morphology in different species of Jasmine by documenting images using scanning electron microscope. Biodiversity can be linked to livelihood also. One such success story with tamarind selection ‘Lakhamna’ is being reported by **Kanupriya et al.** This tamarind selection has been identified from participatory breeding programme. It has a better pod characters and more preferred by consumers.*

*Protected cultivation has seen greater momentum in last two decades. **Adeniji et al.** identified the best varieties of tomato for polyhouse cultivation in Nigeria. **Rao et al.** selected two gladiolus hybrid selections IIHRG-7 and IIHRG-11 with red purple and red coloured flowers respectively. These hybrids have resistance to Fusarium wilt and suitable for cut flower and flower arrangement purposes. **Sankaran et al.** analysed the variance for 6 quantitative and 30 qualitative traits in mango in 400 genotypes and identified 18 clusters. Selected genotypes from specific clusters can be used in hybridization programme.*

*The production aspects are important in perennial crops. It is crop management that needs to be prioritized for enhanced yield. **Adiga et al.** have reviewed the research work carried in “Canopy Management in Cashew”, providing the wholistic view of cultural operations to have a better crop. Use of soilless medium in nursery industry is gaining importance. Best suited potting mixture for mango stone graft of cv. Alphonso has been identified by **Lad et al.** They found that cocopeat + leaf manure + compost (1:1:2) as pot mixture provided better plant growth.*

*Growing Chrysanthemum in pots is practiced in home and terrace gardens. The cultivar Kikiobiory is well suited for this purpose. **Thakur** has studied the nitrogen requirement for this cultivar and has come out with the recommendation of 300 mg of N per pot applied*



twice in September and October in Punjab for best results. In another study, **Singh and Bala** confirmed that use of benzyl adenine at 200 ppm helped in extended vase life of *Chrysanthemum morifolium* flowers. **Nair et al.** recorded that foliar spray of 30:20:20 NPK at weekly interval recorded more number of flowers of *Dendrobium* cv. Singapore White with significantly longer spikes.

Crop production is directly influenced by pollinators. Decline in honey bee population is a serious concern and to conserve the pollinators community approach through ecosystem services is required. **Rami Reddy** reports the benefits of having ornamental plant Mexican Creeper (*Antigonon leptopus*) as forage plant. This creeper attracted all the four species of honey bees studied. This creeper can be used as bioindicator of honey bee population.

Aravindaraj et al. have reported the honey dew secretion by *Thrips palmi* and analysed the composition of it. They had identified different sugars present in the honey dew secretion of *Thrips*. *Thrips* not only cause direct damage but act as vectors of many plant viruses. Management of diseases in perennial crops is a challenge. *Phytophthora* incited root infection in citrus needs concerted efforts. **Ingle et al.** have demonstrated that use of potassium salt of phosphonic acid could help in management of *Phytophthora* root rot in Nagpur Mandarin.

Mushrooms can fill the gaps in nutritional security as they are rich in nutritive value. Iron deficiency is important issue to be addressed. Iron fortified oyster mushroom products have been developed by **Pandey et al.** The bioavailability of iron from Arka Mushroom Fe-Fortified Rasam Powder has been confirmed. In another study, the amino acid profile of 18 isolates of oyster mushroom species belonging to 4 species have been documented by **Azeez et al.** Quantification of essential and non-essential amino acids has been reported. Nutritionally superior isolates can be selected from these isolates.

The editorial team of JHS expresses the sincere efforts of reviewers who really complement the publication processes. All scientists and scholars can utilize the open access of JHS. Recently FAO has made JHS available through AGRIS. It is indexed by Redalyc, CABI_Hort and Scopus. All subscribers, scientists and scholars are requested to continue their support in publishing quality information in **Journal of Horticultural Sciences**.

S. Sriram
Editor in Chief

Original Research Paper

Studies on bioavailability of iron from iron fortified commercial edible mushroom *Hypsizygus ulmarius* and standardization of delivery system for human nutrition

Pandey M.^{*1}, Gowda N.K.S.⁴, Satisha G.C.², Azeez S.³, Chandrashekara C.¹, Zamil M.² and Roy T.³

¹ Division of Crop Protection, ²Division of Natural Resources

³ Division of Basic Sciences, Indian Institute of Horticultural Research, Hesaraghatta, Bengaluru, India

⁴ National Institute of Animal Nutrition and Physiology, Adugodi, Bengaluru, India

*Corresponding author Email: Meera.Pandey1@icar.gov.in

ABSTRACT

Iron is one of the most important micronutrients for human health. Iron deficient diet and defective iron absorption are among the major reasons for iron malnutrition. Mushrooms are edible fungi, which are a very good source of iron and can be easily grown on agricultural residues at home scale, dehydrated and stored in the powder form, which can supplement the daily diet. Although mushrooms in general and oyster mushrooms in particular are a rich source of iron, yet, have not become a recommended diet by the nutritionists due to lack of data pertaining to its bioavailability from mushrooms. Data has been generated in the present study on the *in vitro* bioavailability of iron from non-fortified and iron fortified *Hypsizygus ulmarius*, which is a commercially grown species. This is the first report pertaining to the bioavailability of iron from iron fortified mushrooms. A delivery system for human nutrition was also standardized in the form of Arka Mushroom Fortified Rasam Powder using iron fortified and non-fortified mushrooms, which can be used to mitigate iron malnutrition. The data generated in this study will help in providing application techniques to use mushrooms as dietary source of iron in everyday diet to mitigate iron deficiency.

Key words: Arka Mushroom Rasam, Bioavailability, *Hypsizygus ulmarius* and Iron fortified

INTRODUCTION

Iron is one of the most important micronutrients for human health. It is required in the process of oxygen transport and storage (as a constituent of hemoglobin and myoglobin), for electron transfer (in cytochromes), desaturation of fatty acids, tyrosine iodination (in thyroid peroxidase), prostaglandin synthesis, formation of erythrocytes, maintenance of heat balance, as well as humoral and cellular resistance (Burke *et al.*, 2014; Nadadur and Mudipalli, 2008). Iron deficiency can also result in decreased cognitive health. Nutrition Monitoring Bureau Survey (NNMBS, 2006) showed that anemia prevalence in adolescent girls in India was 69.7% which caused 1.8% loss of GDP. Indian government has been working towards reducing the load of anaemia in girls and women by 50%. Numerous

interventions like biofortification and fortification of diet are being undertaken to mitigate this problem. A very important intervention can be adoption and popularization of alternate foods rich in iron, which have not yet become a part of daily diet. Although mushrooms in general and oyster mushrooms in particular are a rich source of many minerals including iron, yet, have not become a recommended diet by the nutritionists due to lack of data pertaining to bioavailability. Any iron fortification method is incomplete until a proper delivery system is also standardized so that the benefit can reach the target population. The innovation for the process of production of iron fortified mushroom was taken to a logical and meaningful conclusion through the standardization of recipe for iron rich Arka Mushroom Rasam Powder which is available as licensed technology from ICAR-IIHR. The data



generated in this study will fill this gap on bioavailability of iron from mushrooms and provide application techniques to use mushrooms as dietary source of iron in everyday diet.

MATERIALS AND METHODS

Hypsizygus ulmarius (Elm oyster) was cultivated on sterilized paddy straw-based substrate. Paddy straw was cut into small pieces (about 2.5-3.0 cm) using a motorized chaff cutter and soaked in clean water for 4 h. The soaked straw was drained for removal of excess water and spread on meshed tray under sun to reduce the moisture content to 65-70%. One kg wet straw was filled in polypropylene (PP) bags measuring 304 x 406 mm, plugged with non-absorbent cotton and sterilized in autoclave at 121°C, 15lb pressure for 25 min. The moisture content of the wet straw was determined by gravimetric method by drying 100 g of wet straw to a constant weight in oven at 70°C. The sterilized bags were cooled to room temperature and spawned (seeded) using grain spawn of *H. ulmarius* @ 5% (50g spawn per bag) in an aseptic bag inoculation chamber. Spawned bags were incubated at 26±2°C for spawn running (vegetative growth). After 25 days of completion of spawn running, the bags were shifted to cropping room where a temperature of 26±2°C and humidity of 80-85% was maintained throughout the cropping period of 30 days. Four to five holes were made in each bag to induce mushroom initiation. Iron fortified mushroom was developed through an innovative technique using iron salt (Patent application No -201841022601). Mature mushrooms of both the non-fortified *H. ulmarius* (NFHU) and iron fortified *H. ulmarius* (IFHU) were harvested, hand shredded and dehydrated in a tray dryer covered with PP sheet at 48±2°C for 12-14 h. Dry mushrooms were powdered in wooden pestle mortar. Necessary precautions like using gloves while handling mushrooms, using plastic knife to cut mushrooms, were taken in handling of iron enriched mushrooms to avoid external contamination. NFHU powder and IFHU powder prepared was used for the *in vivo* iron bioavailability studies.

***In vivo* studies**

The experiment was conducted in the Laboratory Animal House at Experimental Livestock Unit, National Institute of Animal Nutrition and Physiology (NIANP), Bangalore, India. The animal experiment protocol was as approved by the Institutional Animal

Ethics Committee (IAEC) and carried out as per the guidelines of Committee for the Purpose of Control and Supervision of Experiments in Animals (CPCSEA), Ministry of Environment, Forests and Climate Change, Government of India.

Experimental design

Forty-eight numbers of healthy weaned (40 days old) Wistar strain (*Rattus norvegicus*) albino rats were randomly divided into four dietary groups, with four replicates of three rats in each group. All the rats were housed in polypropylene cages and maintained under similar managemental conditions. Temperature and humidity were maintained at 23±2°C and 50 to 70% respectively in proper ventilated animal house. All rats were fed pelleted feed and offered purified drinking water *ad libitum*.

Preparation of experimental diets

Commercial pelleted rat feed was served as the control diet (A). The control diet pellets were finely powdered and mixed well separately with dried NFHU and FHU mushroom powder in 75:25 proportion to prepare experimental diet B and C respectively. The experimental diet D comprised of control feed supplemented with ferrous sulfate (Analytical grade, SD fine chemicals, Mumbai, India) equivalent to the iron content of fortified mushroom. Diets B, C and D were again pelleted in a manual pelletizer, air dried for 24 hours, later oven dried at 60°C for 24 h and stored in respective air tight containers until fed to the experimental rats. Necessary precautions were taken at each step of diet preparation, pelletizing and storage to avoid contamination.

Sample collection

All the rats of different dietary groups were offered daily weighed amount of respective experimental diets in the morning at 9:30 h. They were provided with fresh and clean deionized water *ad libitum*. The feeding experiment was continued for 30 days and initial as well as final body weight of each rat was recorded. At the end of experiment, a digestibility trial of 5 days duration was conducted to record dry matter (DM) intake, DM digestibility and apparent Fe absorption in gut.

Analytical technique for iron bioavailability studies

The samples of diets offered, residues and faeces were analyzed for DM as per standard procedures of Association of Official Analytical Chemists

(AOAC 2000). The feeds and faeces were taken in pre-weighed silica crucibles and oven dried at 80°C for 24h, decarbonized and ashed at 550-600°C in a muffle furnace. The total ash was digested with 5N HCL over a hot plate for 15 min, cooled and filtered through Whatman filter paper (No. 41) into volumetric flasks of desired volume. Iron content in mushroom, diets and faeces was estimated using Optima 8000 inductively coupled plasma optical emission spectrophotometer (ICP-OES) (Perkin Elmer, Shelton CT#064840, USA).

Statistical analysis

The data were analyzed using the statistical Package for Social Sciences (SPSS, version 20.0 Chicago, USA) by one-way ANOVA and comparison of means was tested using Duncan's multiple range tests (Duncan 1995).

Standardization of delivery system for human nutrition

Standardization for a socially acceptable value-added product with iron fortified mushroom was undertaken. A local daily food accompaniment called 'Rasam or Soup' which is used in every Indian home of all strata was standardized using powder made by dehydration of both iron fortified and non-fortified mushrooms. The mushroom fortified rasam powder was made with traditional spices like coriander, jeera, pepper, fenugreek and tomato powder, chili powder, tur (*Cajanus cajan*) and urad dal (*Vigna mungo*), crushed garlic, salt and fresh coriander. This traditional mix was fortified with mushroom powder made from dehydrated non fortified and iron fortified Elm oyster mushroom. Ready to serve rasam or soup was made by boiling 10g of the rasam or soup powder in 350ml of water for 15 minutes and subjected to acceptance test by 501 people of rural and urban backgrounds. The rasam powder was also analyzed for its nutritive values. Taking a cue from the available literature on low availability of iron in human from legumes (Lynch et al 1984), the Arka rasam powder was also prepared with and without two legume ingredients (*C. cajan* and *V. mungo*). The five kinds of products viz. Rasam without legumes + iron fortified mushroom, Rasam with legumes + iron fortified mushroom, Rasam without legumes+ non fortified mushroom, Rasam with legumes+ non fortified mushroom and Rasam

without mushroom (traditional recipe) were analyzed for iron content as per the standard AOAC method.

Proximate nutritive analysis of Arka Rasam

The total soluble carbohydrates of the dry powders of rasam, rasam powder with mushroom and rasam powder containing iron rich mushroom was determined by the phenol-sulfuric acid method as described by Dubois *et al.* (1956). Protein content was calculated by multiplying the total nitrogen value by the factor 6.25. Fat percentage was determined in the samples by the Soxhlet procedure using petroleum ether (60-80 °C) (AOAC, 1984). Dietary fiber percentage was estimated in dry, defatted samples by the Official AOAC Method No. 982.29 (Prosky *et al.*, 1992). All the results were expressed as mg/100g dry weight or fresh weight basis. The extraction procedure of water-soluble vitamins from mushroom was followed as described earlier by Esteve *et al.* (2001). A composite sample of 2 g of freshly cultivated mushrooms was homogenized using a domestic blender with 5 ml of MilliQ water to which 10 mL of 0.1 N HCl was added and incubated in a water bath at 95–100°C for 30 min. The extract was then allowed to cool at room temperature and the pH was adjusted to 4.0–4.5 using 5 M sodium acetate solution. One ml of takadiastase enzyme solution, contain 200 mg/ml was added to this extract and incubated in a water bath at 45–50°C for 8 hrs. After the enzymatic hydrolysis, 1 mL of 50% trichloroacetic acid was added to the extract and incubated at 95–100°C in a water bath for 5 min. After cooling it was centrifuge at 10000 rpm for 10 min, supernatant was collected and made up the volume to 25ml with 0.1N HCl. To the 5 mL of the stock solution 300 µL of potassium ferricyanide (1% in 15% NaOH) was added and left to react for 10 min in the dark. The extract was then neutralized with 200µL of 15% ortho-phosphoric acid, filtered through a nylon membrane syringe disc filter with a pore size of 0.45 µm followed by 0.2 µm and 1µl was injected immediately into the UPLC-MS/MS.

Mineral analysis of Arka Rasam

Mushroom samples were dried in the oven at 70°C to constant weight and analyzed for mineral nutrients as per the methods described by Piper (1966). Nitrogen content was determined by Kjeldahl method (Kjeltek Auto-Analyzer, Gerhardt, Germany),

phosphorus was estimated calorimetrically by vanado molybdophosphate method and potassium content by flame photometer. Calcium, magnesium, iron and zinc were determined using atomic absorption spectrophotometer (AAS) by Wet digest method with HNO_3 and HClO_4 in 10:4 ratio.

Water soluble vitamins analysis

Reagents

The water-soluble vitamin standards biotin (B7), niacin (B3), pyridoxine (B6), pantothenic acid (B5), folic acid (B9), cyanocobalamin (B12), thiamine (B1) and riboflavin (B2) were from Sigma Chemical Co., USA. The standard vitamin solutions were prepared in 0.01 N HCl. The enzyme takadiastase was purchased from Fluka, Switzerland. The organic solvents used for mobile phase in liquid chromatography were of chromatographic/MS grade and all other reagents were of analytical grade. Milli-Q water (Millipore system) was used to prepare the mobile phases. All mobile phases were filtered through 0.45 μm pore size membranes before use.

Equipment

Freshly harvested mushroom samples were used for the validation of the methodology. An Acquity UPLC-H class coupled with TQD-MS/MS from

Waters, USA with ESI source was used for vitamin determinations; the system was equipped with a degasser, quaternary pump, automatic injection system (0–10 μL), with a diode array detector and a temperature control compartment for the analytical column. The detection system allowed for the simultaneous detection at various wavelengths and MRM for individual masses. The overall system was controlled by the Mass lynx software, which also administered the data collection and treatment system.

MS-MS Methodology

The methodology for estimation of vitamins by UPLC-MS/MS system was validated for linearity, quantification and detection limits, and repeatability; recoveries of standards were 85-95%. The most sensitive, Multiple Reaction Monitoring (MRM) mode was employed for detection. The optimal operational parameters of the MS/MS system, to select the most abundant mass-to-charge ratio (m/z), was determined for each vitamin by direct sample infusion, using positive ionization mode (ES^+): the precursors ion and the product ions, the MS/MS capillary voltage, extractor voltage and RF lens values were set at 3.2 kV, 4 V and 0.1 V respectively, the gas flow for de-solvation and cone was set at 550 and 50 L/h. The cone voltage and the collision energy for individual vitamins were optimized (Table 1).

Table 1. MRM of water-soluble vitamins standards

Water soluble vitamins	Formula/Mass	Parent m/z [M+H] ⁺	Cone Voltage	Daughters	Collision energy (CE)	Ion Mode
Thiamine (B1)	264	265.03	20	122.06(Q)	16	ES+
		265.03	20	144.02	14	ES+
Riboflavin (B2)	376	376.97	40	243.05	24	ES+
Niacin (B3)	123	123.9	34	80.523(Q)	20	ES+
		123.9	34	77.47	18	ES+
Pantothenic acid (B5)	219.03	220.01	28	202.21	12	ES+
		220.01	28	124.16	20	ES+
Pyridoxine (B6)	169	169.97	24	152.09(Q)	12	ES+
		169.97	24	134.04	20	ES+
Biotin (B7)	244	245.03	26	227.14	14	ES+
Folic acid (B9)	441	442.1	24	295.16	16	ES+
Cyanocobalamin (B12)	677	678.29	38	147.18	68	ES+

* “Q” taken as quantified ion.

LC and MS/MS conditions

The mobile phase consisted of the aqueous phase 0.1% formic acid in water (A) and organic phase acetonitrile (B). The initial gradient of 60% A and 40% B was held for 0.5 min; at 6.0 min the gradient was changed to 5% A and 95% B, held for 0.5 min, and a linear gradient then followed, and later 30% A and 70% B in 8.0 min, held for 0.5 min. The system was then returned to initial conditions at 14 min and held for 1 min for equilibrating before the next injection. The flow rate was 0.3 ml/min. The analytical column 2.1x50 mm UPLC BEH-C18 column (Waters) with 1.7 μm particle size, protected by a Vanguard BEH-C18 with 1.7 μm . guard column (Waters). The column temperature was maintained at 25 °C and sample injection volume was 1.0 μl . The vitamins eluted were monitored using a PDA detector and the UPLC column effluent was pumped directly without any split into the TQD-MS/MS (Waters, USA) system, optimized for the vitamins analysis.

Extraction of Water-Soluble Vitamin

The extraction procedure of water-soluble vitamins from mushroom was followed as described earlier by Esteve *et al.* (2001). Two gram of composite sample of freshly harvested mushrooms was homogenized in a domestic blender with 5 ml of MilliQ water, added 10 mL of 0.1 N HCl and incubated in a water bath at 95–100 °C for 30 min. The extract was cooled to room temperature and pH was adjusted to 4.0–4.5 with 5 M sodium acetate solution. 1 ml of takadiastase enzyme solution (200 $\mu\text{g}/\text{ml}$) was added to this extract and incubated in a water bath at 45–50 °C for 8 hrs. After the enzymatic hydrolysis, 1 mL of 50% trichloroacetic acid was added to the extract and incubated at 95–100 °C in a water bath for 5 min. After cooling, the extract was centrifuged at 10000 rpm for 10 min; supernatant was made up to 25 ml with 0.1 N HCl. To 5 mL of this stock solution 300 μL of potassium ferricyanide (1% in 15% NaOH) was added and reacted in the dark for 10 min. The extract was then neutralized with 200 μL of 15% *ortho*-phosphoric acid, filtered through a nylon membrane syringe disc filter (0.45 μm pore size), followed by 0.2 μm filter, 1 μl was injected immediately into the UPLC-MS/MS.

Statistical analysis

The statistical analysis was done by t-test using the Excel (2019) software. The significant differences

($p < 0.01$) were evaluated for the mineral and vitamin contents of each parameter and for each sample individually. The statistical analysis of iron content data of Arka Mushroom Rasam using iron fortified mushroom was analyzed using the software Agres.

RESULTS AND DISCUSSION

The data presented in Table 1 shows that the dry matter intake of the experimental animals reduced significantly with both NFHU (diet B) and IFHU mushroom diet (diet C) as compared to diet without mushroom (diets A & D) by 22.09, 21.40 and 10.46, 9.67% respectively. The reduction was more pronounced in NFHU diet (diet B) as compared to IFHU diet (diet C). Mushrooms are known to be low calorie, high fiber diets with high amounts of non-digestible polysaccharides as β glucans, which act as prebiotic. Such diets can generally reduce the dry matter intake and body weight gain as is revealed in the present study. Hence mushrooms have been recommended as a diet to fight obesity as well (Friedman, 2016). However, no significant reduction in the daily weight gain of the experimental animals was observed in the present investigation. The data in table 1 also reveals that the gut absorption of iron was significantly higher both for NFHU (Diet B) and IFHU (Diet C) as compared with diets A and D without mushrooms. The gut absorption of iron from NHFU diet was 36.30% higher as compared to non-mushroom diet A and 39.52% higher as compared to non-mushroom diet D. The iron absorption from FHU was 66.76% higher as compared to non-mushroom diet A and by 70.70% higher as compared to diet D. This observation is of immense importance as despite its abundance in nature, iron still remains one of the major factors of mineral deficiency leading to numerous health issues. An estimated 20% of maternal deaths are directly related to anemia and another 50% of maternal deaths are associated with it (Anand *et al.*, 2014). Regula *et al.* (2016) studied the effect of feeding iron deficient Wistar rats with cereal products enriched with 10 and 20% *Pleurotus ostreatus* powder. The authors reported a complete restitution of iron deficient rats in a short time. They also reported that iron bioavailability from cereal products enriched with mushroom powder was higher in comparison to standard reference Fe (II) gluconate. Such results further corroborate the

results obtained in the present investigation especially in reference to the higher bioavailability of iron from iron fortified mushrooms (IFHU) as

compared to non- fortified (NFHU) mushrooms which can become an important strategy to mitigate iron deficiency.

Table 2. Performance of rats fed diets with and without iron enriched mushroom

Group	Dry matter intake (g/r/d)	Fecal dry matter outgo (g/r/d)	Dry matter digestibility (%)	Average daily gain (g/r/d)	Iron intake (mg/r/d)	Fecal iron outgo (mg/r/d)	Gut absorption of Iron (%)
A control	17.20 ±0.13 ^a	3.54±0.10	79.66±0.54 ^a	2.09 ±0.22	8.90 ±0.06 ^a	7.74 ±0.06 ^a	13.00± 0.92 ^c
B (NFHU) non fortified mushroom	13.40±0.98 ^b	3.70±0.34	72.38±1.82 ^b	1.66± 0.25	5.76 ±0.42 ^c	4.74 ±0.35 ^b	17.72 ±0.59
C (FHU) iron fortified mushroom	15.40±1.07 ^{ab}	4.23±0.41	72.62 ±1.48 ^b	1.53 ±0.07	6.80± 0.47 ^b	5.30 ±0.31 ^b	21.68 ±1.30 ^a
D equivalent of FeSo4	17.05±0.27 ^a	3.68±0.15	78.42 ±1.00 ^a	2.10± 0.04	9.38± 0.15 ^a	8.18 ±0.11 ^a	12.70 ±0.67 ^c
SEM+ ₋	0.51	0.14	1.03	0.10	0.41	0.40	1.04
F value	5.709	1.12	8.510	2.813	27.646	50.033	21.661
P value	0.012	0.379	0.003	0.084	0.000	0.000	0.000

The delivery system developed in the form of Arka Mushroom fortified Rasam or Soup powder was found to be highly acceptable during the acceptance trials. As evident from Table 2, 91.21% of the respondents liked and accepted the product Arka Mushroom fortified Rasam or Soup while 68.39% liked it very much. None of the respondents disliked the product. This was an important observation as developing iron fortified value-added products acceptable to masses is very challenging due to the color and flavor changes imparted by iron and the protection of the fortified iron from iron inhibitors generally present in cereal-based

foods (Hurrell 1997). Both these issues were resolved in the present delivery system through choosing of the right kind of traditional recipe. The traditional recipe of rasam or soup is originally reddish brown and the addition of iron fortified mushroom does not bring about any color changes and is well accepted. The recipe of rasam includes many traditional Indian spices with very strong aroma and flavor which masks any changes brought about in flavor due to the addition of iron fortified mushroom to the recipe.

Table 3. Acceptance score of Arka Mushroom Rasam or Soup by 501 panelists

9-point Hedonic scale	Acceptance among panelists (Nos)	Acceptance score (%)
9 Like extremely	52	10.379
8 Like very much	291	58.08
7 Like moderately	114	22.75
6 Like slightly	32	6.38
5 Neither like nor dislike	7	1.39
4 Dislike slightly	5	0.998
3 Dislike moderately	NIL	-
2 Dislike very much	NIL	-
1 Dislike extremely	NIL	-
Total	501	100

The traditional recipe has legumes like tur dal (*Cajanus cajan*) and urad dal (*Vigna mungo*), which contain phytates known to inhibit iron absorption. The recipe was suitably modified without the usage of the legumes to enhance iron bioavailability. The proximate and vitamin B composition of the standardized raw Arka Mushroom Rasam powder with and without mushrooms has been tabulated in table 3a & 3b. As

per the data in table 3a, the protein content of the mushroom fortified rasam powder significantly increased by 3.4%, carbohydrate by 2.14% and fiber by 1.43%. The fat content of the sample with mushroom significantly decreased by 3.53%. There was appreciable and significant increase in vitamin B content in the rasam powder with mushroom vis-à-vis without mushroom. (Table 3b).

Table 3a. Comparative proximate composition of raw Rasam or Soup powder

Composition	Raw Arka Mushroom Rasam powder with Mushroom mg/100g raw powder	Raw Rasam powder without mushroom mg/100g raw powder	't' Stat value
Protein	20.85 g	17.45 g	9.89**
Carbohydrate	27.64 g	25.5 g	27.74**
Fat	94.93 g	13.02 g	86.90**
Fiber	10.63 g	92.08 g	5.77**

** Significant at 1%

Table 3b. Comparative vitamin B composition of raw Arka Mushroom rasam or Soup powder

Composition	Raw Arka Mushroom Rasam powder with Mushroom mg/100g raw powder	Raw Rasam powder without mushroom mg/100g raw powder	't' Stat value
Biotin	2.576	0.857	11.15**
Cyanocobalamin	0.003	0	59.00**
Folic acid	0.059	0.003	8476**
Niacin	161.00	39.82	33.61**
Pantothenic acid	270.433	15.09	468.56**
Pyridoxin	8.104	1.829	77.35**
Riboflavin	9.057	1.62	83.90**
Thiamine	2.576	0.396	113.43**

** Significant at 1%

Table 4a & 4b depicts the data obtained of the proximate and vitamin B content of the cooked and ready to use rasam. There was a significant increase in the protein content of cooked and ready to use rasam with mushroom by 7.34%, carbohydrate by 16.19% and fiber by 3.15%. The

fat content in the product with mushroom significantly decreased by 7.3%. There was significant increase in the biotin, niacin, pantothenic acid, riboflavin and thiamin content in the cooked product with mushroom as compared to without mushroom.

Table 4. Comparative proximate composition of cooked Arka Mushroom rasam or Soup

Composition obtained after boiling 10g of the powder in 350ml of water	Cooked & ready to use Arka Mushroom Rasam with Mushroom mg/100ml	Cooked & ready to use Rasam without Mushroom mg/100ml	‘t’ test value
Protein	3.19 g	2.46 g	64.90**
Carbohydrate	8.32 g	6.70 g	105.76**
Fat	0.1 g	1.42 g	29.135**
Fiber	4.31 g	3.99 g	40.93**

** Significant at 1%

Table 4b. Comparative vitamin B composition of cooked Arka Mushroom rasam or Soup

Vitamin B	Cooked & ready to use Arka Mushroom Rasam with Mushroom µg/100ml	Cooked & ready to use Rasam without Mushroom µg/100ml	‘t’ test value
Biotin	2.555	0.316	7.88**
Cyanocobalamin	0	0	65535 ^{NS}
Folic acid	0	0.0035	65535 ^{NS}
Niacin	134.184	34.914	151.18**
Pantothenic acid	49.905	14.50	114.89**
Pyridoxin	4.628	0.677	27.80**
Riboflavin	3.404	0.354	272.92**
Thiamine	2.594	0.268	59.94**

** Significant at 1%

NS – Non significant

Table 5 shows the macro and micro element composition of the Arka rasam powder with and without mushroom. As seen from table 5a, there was a significant increase in Phosphorus, potassium and calcium levels in the rasam powder with mushroom as compared to without mushroom. The levels of other minerals remained same in both

treatments. As per table 5b, the level of micronutrients Manganese significantly increased in the product with mushroom as compared to without mushroom. The level of Zinc decreased significantly by the addition of mushroom. The levels of iron and copper remained statistically unchanged in both treatments.

Table 5a. Comparative macro and micro element composition of raw Arka Mushroom rasam or Soup powder

Composition	Raw Arka Mushroom Rasam powder with Mushroom mg/100g raw powder	Raw Rasam powder without mushroom mg/100g raw powder	't' test value
Macro-elements			
Phosphorus	516.66	341	10.106**
Potassium	2170	1943.33	6.552**
Calcium	348.33	451	19.479**
Magnesium	120	131.33	4.25 ^{NS}
Micro-elements			
Iron	9.23	13.52	4.066 ^{NS}
Manganese	5.43	2.01	80.781**
Copper	1.11	1.05	1.889 ^{NS}
Zinc	1.20	3.20	22.863**

** Significant at 1% NS – Non significant

Lynch *et al.* (1984) reported a low bioavailability of 0.84-1.91% of iron from legumes in human beings.

The data in table 6 shows the iron content of the Arka Rasam Powder made without the use of legumes.

Table. 6. Iron analysis of rasam with iron fortified mushroom

Sl. No.	Treatment	Iron content(mg/100g dry powder)
1	Rasam without dal + with iron fortified mushroom	30.8 ^a
2	Rasam with dal + iron fortified mushroom	29.2 ^a
3	Rasam without dal+ non fortified mushroom	11.36 ^b
4	Rasam with dal+ non fortified mushroom	12.06 ^b
5	Rasam without mushroom (traditional recipe)	11.83 ^b
	Coefficient of variation	8.589
	CD (0.01)	4.234
	CD (0.05)	2.977

As evident from Table 6, the iron content of the product Arka Rasam Powder made with and without legumes but with iron fortified mushroom was statistically at par and significantly higher as compared to the one made with non-fortified mushroom or the traditional rasam without mushroom. The iron content of the rasam with and without non fortified mushroom and without dal were statistically at par. This result is of significance as the bioavailability of iron is higher as depicted in the animal model studies earlier. Bioavailable iron in 10g Arka rasam powder made with iron fortified mushrooms @ 21.68% bioavailability is 0.667m and it takes cae of 28.02% requirement of bioavailable iron (2.3 mg/day) iron measuring women.

Conclusion

The high bioavailability of iron from both non fortified and iron fortified *Hypsizyguis ulmarius* is of immense significance as the predominantly vegetarian population of India suffer with iron deficiency due to very poor bioavailability of 5-8% of non-heme iron from plant sources. The bioavailability of heme iron from meat sources varies from 18-25%. The present data of the bioavailability of 21.68% from iron fortified mushroom can play a significant role in mitigating iron malnutrition in the country. The delivery system standardized in the form of Arka Mushroom Rasam powder was not only significantly higher in

iron content with high bioavailability but also showed enhanced nutritional values in terms of protein, carbohydrate, fiber and B vitamins like biotin, niacin, pantothenic acid, riboflavin and thiamin. Thus, this

product can play an important role in mitigating iron malnutrition as it fits into the taste, flavor and aroma of an already well accepted form of daily food accompaniment already in vogue.

REFERENCES

- Anand, T., Rahi, M., Sharma, P. and Ingle G. K. 2014. Issues in prevention of iron deficiency anaemia in India. *Nutrition*, 30: (7-8), 764-770
- AOAC 1984. Association of Official Analytical Chemists. 14th ed. Official Methods of Analysis: Washington DC. USA.
- AOAC 2000. Association of Official Analytical Chemists. 17th ed. Official Methods of Analysis: Washington DC. USA.
- Burke, R.M., Leon, S.L. and Suchdev P.S. 2014. Identification, prevention and treatment of Iron deficiency during the first 1000 days. *Nutrients*, 6(10): 4093-4224
- DuBois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A. and Smith, F. 1956. Colorimetric method for determination of sugars and related substances. *Analytical Chemistry*. 28 (3), 350-356 DOI: 10.1021/ac60111a017
- Duncan, D.B .1995. Multiple range and multiple F-tests. *Biometrics* II. pp-1-42.
- Esteve, M.J., Farre, R., Frygola, A., & Garcia-Cantabella, J.M. (2001). Simultaneous determination of thiamine and riboflavin in mushrooms by liquid chromatography. *J Agrl. Food Chem.*, 49: 1450–1454.
- Hurrell, R.F., 1997. Preventing iron deficiency through food fortification. *Nutrition Reviews*. 55(6): 210-222
- Lynch S.R., Beard, J. L., Dassenko, S.A. and Cook, J.D. 1984. Iron absorption from legumes in humans. *American J Clinical Nutrition*. 40 (1):42-47
- Friedman, M. 2016. Mushroom polysaccharides: chemistry and anti-obesity, antidiabetic, anticancer, and antibiotic Properties in cells, rodents, and humans. *Foods*, 5(4): 80; <https://doi.org/10.3390/foods5040080>
- Nadadur, S. and Mudipalli, A. 2008. Iron transport and homeostasis mechanisms: Their role in health and disease. *Indian J Med. Res.* 128: 533-544.
- National Nutrition Monitoring Bureau, Diet and Nutritional Status of Rural Population and Prevalence of Hypertension among adults in Rural areas, NNMB 2006. Technical Report No. 24. Hyderabad: NNMB, NIN
- Piper, C. S. 1966. *Soil and Plant Analysis*. Hans Publishers, Bombay, p. 368.
- Prosky, P., Asp, N. G., Schweizer, T. F., De Vries, J. W. and Furda, I. (1992). Determination of insoluble and soluble dietary fiber in foods and food products: Collaborative study. *Journal of Association of Official Analytical Chemists* 75: 360-367.
- Regula, J., Krejpcio, Z., Staniek, H. 2016. Iron bioavailability from cereal products enriched with *Pleurotus ostreatus* mushrooms in rats with induced anaemia. *Annal. Agri. Environ. Med.* 23(2): 310-314. doi: 10.5604/12321966.1203896.
- Manuel otivares MD; Fernodo piz arro MT; Tomas walter MD; miguel arredondo MT; Msci Eva Hertrompf MDI Msci 1999.
- Bioavailability of iron supplements consumed dialy is not different from that of iron supplements consumed weekly. *Nutrition Research, Vol 19, issue 2*, pp 179-190.

(Received on 01.05.2020 and Accepted on 4.11.2020)

INFORMATION TO CONTRIBUTORS

Journal of Horticultural Sciences, an international journal, is the official publication of **Society for Promotion of Horticulture (SPH)**. It covers basic and applied aspect of original research on all branches of horticulture and other cognate disciplines, which promotes horticulture in its broadest sense. Its goals are to apprise horticultural scientists and others interested in horticulture of scientific and industrial developments and extension findings. The area of research include evaluation of germplasm, breeding, agronomic practices, physiology, biochemistry, biotechnology, soils and plant nutrition, plant protection, weed control, pesticide residue, post harvest technology, economics, extension, farm machinery and mechanization, etc. which facilitate in the growth and expansion of horticulture. The journal is published twice a year, in June and December.

The Journal of Horticultural Sciences (JHS) publishes critical reviews, research papers and short communications. Three copies of the manuscript and an electronic form (CD, MS Word) should be submitted to the Chief Editor, JHS, SPH, Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bangalore-560 089. The manuscript should preferably pertain to the research work carried out during the last five years. Author(s) must certify that the manuscript (s) has/have not been sent elsewhere for publication. All the authors have to become the members of SPH when a paper is accepted for publication. All papers will be refereed. Short communications on significant research findings, new record / technology are welcome. Besides invited review papers, scientists with vast experience on a particular field of research can also submit review papers which will be refereed. Decision of the Chief Editor / Editorial board is final. Authors are permitted to photocopy their article for non-commercial and scientific purpose. No reprints shall be provided *gratis*. Acceptance of manuscript for publication in JHS shall automatically mean transfer of copyright to the SPH. The chief editor/ Editorial board assumes no responsibility for the statements, opinion or facts expressed in the journal, which rests entirely with the author(s) thereof. Mention of a pesticide or a commercial or proprietary product does not constitute an endorsement or recommendation for the use.

Title: The title of the article should be bold and in running form. Use the font Times New Roman (14 point). Botanical / scientific names should be italicized. Author name(s) should be in running and bold with full address of the first author including e-mail address (it is mandatory as future correspondence will be only through e-mail). The address of other author(s), if different from the first author, should be given as footnotes and indicated by consecutive superscript numbers. A brief running title should be provided on a separate sheet.

Abstract: The abstract should not exceed 200 words. It should be suitable for indexing and publication in abstracting journal. Very pertinent keywords may be furnished.

Text: The text should be typed in double space on one side of good quality paper (21 x 29 cm) with 3cm margin on all sides **without justifying the text** and in clear and concise English. Use the font Times New Roman (12 point). The paper should be divided into subheadings (placed on the left margin and in upper case) such as Introduction, Material and Methods, Results and Discussion, Acknowledgements, and References. Units and abbreviations should be in metric (SI) system. It is desirable that authors take due care on clarity and brevity of the paper. The length of the paper should not exceed 2500 words.

Tables/ Illustrations/ Photographs: Each table should be on a separate sheet with a short title at the end of the paper, numbered in the order in which it appears in the text. The data reported must be subjected to appropriate statistical analysis. The illustrations should be relevant to the research findings and should not be repeating of data presented in the table. Only very good photographs, mounted on hard paper to avoid folding, given on a separate sheet of paper with title, which are reproducible, will be accepted. Data to be presented in graphical form should be sent on quality glossy contrast paper without folding.

References: References should be cited in the text in the form of (Anon., 1999; Prakash, 2002; Krishnamoorthy and Mani, 2004). The term *et al* should be used when there are more than two authors. The letters, a,b,c,... should be used following the year, to distinguish between two or more papers by the same author(s) in one year. References at the end of the text should be given in the following form:

Shikhamany, S. D. and Satyanarayana, G. 1973. A study on the association of leaf nutrient contents with poor yields in Anab. E.shahi grape (*Vitis vinifera* L.). *Ind. J. Hort.*, **30**: 376 - 380

Panse, V. G. and Sukhatme, P. V. 1978. Statistical methods for Agricultural workers. ICAR, New Delhi, p 108.

Srinivas, K. 1987. Response of watermelon (*Citrullus lanatus* Thunb. Musf) to drip and furrow irrigation under different nitrogen and plant population levels. Ph.D thesis, UAS, Bangalore

Mehta, N. K. and Sharma, S. D. 1986. Studies on flowering and fruit retention in some cultivars of peach (*Prunus persica* Batch). In: Advances in Research on Temperate Fruits. *Proc. Nat'l. Symp. Temp. Fruits*, Solan (India), Dr. Y. S. Parmar Univ. Hort. and Forestry, pp 37-42

Krishnamoorthy, A. and Mani, M. 2000. Biological Control of Pests of Vegetable Crops.p367-78. In: Biocontrol Potential and its exploitation in sustainable Agriculture. Vol. 2: Insect Pests. Upadhyay, R. K. Mukerji, K. G. and Chamola, B.P. (ed.). Kluwer Academic / Plenum Publishers, New York

Cover photo (s) shall be included at the discretion of Editor. Authors may submit photographs/figures/diagrams for cover page while submitting the manuscript.

AUTHOR INDEX - VOL. 15 (1&2) 2020

Name	Page	Name	Page
A			
Adamu, J.T.	136	Gavankar, M. S.	233
Adekoya, M.	136	Gokhale, N. B.	233
Adeniji, O.T.	136	Gowda D. C. S.	161
Aghora T.S.	62	Gowda, N. K. S.	197
Ahamed N.	17	I	
Aravintharaj, R.	229	Ingle Y. V.	153
Aremu, C.A.	136	Ishaka, A.	136
Ashok Kumar J.	45	J	
Asokan, R.	229	Jadhav S.B.	67
Aswath C.	93	Janakiram, T.	147
Aswath, C.	147	Jandong, E.	136
Awcharae, C. M.	177	Jasmin M. R.	207
Azeez, S.	197, 207	Jessy Mol K.K.	52
B			
Babli, M.	127	K	
Bala, M.	191	Kalaivanan D.	9
Bhatt R.M.	62	Kanupriya, C.	221
Bhonde, S. R.	153	Karunakaran, G.	221
Burondkar, M. M.	233	Katwate S.M.	67
C			
Chandran, N. K.	81	Khandekar, R. G.	233
Chandrashekara C.	197, 207	Kshirsagar, P. J.	233
D			
Desai, V. S.	233	Kulkarni, M. M.	233
Dhananjaya, M. V.	147	Kumar D.	17
Dinakara Adiga, J.	127	Kumar, R.	147
Dinesh, M. R.	107, 161	L	
G			
Ganeshamurthy A.N.	9	Lad, O. A.	233
Ganga, M.	183	Lakshmana Reddy D.C	52
		Lakshmi, J.	183
		Laxman R.H.	35
		M	
		Madhavi Reddy K	52
		Manivannan, N.	183
		Manjunath B.L.,	35



Name	Page	Name	Page
Manoj Y.B.	52	Sankar V	177
Meena H.R.	72	Sankaran, M.	107, 161
Mohan N.	62	Satisha G.C.	197, 207
Muralidhara, B. M	177	Shejal A. Porob	97
N		Shilpa Pandurangaiah,	27
Nair A.K.	35	Shivashankar K.S.	27
Negi, S. S.	147	Shivashankara, K. S.	207
P		Singh D. R.	177
Paithankar, D. H.	153	Singh S.R.	17
Pandey, M.	197, 207	Singh, P.	221
Pawar, C. D.	233	Singh, T.	191
Priya Devi S	45, 97	Somasundaram J.	72
R		Sriram S.	81
Rachitha R.	207	Srivastava K.K.	17
Radha T.K.	72	Sudhakar Rao D.V.	27
Ragaji, S. G.	233	Sujatha A. Nair	177
Raghu B.R.	1	Susmita C.	62
Raghupathi H.B.	9	T	
Rajamani, K.	183	Tanya Thakur	173
Rajiv Kumar	93	Tejaswini Prakash	81
Ramachandran, N.	147	Tenebe, A.V.	136
Ramachandrudu K	45	Thangam M	45, 97
Rami Reddy, P. V.	225	Thondaiman, V.	127
Rao, T. M.,	147	V	
Rashmi I.	72	Veena, G.L.	127
Ravishankar K.V	27	Venugopalan, R.	161
Roy, T. K.	197, 207, 229	Vichare S.V	67
Rupa T.R	9	Y	
S		Yousuf S.	17
Sadashiva A.T.	27	Z	
Sadawarte, A. K.	153	Zamil, M.	207
Safeena S.A.	45	Zamzam, M.A.	136

SUBJECT INDEX - VOL. 15 (1&2) 2020

Name	Page	Name	Page
A		Foot rot	152
Alphonso	233	Free amino acids	207
Amino acid score	207	Fruit development	97
Antigonon	225	Fruit trees	9
Anti-senescence compound	191	Fruit quality	136
Apis spp	225	Fruit shape	136
Arka Mushroom Rasam	197	Fruit yield	136
B		Fruits	107
B:C ratio	233	Fusarium wilt	147
Bee flora	225	G	
Bioavailability	197	Garden pea	62
Biplot analysis	161	GCV	161
Bound amino acids	207	Genetic diversity	17
Breeding	62	Genetic analysis	161
Bulb	67	Genetic divergence	45
C		Genotype by environment	136
Canopy management	127	Gerbera	93
Carotene	27	Germplasm	1, 107
Carotenoid	27	GIS	107
CGMS	52	Gladiolus	147
Character correlation	136	Goa	97
Chrysanthemum	173, 191	Groundwater depletion	9
Conservation	107	Growth	67
Copper	72	Growth parameters	233
Correlation coefficient	45	Gummosis	152
Curry leaves	1	H	
Cut flower production	177	Heritability	161
Cut-flower	93	High temperature	62
D		Honey bees	225
Delayed flowering	191	Honeydew	229
Dendrobium	177	Hot pepper	52
Distribution	1	Hybrid	67
Diversity	1	Hypsizygos ulmarius	197
Drought	9	I	
E		In situ	107
Early summer	62	Iron	72
Evaluation	93, 147	Iron fortified	197
Ex situ	107	J	
F		Jasminum spp	183
Flower	67	K	
Flowering	147	Kikiobiory	173



Name	Page	Name	Page
L			
LC-MS-MS	229	Pruning	127
Leaf analysis	72	Pulp recovery	221
Lycopene	27	Q	
M			
Manganese	72	Quality	177
Mango	161, 233	Quantitative character	45
Marker Assisted Selection	52	R	
Micronutrient deficiency	72	Resistance Gene Analogues (RGA)	81
Mitochondria	52	Rootstocks	127
Morphotypes	1	Rose	81
Mushrooms	197	S	
N			
Nagpur mandarin	152	Sapota	72
Nitrogen	173	Scheduling irrigation	35
Novel hybrids	93	Selection	221
Nucleotide Binding Site-Leucine	81	Single linkage cluster analysis	17
Rich Repeats (NBS-LRR)		Single type tuberose	67
Nutrients	177	Soil volume wetting	35
Nutrition	207	Soilless media	233
O			
Onion	17	Solanum lycopersicum	136
Orchid	177	Spacing	35
ORF	52	Standardization	173
Ornamental creeper	225	Stress tolerance	62
P			
Palynology	183	Sugars	229
Papaya yield	35	T	
PBZ	127	Tamarind	221
PCV	161	Thrips palmi	229
Peak water	9	Tomato	27
Perennial crops	9	Training	127
Phytophthora	152	Tropical	107
Pink types	97	V	
Planting geometry	127	Variability	136
Podosphaera pannosa	81	Varieties	107
Policy issue	9	Vase life	147, 191
Pollen germination	183	Vegetable cowpea	45
Pollen morphology	183	W	
Polyhouse	93, 136	Water use efficiency	35
Potassium salt of phosphonic acid (PSPA)	152	Wax apple	97
Potted plants	173	White types	97
Powdery mildew	81	Wild species	107
Principal component analysis	17	Y	
		Yield	221
		Z	
		Zinc	72



**STATEMENT OF OWNERSHIP AND OTHER PARTICULARS ABOUT
JOURNAL OF HORTICULTURAL SCIENCES**

(Form IV)

Place of Publication : Bengaluru

Periodicity of publication : Half-yearly

Printer's Name : Mr. Ravikumar, B.A.

Nationality : Indian

Address : Resolution Print Media
#131, 6th Main, Meenakshinagar
Kamakshipalya, Bengaluru - 560 079.

Publisher's Name : Society for Promotion of Horticulture

Address : ICAR-Indian Institute of Horticultural Research
Hessaraghatta Lake P.O.
Bengaluru - 560 089

Editor-in-Chief : Dr. S. Sriram

Nationality : Indian

Address : ICAR-Indian Institute of Horticultural Research
Hessaraghatta Lake P.O.
Bengaluru - 560 089.

Name and addresses of individuals who own the journal and partners or are shareholders holding more than one per cent of the total capital : Society for Promotion of Horticulture
ICAR-Indian Institute of Horticultural Research
Hessaraghatta Lake P.O.
Bengaluru - 560 089.

I, Dr. S. Sriram, hereby declare that the particulars given above are true to the best of my knowledge and belief.

June 30, 2020

Sd/-
(S. Sriram)
Editor-in-Chief



SOCIETY FOR PROMOTION OF HORTICULTURE

ICAR-Indian Institute of Horticultural Research
Hessaraghatta Lake Post, Bengaluru-560 089, India
sphiihr2005@gmail.com / chiefeditor.jhs@gmail.com
Website : <https://sphindia.org>

ENROLMENT FORM

Name in full (in block letters) :
Dr./Mrs./Mr./Ms.

Designation :

Address for communication :

Phone No. :

E-mail ID :

Type of membership : Patron / Life member / Annual member / Student member*

Payment :

Demand Draft No. / Date :

Bank :

Place :

Date : SIGNATURE

Membership fee structure :

Type of membership	Membership amount	Enrolment fee	Total membership amount payable by Demand Draft (₹)
Patron	20,000/-	200/-	20,200/-
Life Member	5,000/-	200/-	5,200/-
Annual Member (India)	1,000/-	200/-	1,200/-
i. For SAARC authors	US \$ 100	US \$ 5	US \$ 105
ii. For SAARC countries	US \$ 50	US \$ 5	US \$ 55
Student member*	500/-	200/-	700/-

*The application of student members must be certified by their Head of dept. or equivalent and the student member shall not receive a copy of the journal.

Please send the duly filled-in enrolment form along with Demand Draft drawn in favour of Society for Promotion of Horticulture, by post to General Secretary, Society for Promotion of Horticulture ICAR-Indian Institute of Horticultural Research, Hessaraghatta Lake Post, Bengaluru - 560 089.

ACKNOWLEDGEMENTS

The editorial team acknowledges the services of the following reviewers

Dr. Shylesha A.N.

Principal Scientist, ICAR-NBAIR, Bengaluru

Dr. Ashwath Narayan

Associate Professor, UAS, Raichur

Dr. Mohan C.

Principal Scientist, ICAR-CTCRI, Trivandrum

Dr. Chavalli Sarada

Associate Professor, YSRHU, Guntur

Dr. Dinesh R.

Principal Scientist, ICAR-IISR, Calicut

Dr. Kalaivanan D.

Scientist, ICAR-IIHR, Bengaluru

Dr. Sudhakar Rao D.V.

Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Fakrudin B.

Professor, College of Horticulture, UHS, Bengaluru

Dr. Hebbar K.B.

Principal Scientist, ICAR-CPCRI, Kasaragod

Dr. Hima Bindu

Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Satisha J.

Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Krishnamurthy K.S.

Principal Scientist, ICAR-CPCRI, Kasaragod

Dr. Kundan Kishore

Principal Scientist, CHES (ICAR-IIHR), Bhubaneswar

Dr. Sankaran M.

Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Madhu Bala

Associate Professor, PAU, Ludhiana

Dr. Nandeesh P.

Senior Scientist, ICAR-IIHR, Bengaluru



Dr. Venkatarami Reddy P.
Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Prakash Tripathi
Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Prasad R.D.
Principal Scientist, ICAR-IIOR, Hyderabad

Dr. Rajashekar P.E.
Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Rajiv Kumar
Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Ravindran Chandran
Horticulturist, TNAU, Coimbatore

Dr. Ramani S.
Former Project Coordinator, AICRP on Honey Bees and Pollinator,
Bengaluru

Dr. Veena S.S.
Principal Scientist, ICAR-CTCRI, Trivandrum

Dr. Smaranika Mishra
Scientist, ICAR-IIHR, Bengaluru

Dr. Sujatha A. Nair
Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Tejaswini Prakash
Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Usha Bharathi T.
Scientist, ICAR-IIHR, Bengaluru

Dr. Sridhar V.
Principal Scientist, ICAR-IIHR, Bengaluru

Dr. Srinivasan V.
Principal Scientist, ICAR-IISR, Calicut

Sd/-
(S. Sriram)
Editor-in-Chief

New Varieties/ Technologies of ICAR-IIHR



New Water Melon - Arka Shyama variety



Arka Red - New Gerbera variety



Leaf curl resistant chilli varieties Arka Tejaswi, Arka Saanvi and Arka Tanvi



Arka Abhi



Arka Shuba

New Varieties/ Technologies of ICAR-IIHR



Arka Herbiwash - Safe way of removing pesticide residues



Arka Bharath - New teasel gourd variety

Journal of Horticultural Sciences is indexed by the following abstracting and indexing services



Article published in Journal of Horticultural Sciences are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

